# PINAX

Rerum Naturalium
BRITANNICARUM,

CONTINENS

VEGETABILIA, ANIMALIA,

ET

Fossilia,

In hac Infula reperta inehoatus.

Editio Secunda.

AUTHORE

Christophoro Merrett,

Medicinæ Doctore utriusque Societatis Regiæ Socio primoque Musæi Harveani Custode.

Μή τῷ λόγφ μουνον, ἀλλὰ τρος δοι νομίζεως τὰς ἐήτρας. Hipp.

Typis T. Roycroft, Impensis Cave Pulleyn, Prostat apud Sam. Thomson in vico vulgo dicto Duction: 1667.



# Imprimatur,

The Grigg, R.P. D. Humfr. Epifc. Lond. Sacell. Do-mest.





\*\*\*

#### Ornatissimo Viro

#### BALDUINO HAMET

Medicinæ Doctori, Collegii Med. Lond. Socio, Electori, Thesaurario, Singularique Benefactori.



IR Clarifsime biftoriæ naturalis, medicaque materiam tradituro mos sæculi sequendus,no.

meng; aliquod illustre buic libellulo præfigendum fuit. Cumq; bæc res ad Medicos pracipue spectet, seligendus erat aliquis ex Collegis (quorum plures sunt) qui communi litteratorum consensu utrama; calleret, insuperque necessitudine haud vulgari mihi conjunctus, Collegis singulis charus, propriisque

isque dotibus clarus esset. Et quo majus est, insignem te præstitisti erg Collegium nostrum, & summe pium erga templa publica Benefactorem. Praterea quod maximi fuit apud me mo. menti, & ad reipublica literaria utilitatem, ut modestiæ tuæ vim inferrem, & amice cogerem ad edendum ea, qua penes te sunt Adversaria dociissima; qua summo judicio, & trutina pertexuisti. Hac sunt amicissime Hamae qua me impulerunt, ut nomen tuum buic opusculo inscriberem. Is enim es quem carmine propemptico juvenem adhuc celebrare dignatus est Egregius poeta, & Philosophus Caspar Barlæus quem nuncupat----Battavicæ nomina prima Scholæ

F

Cui fua Phillyrides, Amythoniufque Melampus, Cui

Cui sua Pergameus credidit Ausa Senex.

Sacraque præclarus pandit confulta Machaon,

g 7

£.

0.

į.

7.

æ

Et veteres Graium terra reclusit opes.

Hac Encomia per totum vita curriculum ornare sategisti. Quantum in Medicina valueris, testantur viri quam plurimi nobiles, & illustrisima civium familie a te curate, & quanta apud ipsos fueris auctoritate innotescit omnibus. Adeo ut ad hanc artem non facius, sed doctissimo patre in Moscorum aula juvenili atate Archiatro natus esfe videris. Quid dicam de summa tua in Anatome peritia, ingentibusq; in eam sumptibus, quam in prælectionibus tuis in Collegio habitis jamdudum demonfirafii.

strasti. In quibus præclara non pauca eaque nova, a nemine observata dete सं xisti, imo (quod mirum) quæ in gene ca ratione animalium sagacitatem summ it Harvæi fugerunt. Ad Græcas quo attinet litteras quantus in his fueri Criticus, norunt omnes qui tecum familiarius versantur, idque ipsum testantu baud vulgares commentarii in ingeniosum at obscurum poetam Gracum Aristophanem. Quos utinam aliquando (deo valetudinem longævitatemq; tribuente) lucem aspicere dignareris. In latinis litteris quam tensus, politusque fueris bene noverunt quam plurimi ex Characteribus eximiis quos in privatum usum conscripsisti de illustrioribus bujus sæculi viris qui diem suum obierunt. Insuperque ex Annalibus Collegii dum Registrarii

3

CA

te

20.

774

00

r<sub>L</sub>

į.

Registrarii Thesaurariiq; munere fun-Eins fueris. Deinceps pietatem tuam calare piaculum est, munificentiama; in extruenda Ecclesia de Barking in qua sepulta jacent offa Patris tui dile-Elissimi, & in Parochiali tua ornanda, o ad readificandum Templum Divi Pauli, vetustissimi bujus gentis monumenti quibus Centum aliquot libras divino affectu contulisti. At hac minuta sunt præ illis quibus Collegium nostrum beasti. Domum nostram diris temporibus sub hasta positam ante annos aliquot opportune solus redemisti, quam postea syngrapha nobis successoribusque nostris summa largitate in perpetuum consignasti. Ex quo factum est ut præclarissimus Harvæus æternum decus nostrum Museum canaculuma; fundo

A 4

tuo

b

bis

tuo superstruxerit quorum utraque su pellectile, libris, instrumentisque sui Chirurgicis locupletavit, insuperq; Collegas suos ex asse paterno, dum in vivi esset, haredes dixit. Musaumque hoc Nobilissimus doctissimusque Marchio Dorcestrensis centum libris in libros

Just postra Dorcestrensis centum libris in libros de la libros de la libros de la libros coemendos erogandis primus, magnificol. Medico ceque cohonestavit. Quem sequuti alii, legavit

Joann. Seldenus scientissimus hujus seculi antiquarius Manuscripta Arab. in Medicina que collegerat omnia dedit & donavit, & Edoardus Bish Clarencieux eques auratus, & Eliah Ashmole Armiger, utriq; de repub. litteraria optime meriti Richardus Napier eques item auratus, & Dr. Scarburgh Socius dostissimus idem

Mufeum donariis suis auxerunt. Nec

u.

11

1

u

00

0

25

his contentus Hamæus fundos in bonorem nostrum, pauperumque sublevamen pia charitate ultimis tabulis legasti. Praterea ea, (qua apud omnes polles) auctoritate usus apud melioris fortunæ viros, non solum exemplo tanquam elogio actuofo præivifti, sed virtutem tuan imitari amice coegisti, & quod grati animi tui signum statuas ipsis erigi curaveris. Tu autem cum tot tantaque præstiteris, bonorem, quem amicis procuraveris, tibi a Collegio decretum non admisifi, hinc etiam aternis honoribus dignissimus. Cujus animo (ut verum Philosophum decet) nulla inanis gloriolæ cupido insidet, solo benefaciendi conscientia, aliorumque emolumentis contento. Hancque mentis tuæ modestiam testatum facit præsidendi munus sæpius oblatum,

oblatum, recusatumque, quamvis ad alia Collegii onera sustinenda paratissimum semper te ostendisti. Denique unicornu Elegantissimo Collegium nostrum locupletasti, ut haberet quo gratularetur fælicissimo, desideratissimoque reditui Regis in utraque fortuna semper Augusti CAROLI Secundi, quo summa animi serenitate ab ipso accepto gratiam tanta Majestate dignam significavit, quam postea variis modis nobis confirmavit. Et quod superiora superat sine ulla parium tuorum invidia bæc magnisica peregisti. Prætereo minores virtutes tuas, industriam, vigilantiam, egregiam in excolendis Musis industriam, animi grandis moderamen, temperantiam, sobrietatem morum tuorum elegantiam, & in nunerofis tuis apud

7

apud agros congressibus candorem, sine contentionibus Medicastris nimis familiaribus, in praxi judicium solidum, & per tot lustra fælicitatem. Has universas plurimasque alias quas nec per compendium recitare licet, semper in te expertus sum ex quo tibi innotui, insuperque amorem summum in me tuum. Multis enim me beneficiis continenter onerasti, in colloquia, studiaque tua penitius admisisti, laborumque meorum perpetuus fautor, calcarque fuisii, meque de rebus naturalibus balbutientem benigne excepisii, languentemque exuscitasti. Gratus ergo & amoris plenissumus animus justit ut in ipsius testimonium bæc pauca non pro rei dignitate dicerem, o ut bic paucas impleres pagellas cujus memoria tot tabula & monumenta

monumenta in Collegio nostro consecrantur. Ob quæ ingentia merita serena qua offertur fronte nugas has excipias oro uti tesseram obligationis, qua me in aternum tibi devinxisti, patiarisque quafo, te exorem participem ut facias remp. litterariam lucubrationum tu.zram, quod te facturum non sine magno tui honore doctorumq; emolumento spondere audeo. Præsertin cum boc beroicum, & vere regium cateris tuis egregiis facinoribus accumulaveris quod te a tumultu praxeos subduxisti, ut deo & Apollini litares. Deus O.M. te diu incolumem servet, vegetamque senectutem largistur, quod semper vovebit

Fx Muszo meo
in Domo Coll.
M.d. Lond. Cal.
humilimusq; Servus

Aug. 66. V usman Chr. Merrett.

he's long numan habit affect - Chr. Merrett.

Lestora endson Boctifi may - nom on fataly als way mendio go paulo host fort. 2.1666-cooking it - the collegy The surrainy. Hem - for extrem have to a set the standard of the publicary memorally - granter manufaction on numerally - to the grant, cum miles to memorally -

Fifth Dehmatoria.

# \$4\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$

# Lectori Salutem.

Uædam sunt (Amice Lector) de quibus monendus es in vestibu-lo hujus opusculi, quæ ad duo capita reduci poterunt, quid ante me alii præstiterint, & quid ego de novo superinduxerim. Non enim est mihi animus, nec otium in longum rhetoricare de necessitate. & utilitate hujus partis naturalis philosophiæ. Cum per se constent, feceruntq; id non pauci; Sed hie calami temperare nequeo a tenebriorii fanaticorú examinibus in quovis angulo hujus civitatis latitantibus, Qui fine ulla rerum naturalium scientia, fine vel extremis labiis degustata Anatome, duobus artis nostræ cruribus, imo sine vel levissima cujuscunq; philosophiæ artisve liberalis tinctura Medicina humano generi tam salutare aggrediuntur. Hi inqua pleriq; diris temporibus proditione & contumeliis remp. Ecclesiamq; pessundiderunt, qui regiam majestatem, Nobiles, totumq; clerum convitiis suis turpissimis

aunit of alunds advocated a protection of annu for grant som grant some from the gold from the stand of the form of the form of the form of the community of the protection of the form of

prosciderunt, totam jam suam pristinam maledicentiæ fentinam nullatenus laceffiti in Collegium nostrum (qued inventis scriptisc; suis honorem regni tunc dierum conservavit) Evomunt, idque in Collegium quia Collegium uti prius in Regem quia Regem suosq; calamos fædasg; linguas consueti exercent idque in singulos Collegas in priores refusuri, si eadem licentia ac impune ipsis liceret. Non omnino isti mores mutarunt, sed aliorfum converterunt, & fub Medicoru larva nefarii histriones priores agunt partes, in maximum reip. & religionis damnum, & cum non poterint legibus cohibiti in animas fertunasq; hominum debacchari, in corpera tamen credulorum heu nimis licet. Quod hinc evenire palam eft,quia in eodem statu ars nostra fita est, quo non ita pridem religio erat, quando libertate conscientiæ (uti vocabant) freto cuivis in utramque involare fas fuit, sie jam nulla legum impunitate coercici professionem nostram omni avo facratissimam irruunt. Hinc factum est ut nullus pene miles, nullus fanaticus, præcove seditiosus nullus Circulator, anus, nutrix,

nutrix, obstetrix, balneator, veterinarius proditorius inventus fuerit, qui non una Needham nocte fungorum modo in Medicum excreverit, quod majus est quidam Chymiam folo verbo professi unoq; fornaculo eoque in Camino erecto instructi, ingentibusq; promissis turpissimisq; contumeliis sufierti methodum a bismille annis approbatam evertere, novamque fumme perniciosam substituere conantur; Sed ut horum fraudes verbo doceam, scire licet ex Kircheri Oedipo primos hujus artis profesiores Egyptios medicamentum universale collimasse & tentasse solum, at hæc fiducia lapsos suamq; fidem liberare nequefuntes tandem ad Elixir, seu lapidem Philosophicum confugisse quem sub variis anigmatibus a nemine intellectis proposuerunt, quibus technis principum, nobiliumque crumenas diu emunxerunt quousq; legibus nostris sub pæna capitali cautum fuit, ne quis artem multiplicandi profiteretur, uti ex doctifimis Commentariis in eandem legem Edoardi Coke militis luculentissime patet. Sic autem seclusi de-

nuo

nuo nunc dieru ad priores fi audes rede unt, & ignavo vulgo obtrudunt nescio quæ fecreta universalia, grandia ne dici ridicula pollicitantes ex animis deorum Ethnicoru extracta (risum teneatis amici) & multa hujusmodi quibus chartam conspurcare nolo. Adeo ut Paracelsus ipse de his sateri cogatur, Chymiam omniu corum inhoneste se alere agentiu agrum esse, quoi pleriq; fuere suæ professionis desertores, ad incitas redacti, dentistagi, ferrarii verlipelles, fartores, cateriq; decoctores, aliiq; hujus furfuris vani, rudes & imperiti homunciones convitiatores & jactatores in utriusq; impudentissimi mendacissimiq; qui ubiq; tanquam muscæ carnes humanas depascentes circumvolitent, insusurrantes se Monarchas & Medicamentorum divos, seque solos efficacia præparare, quod fi facto & experientia comprobarent ipsis fide adhibenda esset. Quæ quam cito obsolescunt & in fumos abeunt, confirmarunt pupera quædam remedia in omnium fere ore celebratissima quæ intra paucos menses evanuere. Ultima æstate sub initium pestis quam mira prædicarunt falsarii Chymici

Chymici seque totam pestem spiritu oris difflaturos, at reipsa ostenderunt, quanta vanitate. Nam magna pars istorum qui hujus curam profitebantur,occubuerunt, interea superstitibus ad id munus a Collegio laudatis, & a civitate sepositis non fine magno medicamentorum Chymicorum opprobrio. Neque tamen sentias me prorfus hanc artem vituperare utpote ad medicamentorum præparationem aliquatenus utilem, & in detegendis naturæ mysteriis apprime necessariam, solum Agyrtas, fumososq; Empiricos procul habeo, qui veræ Chymiæ penitus ignari fub arcanorum prætextu ventosam plebemingentibus promissis lactant, quæ si detegerentur, authores repropinandos darent. Verbo dicam nonnulla horum artificia. Quidam ex his pseudochymicis ut majorem secretis suis auctoritarem conciliaret, & ad vulgus captandum librum edidit curationum suarum, quas inquisivit hortatu meo Coll. Lond. probatus Chymicus Johnstonus comperites plerasq; plani sictas, quasdam nullius momenti, imo quosdam ab ipso conductos ut falfitatem curationum astrucrent, B uti

ũ

(

f

r

1

r

(

1

uti patet ex Johnstoni dicti in eundem | brum observationibus. Addo non inusita tum esse illis morbos vel ignorantia, ve fraude creare, ut tandem de iis vel spon te sua evanescentibus, vel fortuito curati triumphent, & per præcones suos in locis publicis deprædicent. Præterea sermonibus, scriptisq; suis Chymica prætexunt, usu autem Galenica comprobant. Horum Antefignanus quinq; librarum mercede processus (ut vocant) quosdam vulgares edoctus protinus librum conscripsit specie majorem, at demptis convitiis, tautologiisque, minimum, nihilq; folidi in fe continentem, in quo sua Chymica ad calos elevat, Galenica infernis devovet, cui (ut obiter dicam) tres Collegæ responsum dedere, contra quos ne hiscere quidem ausus est. Idem tamen nuper evulgavit experimenta, & praxin famolissimi Emperici Trigge ( quem pro more suo illiteratissimum licet titulo Doctoris infignivit, funt enim omnes apud nos prohpudor Doctores, ) in quibus nil fore invenire est præter tritissima Galenica,& quorum lectio castas lædit aures. Idem tamen in præfatiuncula plurimum hæc laudet.

li ta

tis

cis

o-It,

m

le

es

e. O-

ſe

2

ui

b

1-

11

1

e

1

laudet, evehitq; & hujusmodo in nupera peste usus est, quamvis optimorum delectu caruerit. Et quod ad praxin suam Chymicam attinet, hic nuper in quæstionem vocatus propter melioris fortunæ virum inter arcanorum suorum operationem mortuum, quem ut Lue Venerea affectum curandum susceperat, (cum revera Arthritide laboraret)omnes quippe Morbi ridicule ab ipfo vel pro hac lue vel pro Scorbuto habentur, in quæstionem (inquam) vocatus, fassus est fe solo Mercurio dulci usum fuisse. Unde palam fit, hos titulares Chymicos medicamenta sua omnia vel saltem quædam ex iis ex pharmacopæo nostro, vel aliunde suffuratos pro suis jactitare. Jamque tandem ad facram Empiricorum anchoram confugerunt, quibus morbificam materiam figunt cum volatilem reddere præstiterit, sicq; Morbos faciunt Chronicos, nisi interea somnum induxerint æternum. Hæc pauca, firem ipfam, nimia (fateor) si Epistolam respexeris, ab horum maledictis (quæ quotidie aures meas ubique circumsonant) abreptus, dixi. Nec opus est ut hic vindicarem

Collegas, cum toti mundo innotuera n quantum Anatomiam, aliafq; Medicin n partes adauxerint, cum interea hi con v vitiatores nihil lectione dignum proti lerint, specimenve ullum dederint f n efficacia, utiliaq; quæ prædicant remedir in medium produxisse. Neque unquan p audivi quemvis virum ingenuum feme horum vanis & ingentibus promissis se v ductum, denuo in eundem errorem in cidisse. Sed manum de hac tabula, ne ceandem cum iis insaniam insanire videar l Hoc tantum adjiciam, si in posterun s Collegii nostri famam lædere perrexe rint; acrem me habebunt; verifimume vindicem, fraudesq; detegam ab artis hujus incunabilis in hunc usq; diem usur patas ides reipsa comprobabo haud suf pectæ fidei testimoniis, simula; omnii stigmata Societati nostræ inusta deter gam. Sed ut ad rem, unde nimis digref fus fum, redeam.

Hanc philosophiæ partem primus excoluit Dioscorides, qui pauca quædam in singulis generibus posteris reliquit. Etiamsi pro more suo Helmontius, ejusqui sequaces nihil post ipsum a bis mille an

nis additum fuisse asserant, quod falsissiim mum esse comparebit cuivis vel leviter or versato in ejus commentatoribus cæteth rifq; Scriptoribus modernis qui hanc materiam tractarunt. Adeo ut affirmadi re contra illos non dubitem viginticuan plo auctiorem esse quam Diasc. ævo.

ne Internaturalium rerum Scriptores inse venti multi qui Botanologiæ suam imin penderunt operam, in cæteris pauci. An n quod præ cæteris plantæ magis tractabiles essent, diutiusq; durarent; An quod in fublevando humano genere. & quoad vitam sustentanda, & quoad salutem tutandam reparandamo; magis in usu essent?

Quidam in universum de his egerunt, alii de iis folis quæ in particularibus reir if ia gionibus reperiuntur. Quod inde factum arbitror, ut scirent lectores quæ cuivis so-2 lo & regioni convenirent & unde harum rerum curiosi peterent, & inventas facilius distinguerent. Cum magnum sit in omni inscientia compendium præscindereab infinito.

£

0

m

t.

Inter posteriores agmen ducit Gesn. qui montem Fractum, sive Pilatum in Helvetia describit; dein Joan. Pona mon-

tem

tem Baldum in agro Veronensi, Cass Bauhinus edidit Catalogum plantarum circa Basileam crescentium, Schoensel scripsit de stirpibus & fossilibus Silesia.

Inter nos hanc provinciam primus ag greditur Thom. Johnsons Pharnacopau Lond. vir egregie industrius, qui in usun sociorum suorum 1632. edidit plantas in Ericeto Hamstediano crescentes, poste vero, quas investigavit in itinere suo in agrum Cantianum ad eam rem suscepto Anno. 1632. & Anno. 1634. Alphaberi ce exhibuit Mercurium Botanicum, & de thermis Bathoniensibus tractatulum & una cum eo Mercurii partem alteran Majora certe præstitisset, nisi decore pro rege patriaque sclopeto transfossus occubuiffet. Hunc fequutus Dr. How, Phytologiam Britannicam Anno. 1650. edidit, auctioremq; reddidit, qui festim morte abreptus quæ ulterius destinaverat, peragere destitit.

Postea sc. Anno. 1660. Ds. Agay Collegii Sanctæ Trinitatis quod apud Cantabrigienses est Socius, Catalogum Cantabrigiensem publicavit, librum apprime utilem, magnoq; cum judicio concinna-

tum,

tum, dignumq; qui omnium manibus teratur, in quo tradit solummodo eas quas ipse cum sociis investigaverat, & intra angustos limites solius provinciæ Cantabrigiensis, hinc sactum ut universalis Ca-

talogi loco inservire non poterat.

un fell ag

in in each to the to the to

Phytologiæ exemplaribus tunc pene distractis, oravit Bibliopola mihi familiariffimus ut in manus fumerem, cujus honestis votis contra niti non poteram, ascito ergo in operis societatem perito Botanologo Dre Dale, ut sociatis humeris novum Catalogum, posteaque junctis impensis historiam earundem ederemus. Et ad hanc rem Octingentas plus minus iconas Johnstono procurante Sculptas emi. Sed dum hæc ago immatura nimis morte abreptus erat. Nec tamen a proposito destiti, & nactus virum scientia Botanica tune mediocriter tinctum, at in distinguendis investigandisq; plantis fælicissimum, hujusq; studii avidissimum, Tho. (inquam) Willeselum propriis meis impensis quinque annorum spatio per universam Angliam totis astivis mensibus misi, qui non paucas collegit huic pinaci additas. Insuperq; filius meus natu minor Christo-

Christophorus aliquot itinera ad ampliandum regnum vegetabilium suscepit.

Sed cum in supra memoratis scriptoribus Animalia fossiliaq; nostra desideren tur, quæ defuerant addidi, Pisces, volt cresq; quotcunq; vel ichyopolæ, vel Ornithopolæ Lond.aucupes piscatoreswe suppeditare poterant conquisivi uti e tiam lapides metallaq; a variis mineris transmissa. Plantas plerasq; hic insertas in horto meo ficco fervo, cuivis ingenio. fo paratus oftendere. Et hæc omnia non parvam rerum naturalium partem continent, adeo ut dicere ausim, qui hæc intimius perpenderit, sibique familiaria reddiderit, cætera exotica quæcunque fuerint, facile comprehendet. Tam dis ves felixq; est folum, regioq; nostra ut vix sentio ullam aliam paris spatii tot & tam diversa continere, atque de nostra noumou cum Euripid. licet.

Οὐρανὸν τοῦ γῶς ἔχρων ἐῦ κεκρομένον "Ιν ἔτ' ἄχαν πυς, ἔτε χείμα συμππυεί Ἡ Α' Ελλάς 'Λοία τε κάλλισα τρόσου Τῶς γε δέλεας ἐντῶυδα συνθης διομέν.

Quæ sic latine reddidit Grotius in ex-

F

1

Bene teperato nostra sub calo est humus Non nimia torret slamma, non minium

Europa quicquid, Asia fert quicquid boni Potissima horum seduli in nos vertimus.

Et optandum sane esset viros doctos ubivis gentium idem mecum conarentur, ut facilius arctiusq; commercium inter viros peritos oriretur, tandemq; certior historia naturalis contexeretur.

.

Neque tamen arbitreris me rem confecisse, procul absit ut ita dicam, largius mihi indies crescit penu, multasq; plantas possideo quarum nulla hic mentio, quoniam non satis mihi constat quænam suerint, sed ab aliis hactenus descriptis dissere hoc mihi satis perspicuum. Restat adhuc multum operis, rectabitque quod cum auctareo expectare poteris.

Sed jam de singulis partibus aliquid dicendum venit. Quod ad Botanicam spectat, Ducentas præter propter plantas omisi utpote neq; a meipso vel ab aliis viris peritissimis ullibi repertas, sed a dubiæ sidei amicis auctoribusve communicatis. Quasdam tamen adjeci a superioribus scriptoribus memoratas quia lo-

cum nullum affignarunt, satisq; novi illo nunquam vidisse. Insuperq; locos addidi rarioribus, quibus deerant. Nomina magis vulgaria selegi eaq; præ a'iis qua in prioribus catalogis inserebantur, mutatis tamen paucis, substitutisq; iis qua apud Gerardum & Parkinsonum habentur, & singulis paginam ascripsi in qua apud ipsos repperiuntur, ut certo sciat lector quas volo esse nostrates. Hos autem duos aliis prætuli utpote nobis maxime familiares, maximeq; universales,& ideo utrosque, ut qui hoc careat, illo uti posset. Cum autem Gerardum dico, nolo librum ab ipso editum, sed a Johnstono emaculatum intelligi. Quæ autem in utroq; horum desunt, aliquem Botanographum alium in quo repperiuntur, indigitavi. Et ut suus cuique honos constet. & fartus testusq; conservetur, scias velim eas plantas quas Johnstonus proposuit poni fine asterisco, & quas Phytologus hoc signo Ph. notari, a me autem additis asteriscum præfigi.

Cum autem ingenuum sit agnoscere per quos profecerim, non prætereundi sunt in hac arte Coryphæi quorum opera

adju-

adjutus sum. Inter quos prælucet Dr. Bowles Collegii Medd. Lond Socius honorarius, & in herbarum cognitione præcellentissimus. Proximum locum vindicat Do. Brown Collegii Magdal. Socius & Sacræ Theolog. Bacchalaureus vir exercitatissimus, & Catalogi Oxoniensis author eruditiffimus, Ds. Witham Eboracen. s, Ds. Gunthorp medicus Cornubiensis, & affinis meus Ds. Jenner Theologus. Sed omnibus palmam præripit Ds. Goodyer Hantoniensis vir incomparabilis, & siquis alius, acerrimi judicii, maximæq; industriæ, imo paucis auctoribus postponendus five exactas descriptiones sive accuratas distinctiones animadverteris, uti cuivis Gerardum emaculatum consulenti facile patebit. Is enim erat qui maximam partem dicto libro & Mercuriis supra memoratis communicavit, uti ex litteris hinc inde missis constat quarum me participem fecit Ds Tauldon Goodyeri nepos uti etiam Manuscriptorum e jus ex quibus quadam in hoc tractatulo me mutuo accepisse in progressu operis liquebit. Deinque ex Catalago Cantabrigiensi supra laudato uti etiam ex appendice transcripsi

scripsi quacunque nondum innotuerunt nostris scriptoribus ipsiq; adscripti quamvis quasdam ex illis prius detexeram.

Sed objiciet fortassis quispiam me nimis audacule quasdam plantas huic Catalogo inferuisse, præsertim Hyoscyanum Peruvianum, Caryophillos Carui, &c. Respondeo id ideo abs me factum ut ostenderem quasnam terra nostra ferat, & si ipse ob hoc vapulem, etiam alios incusandos venire, qui sata plurima proposuerunt, viz. frumenta pastinacas, legumina quædam, linum & Cannabin, &c.

quod nemo in aliis vituperavit.

Ulterius dici potest me plantas variegatas omilisse. In promptu responsum, me observasse plerasq;nostrates hanc mutationem subire, neq; perennem esle, ubi autem duraverit hic color ascititius me non neglexisse. Et quod ad flores attinet, inusitatum non est omnes cæruleos, luteosq; præsertim pallescentes in albos mutari, quod herbis senescentibus accidit uti etiam a loco fertili ad steriliorem tran-Memini biennio elapso Blattariæ femina ex horto meo in murum lateritium vicinum (nescio quo casu) translata

ex flore flavo in candidum permutata fuisse; interim iisdem infra in lætiori solo pristinum retinentibus colorem. Idem etiam eadem æstate observavi in Cichoreo, cujus flores cærulei senescente planta lacteum induebant colorem, ita ut ex quingentis nullus apparuit cæruleus, qui candor etiamnum (dum hac scribo) post triennium perdurat. Cum aliæ plantæ ex prioris seminibus deciduis pronatæ nativum servant Cæruleum.

Tertium a me prætereundum non est, quædam forsan hic inseri quæ in dictis locis vel nuspiam apud nos occurrunt. Fateor sic sieri posse, & nullus dubito quin quædam oriantur de novo, denuoq; pereunt, quod in plerisq; capitibus satis patet. Inter quadrupedes Castor in Fluvio forra Tivy in agro Merionethensi prioribus sæculis abundavit, qui nunc dierum nusquam apud nos comparet. Et Lupi adeo pecoribus infensi quonda erant, ut ad hos exterminandos lege cautu fuerit, ut cujusvis criminis rei vitam stato lupinorum capitu numero redimerent, unde factum erat ut brevi penitus occiderentur. Præterea ex statutis regni manifestum videtur, quosdam

quosdam pisces in illis memoratos, jam nuspiam conspicuos esse. Deinceps aves funt migratoriæ quæ raro redeunt nisi frigoris importunitas & in aliis gentibus victus penuriæ huc compulerint. Insecta etiam nova quotannis occurrere apud omnes in confesso est qui intimius indagarunt. Non quod novas species vel æquivoca (quod aiunt) generatione produci crediderim, sed accidentalibus quibusdam differentiis ab invicem difcriminari compererim. Inter plantas autem excrementa terræ viz.muscos,& fungos certis locis perire, & in aliis oriri ex-tra omnem controversiæ aleam ponitur, quod etiam de vitiis frumentorum nuncupatis præsertim in Avenaceo genere facile quispiam observare poterit, uti eti-am in Arboribus fructiferis quarum semi-na terræ commissa diversos fructus proferre vel rusticis innotuit. De fossilibus dispar est ratio quæ omnia perdurare fentio.

Ultimo forsitan desideraverit lector descriptiones plantarum noviter additarum. Dico id propterea a me omissum ne nimium intumesceret hic liber quem in

pera gestatilem sore proposui, præterea hunc desectum me abunde supplevisse arbitror, talia imponendo nomina quæ descriptionis loco inservire possunt.

In Quadrupedibus, Avibus, piscibusq; tradendis consului Gesnerum Aldrovandum, & Johnstonum qui amplissime de his tractarunt, & in quibusdam Turnerum nostratem inter viros sua atatis exercitatissimum qui librum de Avibus edidit mole parvum at judicio majorem, de piscibus etiam nostris præfixam videas epistolam Gesnero, qui etiam ubivis laudat Caium in materia medica apprime versatum, Pennium & Falconerum Collegii nostri lumina. Insecta quod attinet Mouffetum sequor, a quo cæteri plurima mutuati sunt. Et certe hæc naturalis philosophiæ pars maximæ voluptatis, varietatisq; (in qua natura penu suum exhausissie videtur) pene ab omnibus seculis neglecta jacuit, quousq; Nostrates Pennium dico Wottonumq; qui inchoarunt, & Mouffetum qui perfecit, manus suas huic operi admoverunt. Cujus o-pus pudet sane inter Blattas tineasq; relinqui, cum nemo prius solidiusve hoc penfum

pensum absolverit. De cæteris hujus opusculi partibus non est opus ut plum

dicam quam quæ singulis præmisi.

Finem ergo imponam huic Epistola cum duo candidum lectorem exorave. rim, primum quod omnes ante me scriptores fecerunt, sequuturiq; facturi sunt ut Errata Typographia emendet. Secundum ut si qua in re in tanto itinere cespitaverim, (quod primo conatu in materia tam lubrica vix possibile erat non fieri) me ut moneat, dein siquid omissum, amice communicet, & ad me transmittere dignetur, quod non fiet fine grata Benefactoris memoria, & sine impensis ullis. Sicq; tandem brevi comparabitur fupellex idonea ad compilandam hiftoriam non fine ingenti patriæ nostræ honore, & reip. litterariæ emolumento. Tu interim vale & fruere (quæ fola promitto) fide, impensis industriaque mea.

C. M.



#### A.



Bies, the Fir-tree, in Arboreto Hamstediano, & Alpibus Scoticis, & quondam in plurimis Angliæ locis Borealioribus, ut postea constabit. Ger. 1363. P. 1540.

Abies marina Belgica

Clus. G. 1574. Abies & Cupressus marina, P. 1302. sæpe adnascitur ostrearum testis.

Absinthium album, G. 1101. Absinthium album feu umbelliserum, P.99. in montibus Walliz.

Absinthium tenuisolium Ponticum Gal. Ger.

1096. Absinthium Ponticum sive Roman.

vulgare, P. 98. in falsis & maritimis.

Absimb, latifolium, sive Ponticum, G. 1096. Absinthium vulg. P. 98. in simetis & ruderibus.

Acanthium album, G. 1149. Acanthium vulgare, P. 979. Cotton Thistle, ubique ad viarum margines, variat quandoque flore albo.

Acer majus, the great Maple, G, 1484. Acer majus latifol. Sycomorus falfo dictum, P. 1425. juxta ædes nobilium.

man

Acer

Acer minus, G. 1484. Acer minus five vulg. P. 1425. in fepibus & fylvis.

Acetofa, our ordinary Sorrel, P. 742. Oxalis five

Acetofa, G. 396. ubique in pratis.

Acetofa minima, sive Oxalis minor, in pratis siccioribus, Sheeps Sorrel, G. 398. Acetosa minor Lancelota, P. 744. in montosis.

Acinus, stone Basil, G. 675. Acinus, sive Clinopodium majus, P. 22. Variat flore albo, in

montofis & cretaceis.

\* Acorus verus, seu Calamus officinarum, The sweet smelling flag, P. 140. Acorus verus officinis falso Calamus, G. 62. sound by Doctor Brown neer Lyn, and by Mr. Brown of Oxford, near Hedly in Surry.

Adianthum album, five Ruta muraria, five Salvia Vitæ, Wall rue, and Tentwort, G. 1144.

P. 1050. on old walls at Windfor Castle,
Bristol, and Bath, and many other places.

\*Adianthum album, five Ruta muraria foliis vari egatis, on a wall by Mr. Wades house at Kilnfey in Craven. Mr. Witham. Hanc ideo adject quod plantas Capillares hanc mutationem subiisse à nemine (quod memini) observatum est.

Adianthum aureum majus, seu Muscus Capillaris, Great Godly Locks, or Golden Maidenhair, G. 1559. Polytricum aureum majus, P. 1052.

Adianthum aureum minus, cum priore in mu-

ris paffim, G. 1559. P. 1052.

\* Adianthum aureum minimum, vix semiciuncialis est altitudinis, caulis tamen & flos æquant illos Adianthi aurei maximi, Lanceston Cornwall. Mall, ubi copiose provenit. Dr. Gunthorp Ægilops Bromoides, Bearded Wild Oats, G. 77. Ægilops Bromoides Belgarum, Dutch

Havergrafs, P. 1149. in fatis.

Agrifolium, Holly tree, G. 1338. five Aquifolium P. 1487. passim in locis sterilibus. Hujus baccæ ante maturitatem suam lutescunt, & in eadem arbore, solia tam lenia quàm aspera vulgo conspiciuntur, in annosis & procerioribus pleraque lenia, adeò ut non opus suerit cum Ph. in plures species dirimere.

\* Agrifolium fol. variegato, Neer Highgate,
Ob elegantiam in plurimos hortos Londi-

nenses transfertur.

Agrimonia, Agrimony, G. 712. Agrimonia vulg. P.594. ad sepes, & agrorum margines.

Alcaa vulgaris, fine cut, or Vervain Mallows, P. 301. Malva Verbenaca G. 931. in pratis & sepibus, sed rariùs, variat flore albo.

\* Alcaa foliis non dissectis & pedunculis longioribus, hanc mihi communicavit, Dus

Pink ab Herefordia.

Alchimilla, Lyons foot, or Ladies mantle, G. 949. Alchimilla major vulg. P. 538. in sylvis & umbrosis, & pratis quibusdam.

Alchimilla minor, G. ib. 4. miles Northward from Stanford, in a lane on the right hand,

but not at Wombwell.

Alga, Grass Wrack, G. 1569. Fucus marinus five Alga marina graminea, Wrack, or Sea Weed, or Grass, P. 1292. in maritimis, & folio angustiori, from the Severn Sea.

2 Alga

Alga Bombycina, Silken Wrack, in maritimis, Alga membranacea Ceranoides, Broad leav'd or horned Sea. wrack, maanunipus porosa I.B. Alliaria, Samce alone, or fack by the Hedge. G.

794. P. 112. pa ffim ad sepes.

Alliam Ursinum, Ramsons, G. 179. in sylvis & fepibus.

Allium sylvestre, Crow Garlick, G. 179. P.

871. in pascuis pinguibus & pratis.

Alnus hirfuta, rongh leav'd Alder, G. 1477.

Alnus folio inaco, P. 1409. Ph.

Alaus nigra, sive Frangula, Black Alder tree, G. 1470. Alnus nigra baccifera, sive Frangula, the Black Alder Tree, P. 240. in sylvis humidis.

Alnus, the Alder Tree, G. 1477. Alnus vulg.

P. 1409. juxta aquas.

\* Alnu cortice obscure purpureo soliisque parum dissectis, Betwixt the South gate and the Bridg at Hereford.

Alopeeures maxima Anglica paludosa, Great Marsh Foxtail, G. 872. P. 1166. in a Salt

Marsh by Eriff Church in Kent.

Alfine Palustris, Marish Chickmeed, G. 613. Alfine aquatica, sive Palustris major. P. 1259.

Alsine Baccifera, Berry bearing Chickweed. G. 614. Alsine repens Baccifera, P. 759. in sylvis udis.

Alfine Cochleariæ longæ facie, between the two Parks at Eltham, and divers places neer Len-

don.

Alsine Corniculata Clus, at West-gate Bay in the Isle of Thanet, G. 616.

Alsine

Alsine Cruciata marina, vel Anthyllis lentisolia, G. 622. Sea Pimpernel. Anthyllis lentisolia

marit. P. 281.

\* Alsine flosculis conniventibus, found by Mr Goodyer in Hamshire, and by him properly Named Blinks, as also frequent in Hide-Park.

Alsine foliis Trissaginis, Germander Chick weed, G. 616. P. 764. in arvis & ad agrorum

margines.

Alfine foliis Veronicæ, Speedwell Chickweed, G.
613. Alfine foliis subrotundis Veronicæ, P.

762. in arvis & tectis.

Alsine Hederacea, Ivy Chickmeed, G. 616. Alfine folio Hederulæ minor, P. 762. Hæc planta potius dicenda Veronicæ species.

\* Alfine floribus ad instar Polygoni marini, ad fingulas alas albis. This was fent me from

Cornwall. N. D.

Alsine Hederula altera, Great Henbit, or Codwort à seminis figura, G. 616. Hedurelæ solio major, P. 762. Hæe planta est potius marubii species quam Alsine.

Alsine longisolia uliginosis proveniens locis, 7.

B. Alsine Fontana, G. 613. Fountain, or

[mall long leav'd water-Chick weed.

Alfine major G. 611. Alfine maxima, P. 1259. ad sepes.

Alfine minor media, G. 611. P. 1759. passim

ad sepes.

Alfine minima, G. 612. Fine Chickweed. Alfine aquatica minima, P. 1259. in muris & tedis.

C 3 \* Al-

\* Alsine minor Androsaces alterius facie, Buah. in prodromo, P. 118. 5. found by Mr. Halilah Apothecary in Lincolnshire.

\* Alfine Myositis procerior & longiore folio, datur etiam alia folio obtutiore, Mansear

Chickweed, paffim in agris.

\* Alfine Myositis humilior & rotundo solio cum priore. Both those neer Hamsteed Church, and in Hide-park.

Alsine palustris serpillisolia, Creeping water

Chickweed, G. 614. P. 1260.

Alfine recta, Right Chickweed, G. 612. P. 1260. in locis umbrosis.

Alfine rotundifolia, sive Portulaca aquatica, Round leav'd Chickweed, or Water Purstan,

G. 614. P. 1260.

\* Alfine tenuifolia, 7. B. Facie extern. lino cathartico multum affimilatur, Narrow leav'd Chickweed, on the Hills beyound Guilford, and on the Borders of Triplow Heath, CC.

\* Alyssum mont. Columnæ, Fabius Columna bis Madmort, P.589. in Ericeto Hamstediano.

Althea Ibiscus, Marsh Mallows, G.933. Althea vulg. P. 304. in maritimis.

Ampelopressum, five Portum sylv. Wild Leek, G. 176. Ampeloprasum, P. 871. amongst Barley near Beachenton, betwixt Oxford and

Ranbury, Mr. Stonehouse, Ph.

Anagallis, five Becabunga, Brooklime, G. 620. Anagallis aquatica vulg. five Becabunga, P. 1236. variat quandoque flore albo, ab Herefordensibus Limpwort.

Ana-

Anagallis aquat. folio oblongo crenato, Great water Pimpernel, P. 1237. 3. fine icone.

\* Anagallis aquat. flore parvo viridi caule rubro, in a great Ditch neer the Moor at Pe-

ters field Hamsbire, Mr. Goodyer.

Anagallis aquatica major, Great long leav'd Water Brooklime, G. 620. Anagal. Aquat. five Becabunga major, P. 1237. 2. fine icone.

Anagallis aquatica rotundifolia, G. 620. Anagal. tertia Lob. folio subrotundo non crenato. P. 1237. 5. round leav'd Brooklime, 2 miles below Graves end.

Anagallis aquatica quarta Lob. Lobel's 4th water Brooklime, G. 621. Anagallis aquat. angustifolia sive quarta Lob. P. 1237. in Tuthil-fields.

Anagallis aquat. angustifolia, scutellata, P. 1237.
7. nulla icon. scutellata C. B. & J. B. Nar-

row leav'd Brooklime.

Anagallis mas. Male Pimpernel, G. 617. Anagal, floribus phæniceis, P. 558. in agris.

Anagallis femina, G. 617. Anagallis floribus ceruleis, P. 557. in arvis, plentifully beyond Croyden.

Anagallis lutea, G. 618. Anagallis flore luteo, P. 558. in Charlton wood beyond Greenwich.

and Hamsteed wood.

Anchusa degener sacie Malii Solis. Bastard Gromel, G. 610. Lithospormum arvense radice repente, P. 432. inter segetes.

\* Anchufa lutea Columnæ, by Bexley in Suf-

fex, in the way to Rumny Marsh.

4 Andro-

Androsamum vulg P.575. Clymenum Italorum, G. 543. Tutsan, or Park leav's, on Hamsteed Heath.

Androsamum Hypericoides, G.543. Hypericum majus sive Androsamum Math. P. 575. Tutsan St. Johns wort, in montosis.

\* Androsemum Hypericoides birsutum, hairy

Tutsan, in Charlton Wood.

Vincents Rocks near Bristol, and on the Hedge side beyond Lewsham.

Anemone nemorum, the wood Anemone, or Wind-flower, G. 374. passim in sylvis.

×

Anemone flore toto rubro. in Devonia.

Anethum, Dill, G. 1033. P. 886. feritur in a-gris.

Angelica sylvestris, wild Angelica, G. 999. P.

940. in fylvis humidis.

Angelica fativa, Garden Angelica, G. 999. P. 940. amongst the Rocks not far from Barwick.

Anonis, five Ononis spinosa purpureo flore, Prickly rest harrow, with a purple flower, P. 993. Anonis sive Resta bovis, Cammock, G. 1322. variat flore albo, in pascuis humidis.

Anonis non spinosa purpurea, G. 1323. P.

993. variat etiam flore albo.

Anthyllis Leguminosa, Kidney Vetch, G. 1240. Anthyllis leguminosa vulg. P. 1093. in montosis.

Antirrhinum minimum repens, [mall creeping Snap Dragon, or Calves [nont, G. 549. Antirrhinum Thinum Sylvestre minimum, F. 1334. in agris.

Antirrhinum fylvestre medium, F. 1334. Antirrhinum minus, G. 549 cum priore.

Aparine Clivers, or Goof-grass, G. 1122. Apa-

rine vulg. P. 567. in fepibus.

\* Aparine major, Plin. G. 1122. Boraga fylv. minor, P.765. à quo male describitur, Small wild Buglos, great Goof grass. In the hedge of a lane leading out of the Field to Newmarket, and in the Close adjoyning to the Church-yard. CC.

\* Aparine levis. P. 567. fine icone, in moist

pits on Hounflow Heath.

Aphaca, the yellow mild Vetch, G. 1250. P. 1067. in arvis fed rarius.

Apium vulg. five palustre, Ordinary Smallage, P. 926. Eleofelinum sive Paludapium, G. 1014.

in uliginosis.

Aquilegia Cærula, Blew Columbine, G. 1093.
Aquileg. vulg. fimplex, P. 1367. about
Broadsworth, and Hample woods, betwixt
Leusbam and Bromley in Kent, and many
places of York-shire.

Arachus, siye Cracca major, Great wild Vetch or Tare, P. 1071. Vicia sylv. sive Cracca

major, G. 1227. in agris.

\* Arachus, seu Galega sylvestris, Birds Tare,

passim in pratis sylvis & sepibus.

Arachus, feu Cracca minor, Small wild Tares, P. 1070. Vicia fylv. five Cracca minima, G. 1228. ubique in fepibus.

Arbnins, The Stramberry-tree. G. 1469. P.

1490. with indented leaves, in the West of Ar Ireland.

Argemone capitulo longiore, Long codded will An Poppy, G. 373. P. 370. in arvis frequention,

Argemone capitulo torulo, G. 373. Argem. Capitulo rotundiori, P. 370. in fatis fed rari or, variat flore albo.

A

A

Argemone lutea Cambro-Britannica, yellon baftard wild Poppy. P. 370. in Wales.

Argentina, Silver-weed, or Wild Tanfie, G. 993. Potentilla, P. 593. ubique ad semitas.

Aria Theophrasti, Wild Service Tree, or Cum. berland Hawthorn, G. 1327. Sorbus sylv. Aria Theoph. dicta, P. 1421. On Hamfteel Heath, and the Hedges by.

\* Aria Theoph. fol. obtusis. Pin. 452. At San

driff in Kent, Mr. Goodyer.

Aristolochia longa, Long Birthwort, G. 846. P. 291. near Redding, sed in loco ubi quondam fuit Monasterium, Mr. Brown.

Armeria flore simplici, William with a single

flower, in a Wood beyond Redding.

Armeria prolifera Lob. G. 599. Childing Sweet Williams, in the Meadows betwixt Hampton Court and Tuddington.

Armeria sylv. humilis flore unico rubello.

Armoraria, five Lychnis plumaria mas florealbido elegante, by a Ditch in a long lane betwixt Daventry and Dove-bridge in Northamptonshire, Mr. Stone-house, Ph.

Armoraria fæmina flore albido elegantiori, Neer Gosworth bridg in Oxfordsbire. Idem. Ph. Arme.

Armoraria pratensis mas, male wild Williams, G. 600. variat flore albo.

Armoraria pratensis flore pleno, double flowred williams, G. 600. in Nether-wood hall grounds in Darfield Parish, Yorkshire, Mr. Stonehouse Ph.

Arthemisia, mater herbarum, Common Mugi wort, G. 1103. Arthemis. vulg. P. 90. in

agrorum marginibus.

Pil4

or, Ca-

ri

07

3.

ŋ.

4

ŀ

.

\* Artomisia marina, G, 1104. betwixt Deal and Dover.

Arum vulgare maculatum & fine maculis, Wake Robbin, or Cuckow pint, with and without fpots, G. 834. P. 372. passim ad sepes.

Arundo vallatoria, Common Reed, G. 36. Harundo vulg. five vallatoria, P. 1209. in stagnis & udis.

Arundo vallatoria 30 pedes longa ex insula Vecti.

Asyrum, St. Peters wort, G. 542, Ascyrum

vulg. P. 574. in dumetis.

Ascyrum supinum villosum palustre, P. 574. Ascyrum supinum Exodes G. 542. in a bog East of Croyden.

Asparagus sativus, Garden Speragus, G. 1110.
P. 454, in the Marshes beyond Bristol.

Asperula flore Caruleo, Blew Wood-roof. G. 1124. R. 562. In the Woods neer Hamsteed.

Asperula, Wood-roof, G. 1124. Asperula aut Aspergula, P. 563.

Asperula quinta, G. 1125. Small red fl wr'd

Mood-roof, in Tuthil fields, and Black Heath
Asphodelus Lancastriæ verus, Lancastriæ Asphodelus major, sie
palustris P. 1218. in bogs in Surry, 2 mile
beyond Mr. Moors house.

Asphodelus palustris carinato folio, ND, Smal

mater Asphodil.

Asplemum five Ceterach, Spleen wort, or Mil mast, G.1140. P. 1046. in Saxorum crepid. nibus & templorum muris.

After Conizoides Gesn Fleabane Startwort, G.

487. P. 131. raro occurrit.

Astragalus sylvaticus. Wood or Heath peas, G. 1237. in the Woods and Heath at Hamsteel

Atriplex olida, Stinking Orach, G. 327. Atriplex olida five fylv. fætida, P. 749. in Suburbiis Londini.

\* Atriplex marina latifolia tota rubra, plentifully neer Shoram in Suffex. Mr. Brown.

Atriplex marina semine lato, ND, neer Har. lacham. Ph.

Atriplex sylvestris vulg. G. 326. Atriplex vul.

gatior finuata major, P. 748.

\* Atriplex marina Kalifolia, At Boston Lincoln-Shire, in the Road from the Bridge to the Wash.

Atriplex sylvestris latisolia sive Pes anserinus, Goof foot, G, 328. P. 746. in simetis Lond.

Atriplex sylvestris angustisolia, G. 326. P. 748. Atriplex sylv. altera, G. 326. Atriplex vulg. sylv. sinuata minor P. 748. small wild Orach, icona non congruunt.

Atri-

Atriplex Sylv. five Polyspermon, All-feed Orach, G. 324. Blitum Polyspermon, P. 754. Atriplex fylv. latifolia altera, G. 328.

Atriplex sylv. tota rubra, about St. Giles in

the Fields.

ph. five nile

184

lik di.

G.

G

ed,

ri

b.

į.

\* Atriplex Autumnalis facie Mercurialis, flore

parvo luteo, in St. Georges fields.

Avena vesca, Common Oats, G. 75. P. 1134. Avena caudata Cambro britannica, Commonly fowed in Wales, and there call'd Tailcorn.

\* Avena nuda, G. 75 . P. 1134.

Avena Nigra Casp. & Joan Bauhini.

\* Avena Carula, Blew Haver, feritur ad Kighly

in York Bire.

Auricula muris pulchro flore, Monfe-ear. with flowr's, almost on every dry Bank about Cambridge. CC.

Auricula muris vid. Pilofella.

### B

B Accharis Monspel. Plowmans Spicknard, G. 792. P. 114. in sylvis & montosis.

Barbaraa, Wintercress, G. 243. Barbaræa simplex, P. 820. ad agrorum margines.

Bardana major, Great Burdock, G. 809. Bardana vulg. major, P. 1223,

Bardana minor seu Zanthium, G. 809. P. 1223. In a bog beyond Peckham.

\* Battata flore rubro albo & cinericeo, Potatoes with white, and Ash-coloured flowr's, planted in many fields in Wales.

Behen album, G. 678. papaver spumeum, Spatling Poppy, P. 263.

\* Behen album fl. amplo staminibus purpureis foliisq; magis hirtis, Betwixt Sandgate Cafile, and Hide in Kent, and at Bass Castle, Cornwall.

\* Behen album hispidum, hairy spatling Poppy,

plentifull about Oxford.

\* Behen flore albo elegantion, three miles from Dover, in the way to Rye, on the Beach, all along betwixt Hide and Rumney.

Bellis major, great Daysie, G, 634. Bellis ma-

jor vulg. five fylv. P. 528, in pratis,

Bellis minor fylvestris, G. 636. Bellis fylv.

minor fimplex, P.530.

\* Bellis flore herbaceo globoso, in Mr. Seldens Cops neer his house in Worcestershire, Mr. Morgan.

Berberis, the Barbary bush. P. 561. spina acida

five Oxyacantha, G. 1325. in sepibus.

Beta alba, White Bete, G. 318. in maritimis.
Betonica, Betony, G. 714. Betonica vulg. flore
purpureo, P. 614. Variat flore albo, passim
in sylvis, & locis siccioribus.

\* Betonica flore carneo faturo, in the Cow-

pastures at Garforth Yorkshire, Mr. Witham.
Betonica Aquatica, G. 715. Betonica aquatica
major, P.613. Hac omnino annumeranda
scrophulariis, in stagnantibus passim obvia.

Betula, the Birch-tree, G.1478. P. 1408. in

fylvis.

n.

•

Betulus five Carpinus, the Horn-beam tree, G. 1479. Oftrys five Oftrya Theophr. P. 1406. On the West of Primrose-Hill, and Hamsteed Woods.

Bifolium vulgare sylv. sive Ophrys, Ordinary Twayblade or Bifol. P. 504. Ophrys bisol.

G. 403. in pratis humidis.

\* Bifolium minimum I. B. l. 31. pag. 534, neer the Beacon on Pendle Hill in Lanca-fire.

Bistorta major, Snakiwood. G. 399. Bistorta major vulg. P. 392. neer the Horse Ferry at Battersey, in the Meadows.

Bistorta minor, G. 392. P. 392. in pratis hu-

midis umbrosis, at Crosby in Westmoreland, and Ravensmaith; and two miles from Setle Torksbire, in the pastures towards Inglebo

rough hill in a place called Cromock.

Blattaria Plin. Yellow Moth Mullin. G. 776. Blattaria flore luteo, P. 64. in locis siccioribus. Variat flore albo. Between Deptford and Greenwich.

Blitum minus album, the lesser white Blite. G. 321. Blitum album minus syl. P. 753.

\* Blitum minus rubrum, the small wild Blite. G.
321. Blitum minus rubrum sylv. P. 753. In
some Ofier holts by Cam River, and in a
Ditch on the back of St. Johns Colledge, in
a Close on the North of the back gate. CC.

Bonus Henricus, English Mercury, or Good King Henry. G. 329. Lapathum unctuosum,

P. 1225. paffim in locis incultis.

Brassica marina Anglica, Sea Colemort, G.315.
Brassica marina monospermos, P. 270. By
Hastings in Sussex plentifully.

Braffica sylvestris. G. 316. P. 269. ad sepes.

Bromos sterilis, wild Oat, G. 67. Bromos herba five avena sterilis, P. 1148. in satis.

Bromos altera G. 76. Avena sterilis minor. P. 1148. Drank or Small mild Oats, secalis est vitium inter alias segetes rarissime invenitur.

Bryonia alba, white Brionie. G. 869. Bryonia alba vulg. P. 178. in sepibus.

Bryonia nigra G. 871. Bryonia fylv. nigra P. 170, cum cognomine suo.

\* Bryonia nigra fine fructu, G. 871. In the Pine walks going up to Hamsteed.

Buglofum luteum, Langue de bæuf. G. 798. ad

agrorum margines.

\* Buglossum parvum longifolium minimo flore, At Rumney going into the gate by the Coney Boroughs, An Borago minor fylv. Col. 183.

Buglossa sylvestris minor, small wild Bugloss, G. 799. Buglossum sylv. minus alterum, P.

765. in arvis.

d

,

ft i-

1

P.

7-

Bugula, middle Confound, G.631. Bugula vulg. P. 524. in humidis fylvis & pratis, variat flore albo.

\* Bugula flore carneo, several plants of it grow in a Close called the wood-close, belonging to my Fathers house at Mothley, Mr. Witham.

Bugula sive Consolida media Cærula Alpina C
B. Mountain Bugle or Sicle wort, on Carnedh

Llewellyn in Wales, P. 525.

Bulbocastanum minus, small Earth-Nuts, G. 1064. Nucula terrestris minor, P. 893. in the meadows neer Maribone Park and Hamsteed.

Bunias five Napus fylv. P. 865. fylv. Lob. G. 235. Our wild Navem, ad agrorum margines.

Bupleurum angustisolium Monspeliense, narrow leav'd haresear, G. 608 Bupleur minimum, P. 578. betwixt Bromeley and Eltham in Kent, and at Paddington beyond the Bridge in the way to Harrow upon the Hill.

Bursa pastoris, Shepherds purse, G. 276. Bursa D pastopastoris major vulg. P. 866. ubiq; magna est varietas hujus plantæ quoad ætatem, magnitudinem, hujus folia quandoq; magis minuswe inciduntur, quandoq; verò integra reperiuntur.

Bursa pastoris minor & minima, cum priore.
Buxus, the Boxtree, G. 1410. Buxus arbor vulg.
P. 1429. On Box hill in Surrey.

C

C

C

Ca

#### C.

Calamintha aquatica, mater Calament, G. 684. Calamintha arvensis verticellati five aquatica Belgar. Lob. P. 37. in locis humidis.

Calamintha odore pulegii, G. 687. P. 36. in collibus.

Calamintha vulg. officinarum, G. 687. Calamintha vulgaris, P. 36. common Callaminth, in Saxonis.

Calamintha montana vulg. G. 687. Calamintha minor in cana, P. 36.

Calceolus Marix, our Ladies slipper, G. 443. l. 217. in Helk woods in Lancashire.

G. 817. Caltha palustris vulg. simplex, 1. \*

\* Caltha palustris flore pleno, in divers place in Lancashire, G. 817. 2. Came Camelina, Treacle Wormfeed, G. 273. Camelina five Myagrum alterum amarum, P. 868. about a mile from Redding.

Campanula minor alba five purpurea, little white, or purple Bell-flower, G. 452. in are-

nosis collibus.

Campanula rotundifolia, round leav'd Bellflower, G. 452. Campan, minor fylv. rotundifolia. P. 651. cum priore.

Campanula Cymbalaria foliis, G. 452 .7. P. 652. 16. in Cornwall and Montgomery Shire.

and in Bagly-wood neer Oxford.

Cannabis mas, male or steel hemp, G. 780. Cannabis fativa, P. 598. præftantiffima habetur quæ feritur ad Brudeport in Surrey.

Cannabis fæmina, G. 708.

ta

ŋ.

in

th,

2

ld,

P

CE

Cannabis Spuria, wild Hemp, G. 709. P. 599.

ad fepes & in agris, variat flore albo.

\* Cannabis spuria flore albo magno elegante, along the Ditches from Scrooby to Shir Wood Forest. 2

\* Cannabis spuria flore albo magno staminibus luteis. In the Corn fields in the way from Carnarvon to Bangor, on the top of the Hill, two miles before you come to the Town.

Cannabis spuria flore pallido, labro purpureo elegante fair flower'd Nettle Hemp, in fimetis & ruderibus. N. D. At Marish in Cambridge-Shire, and in Kighly Torkshire plentifully.

Cannabis spuria humilis caule penè ad summitatem foliis viduata. In the Corn fields in

Surrey.

D 2

Ca-

Capillus Veneris, vide Adianthum & Trichomines.

Cardamine, Cuckow flower, or Ladie mock, G. 259, Cardamine altera minor, P. 826. ubiq.

in pratis humidis.

\* Cardamine seu Nasturtium aquat. amarum, bit ter water Cresses. prod. 45. in a bog betwin the Duke of Norfolks garden, and Lambul Church, in the way by Thames side, and in Cornwall.

Cardamine pumila Bellidis folio Alpina. G.260 plantula Cardamines alterius æmula Clus. P. 828. On the Rock neer the Quarry by Bath in variis locis juxta Rippon ad monasterium ejus, and in Denbighsbire.

Gardamine seu Sium minus impatiens, Impatien Ladies Smock, G. 260. Sium Nasturtii sylv folio, P. 1241. In Rils and Ditch banks!

bout Bath.

\* Cardamine minor laciniatis foliis, P. 826.5 fine icone, An Nasturtium aquat, minus prod 104. 4. Hanc ad me misit, Dr. Gunthorpe Cornubia.

Cardiaca, Motherwort, G. 705. P. 42. 0

feminibus projectis.

thiftle, Carduus Tomentosus Anglicus, l. 980 in montosis sed non multum frequens.

Cardnus Lanceatus, Spear, or the greater brown Spear thistle, G. 1174. latifolius sive major, l. 992. in pratis. Datur & angustifolius lanceatus.

\* Car

\* Carduns Lanceatus flore & capite minoribus, On the Ditch fides beyond St. James's.

duus Mariæ vulg. P. 976. ad agrorum margines.

\* Carduns Maria hirfutus non maculatus, in a Ditch neer my Lord Southamptons house,

and about Islington.

ma-

G.

big

bit

VIX

beth

din

60

. 1.

ath,

ien ylr

151

.5.

od

es

cl

de

044

an

47

Cardnus mollis flore caruleo, An mollis foliis Lapathi? G. 1184. on the highest Rock of Snowdon, and in Brearcliff neer Brunly Lancashire.

Cardum moscatus. G. 1174. Carduns Avenarius

seu moscatus, P. 959. in montosis.

Carduns folftitialis, St. Barnabies Thiftle, G. 1166. Carduus folftitialis Lob. P. 989.

Cardum stellatus, betwixt London and Mile end plentifully, G. 1166. Carduus stellatus sive Calcipatravul, P. 689.

\*Cardum spinosiffimus altissimus beath Thiftle.

Cardum palustris, the Marso Thiftle, P. 958.

An palustris caule crispo subrubente. CC.

Carduns Polyacanthus, Thiftle upon Thiftle, G. 1173. P. 982. ad agrorum margines, variat

flore albo ut Cardui plerique.

Carduus vulgatissimus viarum, G. 1173. Ceneanathos seu viarum & vinearum repens. P.

959.

Carlina acaulis minor flore purpureo, G. 1158.
Carlina fylv. Septentrionalium, P. 969.
Dwarf Carline Thistle, in montosis & collibus siccioribus.

D 3

CAYO

Carlina sylvestris major, G. 1159. P. 969. cum

priore.

\* Carum five Careum, Caramays, allata for mihi femina ex agro Lincolniensi ibidem nan quæ nullatenus cedebant Germanicis, G. 1034. Carum vulg. P. 910.

Caryophillata, Avens, or Herb Benet. G. 994. Cariophillata vulg. P. 136. ubique ferè in

sepibus.

Caryophillata montana, G, 994. P. 136. com-

mon in the woods of Lancashire.

\* Cariophillata mont. flore pleno, At Brear-elif, in a wood of Mr. Brearcliffs, below his house, An Caryophillara vulg. flore major C Cap.

Caryophillus montanus purpureus, wild Purple jagged Pink, G. 593. Neer Nottingham, betwixt the Town and the Gallows, Maker

Stoneboufe.

Caryophillus pratensis, G.594. Caryophillus pratensis noster major, P. 1338. 2. sine icon Deptfort Pink, in the Meadow on this fide Deptford, and in a Lane neer Early Heath by Redding, and in Tuddington field.

Caryophillus marinus minimus Lob. G. 602, gramen marinum minimus, P. 1279. Thrift,

in paludibus falsis.

Caryophillus faxatilis Erica fol, umbellatis corymbis C.B. Rock Pinks, with Heath-like leaves, P. 1341. I. Saxifraga alfinefolia major, Great Pearlwort, Ph. in montosis & alperis. In the middle way betwixt Lipport and Peters field.

Carj-

Caryophyllus minimus muscosus nostras, our smallest moss-like Pink, or small Pearl wort, P. 1340. Hujus duas species in horto meo sicco servo ex Cornubiz montibus allatas.

G. 591. On the Castle Walls at Roche-

fter.

mus

fant

lata

G.

94.

m-

if,

ie,

ap,

ple

8,

cr

2

ne

de

th

Castanea, Chesnut tree, G. 1442. P. 1400. in sylvis quibusdam & viridariis præsertim in Essexia.

Catanance seu Ervum sylvestre, Crimson Grass vetch, G. 1249. P. 1079 ad pratorum &

arvorum margines sed rariús.

Cancalis apii foliis flore rubro, G. 1021. Caucalis Angl. flore rubente, P. 920. Bastard parsley, with red flowers.

Cancalis altera purpurascens foliis latioribus & crassioribus semine etiam majore, Lob. Ad-

vers. An Caucalis major, Ph.

Caucalis minor flosculis rubentibus, G. 1022.

Caucalis minor flore rubente, P. 921. ad fepes.

Cancalis nodosa echinato semine, G. 1022. P.

921. ad agrorum margines.

\* Cancalis tenuifolia minor flore cæruleo, at a Chalk hole at Cheme in Surrey.

Caucalis albis floribus, Hedgehog Parsley with white flowers, G. 1021. Caucalis vulg albis

flor. P. 920. inter segetes.

\*Caucalis seu Echinophora tertia Acarique Purpurea Col. pag. 67. Caucalis Echinophora major & minor purp. Col. P. 920. plenci-

D 4 fully

fully in the Corn fields, about Banbury, Ox-

ford, and neer Slough Middlesex.

\* Caucalis minor semipedalis, G. 1023. huju tantum meminit sub finem descriptionis quintæ, Amongst Wheat plentifully neer Peters sield, Mr. Goodyer, who call'd it Caucalis pumila segetum.

Cauda maris, Mouse tail, G. 426, P. 500. in

Chelfey Meadow.

Centaurium parvum, G. \$47. Centaurium minus vulg. flore rubro & albo, P. 272. Small Centory, in montosis pratis & arvis.

Centaurium luteum, G. 547. P. 272. plenti-

fully on Roe-hill in Kent.

Gentaurium luteum perfoliatum, G. & P. ut fupra, in montosis & siccioribus pratis.

Cerasus avium nigra & racemosa, Birds cherry, or Black grape Cherry tree, G. 1504 P. 1517. in the North Hedgeberry tree.

Cerasus nigra, G. 1505. In several places in

Kent.

Cerasus sylvestris minimo fructu cordisormi, neer Stock port, and in other places of Cheshire, called by the Country people the Merry tree, Mr Stonehouse.

Cerasus sylvestris, the wild Cherry tree, in a Wood by Bath, and plentifully not far from

Peters field.

Cerefolium fylv. Hedge Chervel, G. 1038. Mir-

rhis fylv. P. 935. passim ad sepes.

Chamacistus vulg. flore luteo sive Heiianthemum Anglicum luteum & album, English yellow pellow and white Cistus, or little Sun-flowr, in montolis, & ficcis, P. 655, G. 1282. in montibus Surriæ frequens.

Chamadrys sylv. Wild Germander, G. 657. Chamadris spuria sylv. P. 107.11. variat

flore albo in pratis ubique.

ijas

onis

eer

tu-

in

ni-

all

i-

ıt

n

Chamefilix marina Anglica five Filicula petræa fæmina, female dwarf Stone Fern, G. 1143. on the clifts west to Hastings Sussex abundantly.

\* Chamapeuce Turneri part. 1. pag. 129. Dr. Turners Dwarf pine, quam ille bellè depingit, on a very high hill in the North call'd Boulfer.

\* Chamapence feemina seu polyspermos, At Dovestones in the side of Yorkshire, neer widdop, and a mile on this side Lippock in Hamp-shire.

Chamalinum perpusillum, Dwarf wild Flax, G. 559. betwixt Highgate and Barnet, and by the Rivers side neer Exeter.

Chamemelum nudum odoratum, G. 754. Chamemelum nudum, P. 86. In Tutbil fields.

chamamelum, Chamemill, G. 754. Chamamel. vulg, P. 85. in pascuis quibusdam, In the waste ground on this side Hide-Park, and on Hunston Heath every where.

Chamanerion, Rose-Bay, Willow herb, G 477. in sylvis quibusdam humidis, at Greenwich in

the place where Balast is taken up.

Chamenerion angustifolium, Narrow leav'd Willow herb, G. 477. P. 548. on the walls of Comb. Fark neer Ham.

nostrati Spencer, Terebinth, 6.525. Chamæpith. vulg. P. 283. in agro Cantiano.

Chelidonium majus vulg. great Celandine, P. 617. Chelidonium majus, G. 1096. ubique

in sepibus.

Chelidonium minus, Pilewort, G. 618. P. 617. in pratis humidis.

Chelidonium minus flore pleno, In Devonshire, & at Malmsbury Wiltshire, P. 617. 4.

Chryfanthemum segetum, Corn Marigold, G. 743. Chryfanthemum segetum nostras. P. 1369 in plerisque satis in quorum quibusdam nimis luxuriat, segetemque nutrimento spoliat.

\* Chrysanthemum flore pleno, As you go out of Nonesuch Park, towards Letherbead in

great Cornfield.

Cichoreum sylv. wild blew Succory, G. 284.P., 776. in arvis, variat flore albo sed rarissime.

Cienta, Hemlock, G. 1061. Cicuta vulg. ma-

jor, P. 932. ad plerasque sepes.

Gicutaria tenuifolia, thin leav'd wild Hemlock, G. 1063. Cicuta minor seu fatua, P. 933. cum priore.

Cicutaria palustris, G. 1063. Cicutaria palustris tenuifolia, P. 933. in locis paludo-

cicutaria alba, G. 1038. 6. Cicutaria palustris alba, P.932. 4. sine icone, Hist. Ludg. 761. optima icon.

Circaa Lutetiana, Enchanters Nightshade, G.

351. Circea Lutetiana major, P. 351. In locis humidis & umbrosis.

Cirlium montanum Anglicum, fingle beaded Thiftle, G. 1183. P. 961. On Barns common.

\* Cirfium altiffimum fol. latioribus & albidioribus, In several places of wales, and on a Moor two miles East from Croydon, where

they dig peat.

ha-

P. ue

17.

re,

G.

P.

m 10-

at

12

P,

ć,

a.

ŀ

5

.

\* Ciffus, mas breviore folio, P. 659. Ciffus mas dentatus, G. 1276. Small leav'd male Ciftus, Accepit Dr. Bowles infignis Boranicus mihiq; communicavit, ex Cambro-Britannia.

Clematis Daphnoides minor seu Vinca Pervinca minor, Periwincle, G. 894. Vinca Pervinca vulg. P. 380. in umbrofis. On the West of Hamfteed Heath plentifully.

Clinopodium minus five vulg. P. 21. Ocymum fylv. wild Basil, G. 675. in dumetis, variat

flore albo.

Cochlearia vulg. Common Scurvy-gras, P. 285. Cochlearia Britannica, G. 401. in maritimis & Thamisi fluvio-

\* Cochlearia longiori & sinuato folio, On the banks of Thames over against Blackwall.

Cochlearia rotundifolia, Great round leav'd Scurvy-grass, G. 401. Cochlearia rotundifolia five Battavorum, P. 285. Plentifully in the Peak of Derby-faire, and on the Rocks in Cornwall.

Cochlearia rotundifolia minima, Small round leav'd leav'd Sourvy-grafs, in Anglesey, P. 286;

fine icone.

\* Colchinum Anglicum purpureum ut etiam flore albo fed rarius, Meadon Saffron, in Occi. dentalis Angliæ partibus in pratis humidioribus, G. 157.

Colchicum Anglicum duplici ferie petalorum.

\* Colchic am Angl. faturatioris purpura.

\* Colchicum Anglicum petalis ex albo & purpureo dimidiatim mixtis, hæc tria Colchica cref cunt in prato juxta Comb. in agro Oxonienfi, Mr. Brown.

Conferva, River hairy weed, G. 1570. & Alga

aquatilis capillacea, P. 1261.

\* Conferva Plin, setis porcinis similis, Below Charlton in Kent, in the Marsh Ditches neer to Thames.

Consolida major flore purpureo & albo, Comfry with purple and white flowers, G. 806. Symphytum majus vulg. P. 523. in locis humidis.

Consolida major flore rubro, By Redding, and

two miles Eastward from Arundel.

Consolida regalis sylv. Larks beel, G. 1083. in a field by Pershore in Worcester Shire, Ph.

Convolvulus major albus, Great white Bindweed, P. 163. Similax lenis five levis major, G. 861. in sepibus.

\* Convolvulus major flore carneo, At Somerfet Bridge end in the way from London to

Peters field.

Convolvulus minimus spica foliis, G. 862. Convolv. spica foliis, P. 172. In the fields about

great

6:

0-

ci.

ti-

.

u-

W

great Dunmow in Esex, and in Inddington field.

Convolvulus minor purpureus, P. 172. Smilax lenis minor, G. 861. in agris, variat flore albo & rubro.

Convolvulus niger, volubilis nigra, G. 863. Convolvulus niger Atriplicis folio. P. 172. inter segetes.

Conyza major vide Baccharis Monspeliensum.

Conyza media, Middle Flea wort, G. 402. in locis uliginosis.

Conyza minima, G. 482. in aquosis cum priore.

Conyza Carulea acris, G. 484. Conyza odorata carulea, P. 126. in montofis pratis, On Parsons green on the Ditch banks plentifully.

Conyza foliis laciniatis, G. 483. Conyza Helenitis pilosa fol. laciniatis, P. 126. In the lse of Ely, about Marsh and Chatteress, C.C.

\* Conyza acris flore albo, plentifull at Heddington hill by Oxford, and in Yorkshire, Mr. Witham

\*Conyen palustris, P. 1232. In the Ditches betwirt Wisheech, and by Gyhorn in the lse of Ely.

\*Corallina comis ad instar caudæ vulpinæ sparsis. Sheeps tail'd Coralline, found cast upon the Beach betwixt Margate and Dover Castle.

Corallinm album, white Coral, G. 1576. Corall.

album majus & minus, P. 1300. in cautibus maritimis.

\* Coriandrum vulgare, Coriander, G. 1012. P. 2918. fatum erat in Essexia, unde allata mihi

semina Germanicis paria.

Cornu Cervinum, Bucksborn, G. 427. Coronopus vulg. five cornu Cervinum, P. 502. passim juxta vias.

Coronopus Ruellii, Swines cresses, G. 427. Co. ronopus repens vel recta Ruellii, P. 502.

Corallina Anglica, English Coralline, G. 1571. Muscus marinus squammatus sive Corallina squammata, P. 1296. adnascitur testis piscium & saxis.

Corallina alba sive Muscus marinus, White Coralline, or Sea-mols, G. 1571. P. 1294. in.

ter scopulos.

Cornus famina , Dogberry-tree , G. 1467. P.

1521. in dumetis & fepibus.

Corylas fylv. Wild Hedge Nut, G. 1438. Corylus five Nux avellana fylv. P. 1416. in

fylvis & fepibus.

Cotula fætida flore pleno, G. 757. P. 87. In the Isle of Thanet, and betwirt Gillingham and Chattam, in a close adjoyning to the Marsh.

\* Cotula non fætida flore pleno, G. & P. ib. In St. James's field on the upper side neer the Highway.

Cotala fætida, Stinking Maymeed, G. 757. P.

87. in agris.

Cotula non fætida, Dogs Chamemile, P. 87.

Coty-

Cotyledon palustris, Water Penny wort, G. 530. Cotyledon acris septemtrionalium, P. 1214. in uliginosis, On Hamsteed heath in the bogs.

Coyledon hirfuta, Hairy Kidney wort, on the moist Rocks of Snowdon, and several Moun-

tains of Wales.

in sylvis & inter segetes, At Waltham Abby in Hartford-shire.

Crateogonum cubitalis altitudinis flore luteo, In the life of Wight, Mr. Cole, and in the Kings

meadows at Godffone in Surrey.

Crataogonum Euphrosyne, Eye-bright Com wheat, G. 92. Euphragia pratensis rubra major, P. 1329. in pratis & inter segetes, storis color nunc intensius aliàs dilutius rubescit.

Crista Galli, Tellow rattle, or Coxcomb, G.
1071. Pedicularis sive Crista Galli mas, Ger.
1071. P. 713. ubique in pratis, variat flore
albo.

Crithmum marinum, Rock Sampire, G. 533.
Crithmum mar. vulgatius, P. 1286. On the clifts of the Rocks, in most parts of England.

Crithmum Chrysanthemum, G. 533. P. 1287.
Crithmum spinosum, G. 533. Cirthmum maritimum spinosum seu Pastinaca marina, P. 1286. By Lee in Essex, ut Ph 'tis not found there, but below Feversham.

Crocus sativus, True or common Saffron, seritur in agro Cantabrigiensi & Essexia, G.

ISI.

Cruciata, Crofs Wort, G. 1123. Cruciata vulg.

P. 566. in Hamfteed Church yard, & inoc.

cidentalibus partibus in dumetis.

Cuscata five Cossutha, Dodder, G. 577. P. 10. in agris & Ericetis, circa varias plantas, tik. Ericam, genistam lupulum, urticam, sabis quas (nisi avellatur) strangulat, hinc Epierica, Epigenista, Epiurtica, & c.

Cyanus vulg. common blew Bottle, G. 73 2. Cyanus minor vulg. diversorum colorum, P.982, inter segetes, coloris sc. cærulei, albi, purpurei, in the corn fields neer Cambridge.

Cyanus flore caruleo medio purpureo, In Stone

bersley field Yorkshire. Mr. Witham.

Cyanus flore albo medio caruleo. In the field by Hocley in the Hole, neer Oxford, Mr. Witham.

Cynocrambe, Dogs Mercury, G. 333. Cynocrambe vulg. P. 296. in fylvis & fepibus.

Cynoglossum majus vulg. Hounds tongue, G. 804.
P. 511. passim ad agrorum margines, variat flore albo.

Cynoglossum minus folio virente, G. 805. Cynoglossum semper virens, P. 512. At Norbury in Surry, a mile from Letherhead plentifully.

\* Cynosorchis minor Pannonica, the lesser Austrian dog stones, G. 207. Cynosorchis militaris Pannonica, P. 1345. On Chiswell hills neer

Oxford, and neer Hereford.

Cyperus longus inodorus, unsavory Cyperus With long roots, P. 1264. On Hinton Moor in Cambridge shire, and several Fenny places.

Cype-

00-

10.

bas

Ca,

ya-

32,

tt.

ld

1.

peroides miliaceum C B. 90. In the Meadows on the East-side of Hampsted heath, and in the North plentifully.

\* Cyperus Typhinus, P, 171. At the farther end of the Neat house Garden in the way to Chelly.

\* Cyperus repens rad. longa unicoq; capite, Betwist Sandwich and Deal, ad littus maris.

# D.

Dans vide pastinaca.

Dens Leonis, Dandelyon, G.290.P.78c.
est magna varietas foliorum hujus plantæ, sed
non tanta ut inde novæ species constituantur.

Dentaria, Tooth wort, G. 1585. P.1363. In a Wood three miles beyond Croydon, near Wad-dington towards the Down:

Digitalis purpurea, Purple fox glove, G. 790. P. 653. in montofis, variat flore albo.

Disfacus sylv mild Teasel, or Venus basin, G. 1167. P. 984. ad agrorum margines.

Diplacus minor five virga pastoris Sheeheards rod G. 1186. P. 984. ad sepes & vias regias sed rarius, In a lane by Depiford, and beyond Fulham in the way to Hammer smith.

Dorenicum majus Officinarum, great Leopards bane, G. 759. P. 319. On the high mountains

of Northumberland.

E

Draba

Draba lutea, Yellow Arabian mustard, P. 851, Draba quarta, G. 274.

Dryopteris Adv. true Oke fern , G. 1135. filir querna repens, P. 1041. in uliginofis.

Dryopteris Tragi, Tree fern, G. 1135. At the bottom of Pendle hills, towards the new Church

Dryopteris nigra five Onopeteris mas, G. 1137.
adianthum nigrum vulg. P. 1049 in umbross
& asperis, At Darkin Towns end, just over
the River.

\* Dryopteris fol. minutim incisis, On the Rock of the Fels, three miles from Hawks-head,

Lancashire.

\* Dryopteris marina, In the clifts of the Rock and shady places, half a mile West from Hafings, by the Sea shore.

Dulcamara, seu Amaradulcis, Bitter-sweet, G. 350. P. 350. in sepibus, variat flore albo.

\* Dulcamara marina, variat etiam flore albo, 44 littora Cantiana, Both forts grow plentifully at Shoram in Suffex. Mr. Brown.

## E.

E Bulus five Sambucus humilis, Danewort, a Danes-blood, or Dwarf-elder, G. 1426. Ebul. vulg. P. 206. in agris, sed rarius.

Echium vulg. G. 802. P. 519. In muris & los ficcis variat fl. albo, Without Frier Bacons Str dy, Oxford. \* Echim

\* Echium fl. carneo cæruleo & albo, In many places twixt Aylesbury and Evesham.

Echium alterum feu Lycopsis Anglica, Wall buglos, G. 802. P. 519. cum priore.

Elaphoboscum vid. pastinaca.

Elatine sive veronica semina Fuch. female sluellin, G. 625. Elat. folio subrotundo, P. 550. inter segetes & ad agrorum margines.

Elatine altera, G. 625. Elat. fol. acuminato, P.

553. cum priore.

Enulacampana five Helenium, Elicampane, G. 793. P. 654. In pratis & pascuis quibusdam. Equiseum majus, G. 1113. Equis palustre majus, P. 1200. Great marsh Horstail.

P. 1201. In Saint George's Fields in all the

Ditches.

51,

)Ot-

37. Ofi

Ver

ck

14,

cks

4.

G

40

lly

Equisetum lacultre foliis mansu arenosis seu Hippuris Coralloides, &c. in uliginosis.

In Pits on Hampsted Heath, and at Hammersmith near Sir Nicholas Crisps Brick-pit.

Equiserum segetale seu arvense longioribus setis G. 1114. P. 1202. 12. ubiq; inter segetes.

Equifetum fylvaticum, G. 1114. Equif. pratenfe majus, P. 1202. In Charlton woods, and the Meadows by Hampsted heath.

Equiserum palustre alterum brevioribus soliis, P.1203. Cauda equina famina, G. 1114.

\* Equifetum pratenfe longissimis setis, C B.247. betwirt wandfor and wimbleton, in the midway in the Meadows.

E 2 \* Equi-

\* Equisetum primum Math. quo utuntur arcusrii ad sagittas poliendas, At Skippon and Craven in Lancashire, and in Rigby Woods.

\* Equisetum palustre ramosum & aquis immersum seu mille solium aquaticum equiseti solium, in aquis pigrioribus, CC. On the back side of the Grange in the ditches beyond Southwark, and betwixt Limehouse end and Black Wall.

Erica baccifera tenuifol.G.1383. Erica baccife. ra nigra, P. 1485. Plentifully in Lancashin and Torkshire, and many other places.

Erica vulgaris five pumila, common heath, G. 1380. Erica pumila Belgarum, Scoparia no stras, P. 1482. fed non congruunt icones, variat flore albo, uti pleræque.

Erica major floribus purpureis, G. 1382. Eria coris folio maxima purpurafcens, P. 1481.

Erica tenuifolia. Erica vergata feu fexta Clufi, P. 1488. in feveral heaths in Surrey.

Erigerum vulg. common Groundsel, G. 278. h. 671. ubivis locorum sese offert.

Erigerum tomentorum alterum, G. 279 Sene cio feetidus, P. 672. ad agrorum margine.

Ernea aquatica, Water-rocket, G. 248. P. 1242. Ernea marina, G. 248. P. 821. ad littus maris in Cornubia.

Eruca fylv. vulgatior, G. 246.P.818.in filluis murorum.

Eryngium marinum, Sea holly, G. 1162. P.986. Erysemum vulg. seu Irio, Bank cresses, G. 254 P. 833. juxta semitas.

E[u|u

7

Essla exigua Tragi, small annual purge, G.503. Tithymalus leptophyllus, P. 193. inter segetes.

Esula ma jor Germanica, Quaksalvers Turbith water spurge, G. 501. by a Wood side a mile from Bath, and betwixt Guilford and Godliman near Compton, in a wheat field by the side of a Moor, near Mr. Yaldons house.

Esula minor seu Pithusa, G.502. In divers corn fields near Petersfield, Hampsbire.

Elula rotunda vid. peplus.

1.

ŀ

P.

78

0.

s;

ca

îi,

P.

16.

Š,

2.

ris

TB

ulı

Evonymus Theophr. Prick timber, or spindle-tree

G. 1468. P. 241. in sepibus.

Enpatorium Cannabinum fæmina, Water-hemp, or Water-Agrimony, G. 711. Eupat. Aquaticum, P. 595. in aquosis.

Eupatorium Cannabinum mas, common Dutch

Agrimony, G. 711. ad rivulos.

\* Enpatorium aquat. folio integro. In the ditches near the Bridge at Amdrey Causey, and many other places. CC.

\* Eupatorium Cannabinum palmare & angustifolium, In Surrey near to Sommer set-bridge,

in the Fish-pond upon the Moor,

Euphragia sive Euphrasia, Eye-bright, G. 663. Euphragia vulg, P. 1390. variat flore Caruleo in pratis humidis.

### F.

F Aba ma jor hortenfis, Great garden Beans, G.

Faba sylv. G. 1209. P. 1054. cum priore, plu.

rimæq; ejus species in arvis seruntur.

Fagus, The Beech, in fylvis & montibus, G. 1444
P. 1403 hujus folia in aliis arboribus magis,
in aliis minus inciduntur, magis, on the first
heath behind Redding, in the way to Oxford

Ferrum equinum, Horshooe, G. 1235. Ferrum equin. majus, P. 1091. On the hills about Bath, and between Bath and Marleboron.

Ferrum equinum comosum, P.1092. 4. Bushyheaded Horsbooe, vetch. On Gogmagog hill, and New-market heath, and in a Chalk-pit Close at Cherry-Hinton. CC. app.

Filipendula, Dropwort, G. 1058. P. 435. in

umbrosis.

Filipendula aquatica, G.1060. Oenanthe aquatica major, P. 1233. In fossis quam plurimis. Filicula petræa mas, G. 1142. in locis siccion

bus.

Filix mas ramosa pinnulis dentatis, Male ferm G. 1128. P. 1036. In umbrosis uliginosis. Filix sæmina, G. 1128. Filix sæmina vulg. I.

1039. cum superiore.

Filix florida seu Osmunda regalis, water fern,or

Ofmond Royal, G. 1131. P. 1038. in uliginofis.

Frequent in most parts of Cornwall.

\* Filix Dauci foliis. On old walls a mile from Hastenden, going over the hill towards Good-scham Chappel.

Filix fontana mas, G. 1142. Filicula fontana, P.

1044. 6.

G.

þ.

44 15,

1

m

ot

.

0

Filix mas non ramosa pinnulis angustis raris profunde dentatis, Rock male fern, G. 1130. in locis asperis & umbrosis.

Filix mas non ramosa pinnulis latis auriculatis spinosis, Pointed male fern, G. 1130, ad sepes

in locis faxofis.

\* Filix faxatilis crispa, P. 1686.ut opinor, Curled wall fern, over the Gate as you go into King-street.

Filix mas non ramofa pinnulis latis denfis minutim dentatis, G. 1129. ubiq; ad fepes.

flowers, G. 387, inter segetes. Plentifully in Kent near Roe-bill,

Fragaria fructu hispido, Prickly Strawberry, G. 998. 2. In Hidepark and Hampsted woods.

Fragaria minima vesca seu sterilis, G. 998. P. 758. 4. in sylvis.

Fraxinus, Afterree, G. 1472. fraxinus vulg. P.

1419. in sylvis & sepibus.

Fraxinus Bubula sive sorbus vulg. G. 1473. Ornus seu fraxinus sylv. P. 1419. the Quickentree, cum priore at rarior.

Fuem five Algatin coria, & muscus marinus,
Dying red sea moss, P.1289.near Bridlington on

4 the

Fuensmarinus rotundus, Sea points or laces, on the fouth fide of the life of Wight on the rocks.

Fuens polyschides, G. 1570. Fucus maximus polyschides, P. 1292. Sea girdles with many la-

bels.

Fucus spongiosus nodosus, Sea ragged staff, G. 1570. spongia ramosa altera Anglica, P. 1304

betwixt Sandwich and Deal.

\* Fucus phasganoides fol. indivisis penes me magna est variatas hujus Fuci quoad fol. magnitudinem, crassitiem & colorem, Quorundam insuper solia serrata aliorum integra, quandoq, crispa vel plana.

Fumaria purpurea, Fumitory, G. 1088.P.287.

in fatis & arvis.

\* Fumaria alba fine claviculis. In Pendle Forest Lancasbire, and at Crediton in Devonsbire.

Fumaria alba latifolia claviculata, G. 1088. Fumar. latifolia alba, P.288. in Charlton wood, and on Black-heath.

\* Fungi multi ex uno pede perniciosi, I B. t. 3.

834. ubiq; fere obvii.

\* Fungi multi ex uno pede Clypei formes variorum colorum per oras crenati vel non, I B. t. 3,836. In rusticorum tuguriolis vetustis putrescentibus.

Fungi esculenti & non esculenti, Mushrooms pojfonous, and not poysonous, G. 1580. P. 1316.

in pratis.

\*Fungi ex putrescentibus carnibus, &c. enati plurimorum generum, Anglice Mouldiness. Hos fungos fungos detexit sagacissimus vir Dominus Hook in suo singulari libro micrographico nuper edito, ibidemque belle depinxit. Quam plurimas alias sungorum species vidi, possideoque quas in præsentiarum omitto, quassam autem adjeci, quod non ita pridem inaudiverim prodiisse librum Romæ qui trecentas & ultra describit.

\*Fungus campani formis, niger, parvus multa semina plana in se continens, Mr. Morgans Garden Westminster, call'd in Wostershire Corn bells, where it grows plentifully.

\* Fungus corallii rubri colore, multis lineis nigricantibus masculisque luteis insignitus, P.

1321. apud Cambro. Britannos.

\* Fungus disci formis, tuberis species, betwixt Highgate and Hampsted in the valley, and on the downs of Suffex and Surrey.

\* Fungus Hepatis facie & colore, As you go from Sir Robert Dillingtons house to the new

Church in the Ifle of Wight.

\* Fungus luteus Sambucino similis, colore suo manus inficiens, Genista spinosa vulg adnascitur, Dr. Cumborp.

Fungus Oviformis, On a Moor betwixt Readlane and Sommerset-bridge in Hampshire.

\* Fungus pyriformis, In old Pastures not far from Yarmouth, on the North of the Isle of Wight.

Fungus pediculo bulboso innixus pallidus & maculatus, 1B, t. 3, 3,43. In some antient

Paftures.

5.

la-

G.

04

ne

n.

7.

f

d,

.

\*Fungus ramofus flavus & albidus, I B.t. 3.837. At Fulham Towns end in the way to Ham.

mer (mith.

\* Fungus rotundus supernè concavus & translucidus coloris fuccini in Mr. Morgans garden, & alter coccinei coloris, in St. James's Park in the Winter time on old decayed Trees.

\* Fungus semicircularis durus, multos durat per annos, originem debet pyro domestica &

fylv.

\* Fungus sterquilinii albus & niger perniciolus,

I B. t. 3. 845.

\* Fungus tertius quinti generis Clus. pag. 265, On a decay'd Willow on the North of St. No. ets near the Mill.

\* Fungus faviginosus, Hony-comb Mushroom, G. 1583.P.1371. Beyond Ditton in Cambridge fire, and on the bottom of an old Oak three miles short of Guilford as you go from Cob. ham, crescit ab Aprili ad Maii finem.

\* Fungus durus arborum five igniarius, Touchwood, P. 1323. Antiquis adnascitur arboribus,

\* Fungus pulverulentus seu tubera terræ, Fus, or Puff-balls, G. 1583. P. 1319. datur minimus, nucis magnitudine, coloris purpurei, la Hampton Court Park.

\* Fungus stelliformis, At Hampton Court below

the house near King Henry's gate.

\* Fungus scarlatinus calice amplo, ex absynthio marino post olei Chymici extractionem projecto in hortum Coll, Medd. Lond. & ibidem fepulto ortus

\*Fungus folidus ex betula & quandoq;ulmo, IB.
t. 3. 841. in plaga boreali, Birch ball, in Surrey Swans-ball, hoc Indunt pueri loco pilæ
palmariæ, Oritur ex putridis partibus dictarum arborum.

Fungus Ophioglossides, On graffie Ant-hills, in a Close next Hample-wood, Mr. Stonehouse, and at Comb-park in the way to Kingston.

Fungus Sambucinus five Auricula Ludz, Jews-ear, G. 1581. P. 1320.32. On old Alders.

\* Fungus clavatus, At Newington beyond Southwark, ex ligno putrido hortos vel areas muniente.

\* Fungus Cucurbiti formis magnitudine capitis humani & ponderofus candidusque instar nivis, Five miles from Peterburgh in the way to Gyhorn.

\* Fungus penis effigie albus fætidus. Hunc maturum devorat musca carnivora, raro inveni-

tur. In Lancasbire.

37.

47.

flu-

len,

ark

per &

us,

5.

10-

\*Fungus circulum gradatim perficiens, cujus diameter quandoque 30 vel plures pedes conficit. In montofis pascuis non infrequens.

\*Fungus Corallinus ad antiquarum arborum ra., dices, In the Woods near Peters Field Hampspire.

\* Fungus piperi Ethiopico similis, vel digitatus niger.

\* Fungus digitatus citrinus minor.

\* Fungus rotundus scarlatinus odoratus, At Church Lench Wostershire.

## G.

Alcopsis vera, Hedge nettle, G. 704. Galeopsis Legitima Diosc. P. 608. variat quandoque flore albo, sed rarissime.

Galium album White Ladies Bed straw, G. 1127.
In the fuz bushes on Bansted downs, and many

other places.

five mollugo vulgatior, P. 565. On wimble ton Common, and other barren places.

Galium luteum, Cheese rennet, and in Yorksbin Maiden hair, G. 1126. P. 564. Ubique in

pratis.

Galium rubro fiore. P. 564. Galium rubrum, G. 1127. Near Minhead in Sommer setsbire.

Genista spinosa vulg. Great Furzbush, G. 1319.

P. 103. ubique in locis sterilibus.

Genista, Broom, G. 1311. P. 229, variat flore albo; quandoque inter alias genistas.

Genistella aculeata, Needle furz, or petty whin, G. 1320. P. 1004. in Ericetis, sed iconas non probo.

Genistella tinctoria, Green weed, or Dyers meed, G. 1316. Genista tinctoria vulgaris, P. 229.

in pascuis & agrorum marginibus-

Gentiauella fugax minor, Bastara, or awarf selwort, G. 437. Gentian. Autumnalis Centauri minoris minoris folio, P. 406.in montofis, variat flore albo.

gentianella Alpina verna, G. 436. Gentian verna major, P. 403. In the Mountains betwixt Gort and Galloway, Mr. Heaton, Ph.

\* Gentianella Autumnalis longifolia, On the

Downs by Lewis in Suffex.

Gentianella minima Bavarica Cam. P. 407. By

Pontfract plentifully.

iat

ny

re

n

\*Gentianella autumnalis flore albo, fol. longis, angustis, In old Pastures on the North west of Church Lench Wostershire plentifully, multum variat Gentian. autumnalis quoad foliorum longitudinem latitudinem, florumq; magnitudinem & colorem dilutius profundius ve caruleorum.

Geranium Cicutæ folio inodorum, unfavoury field Cranes-bill, G. 945. variat flore albo. Geran. moschatum inodorum, P. 708- juxta vias.

Geranium Batrachoides, G. 942, in pratis humidis.

Geranium Columbinum, Doves-foot, or Cranesbill, G. 938. Geran. Columb. vulg. P. 706. fuper tectis & ad agrorum margines ficciores.

Geranium Columbinum fol, magis diffectis, G. 938. Geran. Malacoides laciniatum five Co-

lumbinum alterum, P. 7c6.

Geranium Columbinum fol. magis diffectis, pediculis longissimis flore magno, In several

places of Hampshire, Mr. Goodyer.

\*Geranium Columb fol megis diffectis, pediculis brevibus flore parvo, At Guilford Towns end in the way to London. Gera-

Geranium malacoides fol. dissedis minimum, p.

Geranium fanguinarium, G. 945. On Saint Vin.

cents Rocks, and Ingleborem hill.

Geranium Rupertianum, Herb Robert, G.939. Geran. Robertianum vulg. P.710.in umbro. sis & sepibus, variat flore albo.

Geranium sexatile, G. 938 P. 707. in muris antiquis & saxorum fissuris. Betwixt Branson

and Sion-house plentifully.

Vincents Rocks on the farther fide of the River, and at Bass Castle in Cornwall, and twixt Milford Haven and Haverford Well, on old walls by the way side.

Gladiolus palustris Cordi, Water sword flag, G. 29. Juneus floridus, Flouring rush, P. 1197.

Glandes terræ, peas Earth nut, G.1236. Lathyrus arvensis, sive terræ glandes, P. 1061.

Glastum fativum, Garden woad, G. 491. P.600

seritur in latioribus solis.

Glaux Diascordis, Diaschis, Milk-tare, G.1244. Glaux Hispanica Clusii, P. 1075. in monto-fis.

Claux exigua maritima, Black Saltwork, G.561. P. 1283. In the Downs. vid. Icon. Dod.

Glychyrhiza echinata Diasc. Hedge bogg licoria, G. 1302. P. 1099. Its planted in many places.

Glychyrbiza vulg.G.1302 Glychir. vulgarishliquata, P. 1099. in quibusdam sylvis, Common in Yorksbire.

\*Gnaphalinn

\* Gnaphalium vulg Cudweed, G. 639. filago minor, P. 686. in locis incultis.

Gnaphalium Angl. G. 639. On barren High-

waies, almost every where.

,P.

in.

olo.

an-

aint

Ri

be.

A,

G.

97. by

00

12,

0.

ce,

Dy

fi-

M-

K#

\*Gnaphalium montanum fl, rubro, & purpurascente, In great plenty mixed with the white, on a great Mountanous place on the lest hand of the way as one goes from Pontfrast to Wembersly, and almost directly opposite to Stapleton Town on the right hand, Mr. Witham.

Gnaphalium montanum album, White mountain Cotton weed, G. 640. In Scosby Leas near Don-caster, and Shirmood Forest, & in montibus Cambobritannicis, Gnaphalium montanum sive pes Cati, P. 691. variat store purpureo.

Gnaphalium marinum, G.640. Gnaphalium five cotonaria, P.687. on the sea Coasts of Wales.

Gnaphalium filago seu herba impia, Herb impions, or wicked Cudweed, G.642. Gnaphalium minus, seu herba impia, P. 685. in arenosis & sterilibus.

Gnaphalium odoratum flore albo elegans pusilla planta, On a Heath by Barneck, Dr. Bowles, Ph.

Gramen Alorecuroides minus, Small foxtail grafs, G. 10. P. 1167. Gr. Typhoides molle, CB. 50. Passim ad semitas.

\* Gr. Alopecuroides minus spica aspera brevi, P.

1168. in pascuis frequens.

\*Gr. Alopecarinum majus, G. 11. Gr. Typhoides culmo reclinato, CB. 53.

Gr. Alopecurinum minus, Small bastard foxtail grafs.

grass, G. 11. Gr. Alopecuroides spica longa majus & minus, P. 1168. Gr. Typhoides spica angustiore, CB. 53. in satorum sulcis.

"Gr. Anthozanthum spicatum, I B. tom. 2. pag,

466, locum non memini.

Gr. Aquat. spicatum, G. 13. Gr. marinum spice.

tum, P. 1278.

Gr. Aquat. G. 13. Gr. Aquat. alcerum, P. 1275, nihil aliud esse videtur quam gr. junceum so. lio articulato.

" Gr. Aquat. magis spar sa panicula, CC.

Gr Aquat.paniculatum Tab. seu Calamogrossis sylvæ Divi Joan. P. 1181. Red-grass of Saint Fohns Wood near London, and on the back side of Southwark towards Peckham.

Gr. Arundinaceum Aquat. Great water Rud

grafs, P. 1274.

Gr. Arundinaceum minus, G. 8. P. 1181. Gr. A.

rundinaceum spicatum, CB. 94.

Gr. Arundinaceum acerola gluma nostras, Ou great Reed grass with chassie heads, P. 1273. Ubique ad majorum sluviorum ripas.

Gr. Arundinaceum panicularum, wild Reed, G.;. Calamogroftis five Gr. Arundinaceum majus, P. 1181. Gr. paniculatum aquat. latifol. C B. 40.

Gr. Arundinaceum, Reed-grass, or bent, G.5.Gr. arvorum venti spica, P. 1158. Gr. segetum panicula arundinacea, C.B.5.

Gr. Arundinaceum torosa panicula, sive calamogrostis, Round tusted Reed-grass, P. 1182.

\* Gr. Arandinaceum 30. pedes longum, On the

South of the Isle of Wight, by the Sea fide to-

wards the point.

g.

1

۶. ٥.

int

ck

ed

A-

ut

3.

7.

12.

ol.

m

0-

he

ich

Gr. Avenaceum altero alteri innato, P. 1 50.12. Gr. Avenaceum capillaceum minoribus glumis, Small baver-grass, C.B. 155. 4. in pratis.

Gr. Avenaceum incanum, P. 1149.

Gr. Avenaceum innata longistima spica, Ph.

Gr. Avenaceum lanuginosum glumis ratioribus, CB. 157.

Gr. Avenaceum caule folissq, hirsutis, An Bro-

moides maximum hirfutum? P. 1149.

Gr. Avenaceum rariori grano nemorense Danicum, I B. tom. 2. 434. In Hampsted, and many other Woods and Ditches.

\*Gr. Avenaceum rariore gluma spicatum, Oatgrass of Denmark, P. 115143. C B. 155.

Gr. Avenaceum zerampelinum Danicum, P.

\*Gr. Avenaceum locustis rubris, Oat-grass with red husks, P. 152. gr. montanum avenaceum locustis rubris, CB. 1556

Gr. Avenaceum squammofa gluma, P. 1149. 5.

\*Gr. Avenaceum squammolis glumis ramosum; \*Gr. Brize spica minus, C. B. 133. gr. angustifol spica tritici mutici, P. 1144.

Gr. Brize Spica majus, G. 29. 5.

\* Gr. Bromoides latiore panniculâ, P. 1 1 50. Sex pedes longitudine sua æquat. Plentisul in the Woods below Hampsted.

Gr. Bulbosum Aquat CB.21.P.1249. Est fagittariz species, In several places by the Thames

banks near London.

Gr. Bromoides copiosa pannicula. Festuca graminea essusa juba, C B. 144-3.

Gr. Bromeides vernum spicis erecis, Early

meadow arauk.

Gr. Caninum, Couch-grafs, or Dogs grafs, G.23, Gr. Caninum vulgatius common Quich-grafs, P. 1173. gr. Caninum arvense seu primum, seu gr. Diosc. & officinarum, C B. 7.

Gi

\*0

G

0

Gr. Caninam nodosum Avenacea pannicula, CB. 18. gr. Caninum nodosum, G. 23. gr. bulbosum nodosum vulg. P. 1176. summita non bene depingitur nec à G. nec P.

Gr. Caninum marinum spica siliginea, g. Caninum longius radicatum, P. 1 173. ubi beli

habetur icon summitatis solius.

\*Gr. Caninum nodosum duobus nodis major femper superimposito minori. Below Brish in the Meadows on the North of the River by the Ferry.

Gr. Caninum supinum, Creeping doggs grass, G

26. P. 1147. CB. 13.

\*Gr. Caninum Avenacea pannicula non nodosum ad agrorum margines. Frequent in Middle sex.

Gr. Caninum supinum marinum alterum, G. 25.
gr. Caninum maritimum alterum longus
radicatum, P. 1277.gr. Caninum maritimum

spica foliacea, C B. 15.

Gr. Caninum supinum longissimum, ND. two miles from Salisbury, by Mr. Tuckers & Maddington wherewith they fat Hoggs, and 'tis 24 foot long, and in some places of Walu,

h

re fol er

m

ius

\*Gr. Caninum marinum ad aliquot cubitos extensum, allatum est mihi ex littore Contiano.

Gr. (yperoides, Cyperus grafs, G. 12. gr. majus latifolium, P. 1266. gr. latifolium spica russa seu eaule triangulo, C. B. 83. passim in pigris & stagnantibus aquis & pratis etiam udis.

Gr. Cyperoides angultifolium majus, G. 29 gr. fpicis longis erectis, C B. 84. gr. minus angu-

flifolium, P. 1266. in fylvosis humidis.

\*Gr. Cyperoides albis glumis, on a bog a mile from Sandy Church in the way to Cobham in Surrey.
\*Gr. Cyperoides angustifolium pannicula multiplici CCapp.

Gr. Cyperoides fol. caulem ambientibus. Within two miles of Manchester, in the place where they dig for Marl in the way to Rochdale.

Gr. Cyperoides maximum spicis pendulis, at Bocknam in Surrey, in a bog. Hujus radicibus utuntur pro sedilibus in agro Eboracensi.

Gr. Cyperoides palustre, Great Cyperus grass, G. 21. gr. Cyperoides palustre majus, P. 1266. gr. palustre majus spica compacta, CB. 87.

Gri Cyperoides palustre minus, P. 1267. gr. Cyperoides parvum, G. 21. gr. palustre

majus spica compacta, CB. 88.

gr. Cyperoides aquat. vulgatius, G. 32. gr. Cyperoides palustre pannicula sparsa, P. 1266 gr. pannicula sparsa majus, C B. 86. in humidis paludosis canosis rivorumque margini. bus.

Gr. Czperoides spicatum, G. 22. gr. foliis Caryophylleis spica squammata, P. 1160. variat spica divulsa & simplici, In a bogg a mit

Westward from Oxford:

Gr. Cyperoides spica pendula breviore, CB.87, pseudocyperus, G. 29 Cyperus sive pseudocyperus spica brevi pendula, P. 1266. Ubique ad rivulos.

\* Gr. Cyperoides spica pendula longiore & angu. stiore, C B. 87. 6. On the west side of Prim

rose hill.

"Gr. Cyperoides pulicare, hoc nomen impono, quoniam fingula famina pulicem quoad figuram magnitudinem & colorem belle expimunt, A mile East from Oxford, at Hockly of the Hole.

"Gr. Cyperoides spica echinata simplici, Tm miles southward from Oxford in the bogg.

Gr. Cyperoides Miliaceum, C B. 90. Cyperus gramineus miliaceus, G. 80 pseudocyperus gramineus sive miliaceus, P. 1171. In the Meadom betwixt Hampsted heath and Highgate.

\* Gr. Cyperoides tricubitale glumis ad singula

alas angustis longè distantibus.

\*Gr. Cyperoides spicis minoribus, minusque compactis, C B. 87.9.

\* Gr. Cyperoides cum glumis alternatim litis, Be twixt Kensington and Hammersmith.

Gramen Cyperinum nemerosum, G. 19. gr.nemerosum spica nigra recurva, C B. 98. in sylm humidis.

Gr. Dally oides radice repente. Cocks foot graft with creeping roots, G. 28. gr. Canarun Ischæmi panicula, P. 1179.

Gra

G

(

6

Gr. Fluviatile, Flote-gras, G. 14. P. 1275. gr. Aquat. Fluitans multiplici spica, C B. 41. In aquis stagnantibus & rivulis.

Gr. Fluviatile spicatum, G. 14. P. 1275. gr.
Aquat. geniculatum spicatum, C B. 42. Passim

in aquis & rivulis.

nik

87.

do.

que

gn.

in-

10,

go. pri

ly

WO

i,

ra

mi-

W

18

que

Be

ne.

vis

di

UÜ

Gli

\*Gr. Fluviatile simplici spica nigra. In Chelsey Meadows.

\*Gr. Flore Cæruleo minimum, On Bodle Downs Cheshire, it makes a pretty show being less then an inch long.

\*Gr. Fluviatile cause foliifque capillaribus ad articulos simul ortis, In the River near Snowdon.

\*Gr. Fluviatile facie Cornu Cervim, radice morfus diaboli cum priore.

Gr. Miliaceum fluitans suavis saporis, At Knights-bridge near London, and many stand-

ing waters.

Gr. Geniculatum, Kneed-grass, G. 15. gr. Geniculatum minus, P. 1177. gr. paniceum spica aspera, CB. 139.

Gr. Geniculatum aquat. G. 15. gr. Geniculatum majus, P. 1177. gr. Dactylum aquat. C B.

118. in lacunis.

Gr. Hirsutum exile, G.17. gr. Nemorum hirsutum minus angustisolium, P. 1185. gr. hirsutum capitulis psyllii, C B. 103. in arenosis & pascuis humidis, In Hide-park.

\*Gr. Hirfutum capite globoso simplici. On the

Moors in Lancashire near Hastenden.

Gr. Hirsutum nemerosum, G. 19. gr. Nemorum hirsutum majus, P. 1184. gr. Hirsutum latisosium majus, CB. 101. F 3 \* Gr.

\* Gr. Nemorum birfatum latifolium minus, P. 1185. gr. Hirfetum latifolium minus, CB, 102. below the Pine-walk at Hampsted.

Gr. Nemorum bir sutum majus angustifolium,P.

1185.

Gr. Junceum, Rush. grass, or Toad-grass, G.4. gr. Junceum parvum feu Holosteum, Math. P. 1190. ubique in stagnantibus.

Gr. Junceum Aquat. G. 1 2. gr. Aquat. junceum vulg. P. 1269.gr. Junceum polystachion, CB.

Gr. Junceum Sylvat. G. 22. gr. Junceum Aquat. Bauhini folio articulato, P. 1270. C B. 76. li the bogs on Putney beath.

Gr. Junceum articulatum tricubitale, solio pedali. In the Meadows by the Bridge near

Cafalton.

Gr. Junceum folio articulato aquat, cum panicula & utriculis, C B. 77. P. 1270. In the bog

on Putney-heath.

Gr. Junceum foliis oblongis radice alba, CB.73. gr.minus duriusculum, G.4.gr. Nemorumerik durum, P. 1183 super muris & in aridis.

\* Gr. Junceum foliis & spica Junci, C B. 78. gr. Junceum maritimum majus, P.1270 G.M.

Gr. Junceum spicatum seu Triglochin, C B. 81. gr. Triglochin, P. 1187.gr. marinum spicatum, G. 20.

Gr. Junceum marinum, G. 21.gr. Junceum m ritimum dense stipatum, P. 1270. gf. Junceum foliis & spica junci minus, C B. 79.

\*Gr. Junceum parvum five Holosteum Math.

spicis magis compactis, near Highgate.

Gr. Junceum maritimum exile, Plimostii, P.
1270. Plentisul where a small dril issueth out
betwixt the rocks, near the South-east point,
in the Isle of Wight.

Gr. Junceum montanum subcarulea spica, Mosscrops, In locis uliginosis prasentim montanis.

Gr. Junceum marinum spicatum, On this side Woolieh below Charlton in the marsh Meadows plentifully.

\* Gr. Junceum vix unciale flore Holostei.

Gr. Loliaceum, Darnel-grass, P. 1145. C B. ubique inter segetes.

\*Gr. Loliaceum seu Phœnix acerosa gluma, small

prickley Darnel grass, P. 1146.

Gr. Marinum spicatum, Sea Spike-grass, G.20' gr. Marinum spicatum alterum, P. 1279. est solummodo gr. Triglochin quando juvenescit.

Gr. Marinum spicatum, P. 1278. gr. Aquat.

spicatum, G. 13.

P.

P.

um

B.

ati

lo

olio

ear

nla

ogs

73.

rile

78.

21,

ım,

112-

gr.

791

ith.

Gr. Minimum rubrum seu xerampelinum, G. 3. gr. Arvense pannicula crispa, CB. 32. P.1157. nondum vidi.

\*Gr. Miliaceam, Millet grass, G.6.gr. Miliaceum vulg. P. 1153. gr. sylvat. pannicula miliacea sparsa, CB. 141.

\* Gr. Montanum pannicula spadicea delicatiore,

P. 1159.

Gr. Marorum spica longissima, Capon tail grass, G. 29. gr. spica nutante longissima, P. 1162. On the walls by Pickadylle, and beyond slough on the ground.

FA

Gro

Gr. Nemerosum majus spica rusescente, P.1183, gr. sylv, majus, G. 9. gr. nemerosum spica rusescente molli, C. B. 99.

Gr. Nemerosum minus, P. 1184. 3. gr. sylvat minus, G. 9. 4. gr. nemerosum spicis parvis

asperis, C B. 100.

Gr. Palustre Echinatum, Hedge hog-grass, G.17. gr. Aculeatum Germanicum, Prickly headel grass, P.1184. C B.109. In humidis paludoss.

\*Gr. Paniceum, Bearded panick grass, G. 16.gr. Panicum sylv.aristis longis, P. 1154. gr. panicum seum seum seum seum sylv.aristis armatum, C. B. 137. By a Rivolet side near Petersfield Hampshire, Mr. Goodyer.

\*Gr. Paniceum fylv. spica simplici, P. 1154.2. C B. 138. In hortorum areis, From Mr. Chau the Kings Apothecaries Garden in Kent.

\* Gr. Paniceum procumbens, seu chamæ paniceum palustre, In a Lane, and watery places, and Ditches near Petersfield.

\* Gr. Paniceum parvum, G. 16.

\* Gr. Panici effigie spica simplici, G. 17. In a field betwixt Tuddington, and Hampton-Court,

\* Gr. Paniceum sylv. spica divisa, CB. 136.gr. Panicum sylv. Herbariorum, P. 1154. Ina Lane by the Neat house Gardens.

Gr. Parnaffi, Grafs of Parnaffus, G. 840.P.419.

In pratis humidis.

\* Gr. Parnass. flore pleno, In Lancasbire.

Gr. Parvum Marinum panicula loliaca, an loliaca panicula ramosa maritimum, CB. 130? G. 30. 8. Gr. Phalaroides spica hirsutiore, P. 1164. gr. phalaroides, CB. 55.

Gr. Piperinum, Pepper-gras, Intermiscentur enim foliolis plurima grana magnitudine, & figura grani piperis, near Petersfield.

Gr. Plumosum elegans, Fethered grass, In Ellef-

more-Moor, Dr. Bowles.

gr. Pratense cristatum, Cox-comb grass, CB 43. gr. Cristatum Angl. P. 1159. 3. G. 29. 3.

Gr. Pratense, G. 2. gr. Pratense vulgatius majus, P. 1156. gr. Pratense paniculatum majus, C.B. 28.

Gr. Pratense minus, G. 2. P. 1156. gr. Pratense

paniculatum medium, C B. 30.

Gr. Pratense minimum album, P. 1156. gr. minimum album, G. 3. gr. Pratense panaculatum minus, C. B. 31.

Gr. Pratense paniculatum molle, P. 1155. CB.

27.

t

5

1

5,

r.

į.

14

2.

33

end

ld

11

29.

13-

69

170

on the farther fide of Micham Common by the foot way to the River leading to Calalton.

\*Gr. Pratenfe spica multiplici rubra.

\*Gr. Pratenfe panicula multiplici, P. 1157.

Gr. Secalinum maximum, The greatest Rye grafs, P. 1144. In the Woods a mile west from Petersfield.

Gr. Secalinum majus & minus. P. 1144. gr.

Spica Secalina, G. 29. 4.

Gr. Segetale, Corn grass, G. g. gr. Segetum panicula speciosa, P. 1158 gr. Segetum altissimum panicula sparsa, C. B. 34.

Gr.

Gr. Sorghinum, Darnel-grass, G. 9. P. 1154. Gr. Sparteum montanum spica soliacea graminea majus, The greater Grass-eard grass, amongst the Rocks at Carnedh Libewelyn.

\* Gr. Sparteum, &c. minus, On the Fels between.

Pereth and Kendal, and also on Snowdon.

\* Gr. Sparteum capite bifido vel gemino, At

Crixey Ferry in Effex.

\*Gr. Sparteum Batavicum & Anglicum, P. 1199
gr. Sparteum nostras parvum Lob. G. 1631.
gr. Hollandicum seu capillaceo solio minus,
CB. 70. On Wimbleton Common, and
Chiselburt Heath.

Gr. Sparteum Narbonense, French Matmeed,Pi

1199.

\*Gr. Sparteum nigro capitulo, A mile from Rochdale on the Turbery, the Country people call it Lank, they strew their houses with it in Lancashire.

\*Gr. Sparteum minimum Anglicum, P. 1199. On the next place betwixt Windfor Forrelt

and Redding, where they dig Furz.

\*Gr. Sparteum capillacio folio minimum, CB. 69. 9. G. cap. 22. 10. On Hampsted beath.

"Gr. Sparteum aquat. annuum, On Hunston heath in the new River running to Hampton Court.

\* Gr. Sparteum capitulis Equiseti, Beyond the Windmill at Adington in Surrey, where Peat is dig'd.

Gr. Spicatum alterum, G. 20. gr. Montanum

spicatum Clus. P. 1162.

Gr.

for. Spicatum aspero folio, CB. 45. Invenitur spica compactioni, & alterum spica maxime dispersa.

\*Gr. Spicatum tritici mutici, P. 1144. gr. tritici fpica muticæ fimili angustifolium, CB. 132.

In agris frumentaceis.

Gr. Tomentarium, Cotton-grafs, G. 29. gr. Juncoides lanatum five Juneus bombycinus vulg. P. 1272.

\*Gr. Tomentosum pratense pannicula sparsa, CB.
61. On the greatest bogg on Hampsted heath.

Gr. Tremulum majus, C B. 22. Phalaris pratenfis

Quaking grafs, G. 86. P. 1165.

Gr. Tremulum seu Phalaris pratensis media elatior, albis glumis, N.D. In a hollow Lane bewixt Peasly and Mansfield in Notting hamshire, Ph. Mr. Stonehouse.

\*Gr. Tremulum glumis albis. In a Meadow in Garforth in Yorkshire, Mr. Witham, and

betwixt Highgate and Hampsted.

Gr. Thyphinum minus, small Cats-tail grass, G. 1 12. gr. Typhoides asperum alterum, C B. 52. gr. Typhinum medium seu vulgatissimum, P. 1170.

Gr. Typhinum majus, G. 11. gr. Typhinum maximum, P. 1170 gr. Typhoides asperum

primum, C B. 51.
Gr. Typhoides maximum alterum, Betwixt Deal

and Sandwich.

.

C

ń

.

Gratiela angustisolia, Grass poley, G 581. Gratitiola angustisolia minor, P 220. Betwixt Aston and Oxbridge amongst the Corn, and in many places of Cambridge shire.

Gra-

Gratiola latifolia nostras, P.220. Gratiola latifo. lia, A broad leav'd hedge Hyssop, G. 581. est Lysimachiæ galericulatæ minoris species, Passim in uliginosis & ad rivos, variat fl. albo, near Purbright in Surrey, Mr. Brown.

### H.

Halimus vulg. five portulaci marina, common Sea-Purstan, G. 523. P. 724. Near Gillingham in Rumney-Marsh in Kent.

Hedera Corimbola, Climing or Berried Ivie, G. 858. Hedera Arborea noltras, P.678. pallim, Hedera helix, Barren or Creeping Ivie, G.858.

P. 679. scandit super arbores.

Hedera terrestris, Ground Ivie, Ale-hoof, G. 856. P. 677. ubique ad sepes & in soffis.

\*Hedera terrestris montana, P.677. Vide iconem in Cam. epitome pag. 401. At wigles-worth

near Setle in Yorksbire inter Segetes.

With Ivie-leaves, G. 830. Ranunculus hederaceus aquat.) P. 1216. Passim in sossis humidis, & aquis stagnantibus.

Helleborine, Wild white Hellebor, G. 442. Elleborine minor flore albo, P. 218. On Roe-bill

in Kent, not far from Dart ford.

Helleborine angustifolia sexta Clusii, G. 442. Elleborine angustifolia, P. 218. In montosis cum priore in variis locis in agro Surriensi.

Hele Helleborine flore atro-rubente, P. 318.7. At

Lyfnegorab, Mr. Heaton, Ph.

\*Helleborine multicaulis radice perplexa, By Cumner Wood in the way from Oxford to Eynsham Ferry, Mr. Pink.

\* Helleberine flore viridi, On the fouth fide of the great Beech-wood of Mr. Evelyn, two

miles from Darkin in Surrey.

\* Helleborine angustifol. fl. albo oblongo, In Helkwood in Yorksbire not far from Inglebo-

rough.

).

ft

ľ

.

.

\*Hellehorine nervo medio candido, In Stocken Church Meadows, Mr. Witham. Lusum esse naturæ non sine ratione suspicor.

Helleborastrum, wild black Hellebor, G. 976. Helleborastor minor flore viridante, P. 212.

In the Woods near Arundel in Surrey.

Herba Paris, One-berry, or Herb True Love, G. 405. P.390. variat foliorum numero à tribus ad septem, In a Wood near Hampsted-beath.

Herba Gerardi, Herb Gerard, or Ashmeed, G.
1001. Podagraria vulg P. 943. In viridariis
& juxta oppida, At Wallingford house, and in
Moorfields.

Hieracium asperum, sharp Hank-meed, G. 298. Cichoreum pratense luteum asperum, P. 777.

Adagrorum margines.

Hieracium longius radicatum, G.298.P.790. In

pratis.

Hieracism minus seu Leporinum, small Hares Hawk weed, or Yellow Divels-bit, G. 296.

Hierac.

Hierac.minus pramorfa radice, P.794.ubiq in pratis, Est hujus altera varietas paulo minoris foliifque magis dissectis.

Hieracium majus Diasc. G. 296. Hierac. majus

Sonchites, P. 788.

Hieracium Aphacoides, G. 297. Hierac. fol & facie Chondrillæ, P. 794.

\* Hieracium minus Cichorei vel potius stabes,

fol. hirfutum, CC.

\* Hieracium mont. Caule aphyllo non ramolo, flore pallidiore, an Hieracium mont. hir sutum

minus, CB? CCapp.

\*Hieracium angusto Tragopogi fol. On the back fide of Chelsey Colledge, Hide-park, and on Banstead Downs.

Hieracium asperum Hypochæris seu Porcellia dictum, Simnes Hawkw. P.791. Hypochæris Porcellia, G. 288.

Hieracium minimum Clus. P.791. Hyoseris mascula, G. 288. cum priore in montosis satis.

Hieracium luteum glabrum, sive minus hirsutum I B. pro loci ratione insigniter variat, viz in montosis vix triunciale, in pascuis Cubitum superat, C C. Cichoreum pratense luteum lavius, P. 778. ubique obvium.

Hieracium Intybaceum, G.298 Hierac.frutico.

fum angustifolium majus, P. 801.

Hieracium Dentis leonis folio hirsutum, Dandelyon Hawk-weed, G. 303. Hierac. Asperumsoliis & storibus dentis leonis bulbosi, P. 788. Hieracium caule aphyllo hirsutum, 1 B. Dentis Leonis solio acuto ix, sive asperum store magno Dentis Leonis, C B. In pratis & pascuis.

Hippolapathum rotundifolium, Bastard Rubarb. G. 389. In the Woods in Cornwall, Dr. Gunthorp.

,

1

Ŋ

(

m

in

m

m

0.

ie-

0.

li-

tis

g-

no

\* Hipposelinum, Alexanders, G. 1019 Hipposelinum five Smyrnium vulg. P. 930. At Hinton in a Hedge, CC.

Holosteum vernum, seu gramen Leucantheum,

Stichwort, G. 47. P. 1325. In Sepibus.

\* Holosteum vernum minus flore majore, In the Ditches betwixt Knights bridge and Kenfing-

\* Holosteum minimum, In Hide-park, and about

the Mill on Hampsted heath.

\* Holosteum hirsutum flore amplo, In the field betwixt Hampton-week and Tuddington.

Holosteum seu gr. Leucanthemum alterum, G.47 gr. Leucanthemum minus, P. 1325. On some

Banks and Pastures.

\* Holosteum repens juncifolium, At the bottom of the Moor on the East fide of Peters-field, and in standing waters in and about Stretham Ferry.

Hordeum distichon, Common Barley, G. 70. P.

1130. ubique feritur in agris,

Hordeum distichum minus, Sprit, or Battle-door Barley, so call'd in Kent, where 'tis fowed, and

in Lancashire, P. 1129.

Hordenm polyflichum five hibernum, P.1130. Winter Square Barley, In Glocefter-Shire, Bonder beer Barley.

Hordeum pracox, 'tis ripe three weeks fooner

then

then common Barley, at Evesham in Worcester Bire.

\* Hordenm spica longissima. In many places

of Wiltsbire on rich foil, Mr. Jenner.

Hordeum Spurium, Wall-barley, G.73. Hordeum spurium five spontaneum, P. 1147. passim in muris & viarum marginibus.

Horminum Sylv. wild Clary, or Oculus Christi, G. 771. Horminum fylv. vulg. P. 55. ad

agrorum margines & muros.

Hyacinthus Anglicus, English-Harebels, G. 111. ubique ad sepes & in sylvis, Hujus varietas flore albo, at the farther end of Black-heath, & quandoque flore carneo reppentur.

Hyacinthus, autumnalis minor, G. 110. Near Ditton, and on Kingstone-bridge, and going

to Hunstow heath.

Hyacinthus stellaris vernus pumilus Lob. In Barge Island in North-wales, and at the Kings-end near Dublin, Mr. Heaton, and in Anglesey, Mr. Willoughby.

Hyofeyamus niger, Black Henhane, G. 353.

P. 362. in fimetis.

\* Hyoscyamus luceus, Yellow Henbane, G. 356. Tabacc. Angl. P. 712. sponte nascitur in quamplurimis horcis, and betwixt St. Jamei's and Hide-park.

Hyofcyamus Paravianus, Tobacco, or Henbane of Peru, G.357. Colitur in quamplurimis locis, præsertim Winchcombiæ in agro Gloucestr.

Hypericum, St. Johns-wort, G 539. Hypericum vulg. P. 572. passim in sep.bus.

Hypi.

Hypericum minus erectum, P. 572.5. Hyper. pulchrum Tragi, G. 540. in dumetis & fylvis. Hypericum tomentolum, Lob. G. 540. Hypericum Tomentosum majus Hispan. P. 572. Hypericum fupinum glabrum, G. 541. Hypericum minus supinum, P. 572. 3. cum superioribus.

Y Acea five herba Trinitatis Elegantissima flore luteo amplissimo Pansie with a most ample yellow flour. Near Eldenhole in the Peak, and about Buxton in Darby Shire, and on the Black Monntain in Wales.

\* facea five herba Trinitatis fol. aureis striatis. This mightily beautifieth the border of an hedge in our own Garden at Methly in Yorkshire, where it groweth spontaneously, Mr.

Witham.

3

n

S 9

> Jaces nigra vulg. Black Matfellon, G. 727. P. 468. ubique in pratis, variat flore albicante.

> facea major, Great Matfellon, G. 727. 1. Jacza Nigra vulg. laciniata, P. 470. inter fegetes. facea flore albo, G. 728. Jacaa laciniata alba,

P. 471. at Martin Abbey in Surrey.

\* lacea nigra vulg. fol. diffectis flore ampliffimo, a mile below Bristol, on the South fide of the River, and in Viland Meadows and Pastures near the Bath.

G

\* Facaa

\* Ideaa binis perpetuo capitibus infignita majo.
re & minore, two miles on this fide Hereford, where all the Knapmeeds are fo.

Jacobaa Ragwort, G. 280. Jacobaa vulg. P. 668.

\* Jacobaa angustisolia, G. 280. Mountain Ragmort, On Gogmagog hills, and New-Market.
heath, and on the ditch sides at Willinborough.

\* Jacobaa latifolia, G. 280 Jacobaa latifolia Pannonica prima, P. 668. In humidis & aquosis,

\* Iberis Nacturtii fol. P.8 54. On the Clifts be-

yond Deal Castle in Kent.

\* Irio lævis Apula Col. 265. Erysemum latisolium Neapolitanum, Hedge Must ard of Naples, P. 834. Ubique sere in Suburbiis Lond, supra muros & juxta sossas.

Iris bulbosa Anglica, Onion Floure de luce, or water flagg, G. 99. Acorus palustris sive pseudo Iris lutea palustris, water Yellow flagg,

P. 1219.

Ischamon vulg Cocks-foot grass, G. 27. Ischam. fyl. latiore folio, P. 1178. in maritimis.

Juglans nux the Walnut-tree, G. 1440. P.

Juncus acutus, sharp or hard Rush, G.35. Junc. acutus vulg. P. 1193. passim in paludoss.

Juneus minor capitulis equifeti, Club. Rush, G. 1631. Juneus Aquaticus capitulis equifeti, R. 1596. datur varietas quoad magnitudinem, P. ibid.

Juneus aquaticus maximus, Bull-Rush, G. 35. June. lavis maximus, P. 1191.

funcia lavis vulgatior pannicula sparsa, P. 1191. in & juxta aquas.

funcus acutus Cambrobritannicus, P. 1193.

\* funcus caule Triangulati, At the Horse-ferry
at Westminster, Hunc mihi primus ostendic
Dr. Dale insignis Britannicus.

\*funcus palustris pannicula glomerata ex rubro nigricante, Teversham Moor in Cambridge-

Bire.

.

t.

6.

n-

e.

į.

4-

d.

10

ve

g,

M.

P.

nc.

G.

m,

35.

CW

Juncus maritimus capitulis forbi, P. 1 193.

funcion ad instar spinæ acutus rigidusq; In the mid way twixt Sandwich and Dale in a sandy

Cony Warren.

\* Juneus capite globoso amplo. Half a mile beyond Atton in Middlesex, and on Shooters hill. Juneus lævis medulå solidiore, Candle-Rush, an juneus lævis alter pin 12. (a 7?)

funces five scirpus medius sylvat. Tab. In the Mill at Cobgrave not far from Rippon, and at

Nostal Abbey in York Shire.

Juniperus The Juniper Tree, G. 1732. Juniperus vulg. seu minor, P. 1029. In many hills in Surrey and Oxford-sbire.

Juniperus sterilis, frequent on the mountainous

parts of Surrey, Mr. Goodyer.



## K.

Ali geniculatum, five Salicornia, Sea-grape, Glafs-wort, or Salt-wort, G. 535. Glafs. wort joynted, P. 280. in falsis & maritimis. Kali minus, G. 535. Kali minus album, P.279. cum priore.

Kali spinosum, Prickly Sea-grape below Whit-

Staple in Kent and in Cornwal.

## L.

Actuca Agnina , Lambs-Lettuce, G. 310. Corn-Sallet, P. 812.

Lattuca fylv, foliis diffectis, G. 309. Lattuca fylv. laciniata, P.814. ad arvorum margins, Lactuca sylv. major odore Opii, G. 309. Lactu. ca fylv. Endiviæ folio odore viroso, P.813.10 sepibus & juxta vias circumcirca Londinum.

\* Lattuca fylv. laciniata minima, ND. A quarter of a mile beyond the Spittle, in a lane lead ing from London Road to the River in a Ditch, C C. and near Church lench in Wor cester-fbire in great plenty.

\* Lactuca fylv. altera folio non laciniato, odor

minus vehemente.

Ladaum segetum Plin. G. 699.7. Sideritis arvensis flore rubro Cam. P.58. In the Cornfields of Kent, and at Purfleet in Esfex.

Lagopus vulg. G. 1193. Lagopodium seu pes leporis, Little Hare-foot Tresoil, P. 1107. In-

ter fegetes.

J.

19:

oit-

10,

tuca

nes,

Au.

3.In

₩.

var-

ead.

in a

py or

dore

ada

Lagopus altera angustifolia Advers. 384. G. 1193. 3. at Stretham in Surrey.

Lamium album, White Arch angel, G. 702.

Lamium flore albo, P. 605. ubiq;

Lamium luteum, G. 702, P. 606. In sepibus, fed rarius in Hamsted Church-yard, and in the west side of Primrose-hill.

Lamium rubrum, G. 703. Lamium vulg. flore

rubro, P. 605. variat flore albo.

\* Lamium rubrum foliis profunde incisis, In the Kings new Garden near Goring-honse.

Lampfana, Dock-cresses, G.255.P.810.ordinary Niple-wort, Passim ad agrorum margines.

Lapathum acutum, Sharp pointed Dock, G. 338. Lapath, acutum majus Red Dock, P. 1224.

Lapathum acutum minimum, G. 388. P. 1224. Lapathum sylv. folio minus acuto, G. 388. La-

path. fylv. vulgatius, P. 1225.

Lapathum pulchrum Bononiense, I B. quod non ineptè vocari potest, The Fidle-Dock propter figuram soliorum inseriorum, Common in St. George's Fields, and other places about London

\*Lapathum angustifolium polyspernum, Betwixt Wishich and Giborn, in the way to Peter-

burgh.

G 3 \* Lapa

Lapathum foliis crifpis, In St. George's Fields.

\* Lapathum fanguineum fativum, Blood-wort, G. 340. Lapath. sanguineum, P. 1226. In the Woods about Hampfted.

Lapathum aquaticum seu Hydrolapathum magnum, G. 389. P. 1225. In fossis & stagnis.

Lapathum seu Hydrolapathum minus, G. 389. P. 1225. cum priore.

Lathyris feu cataputia minor, Garden. Spurge, G. 503. P. 191. qui dividit in majorem minorem & minimam, nescio quo jure, sponte sua micitur in hortis.

Lathyrus major latifolius, Peas- Everlasting, G. 1229. Lathyrus major perennis, P. 1061. In fylvis & dumetis, fed minus frequens.

Lathyrus fylv, flore luteo, Tare- Everlasting, G. 1231. P. 1062.7. ubique in pratis & fylvis,

\* Latbyrus binis ad fingulas alas filiquis, On the banks of the Fields near Pickadilley.

\* Lathyrus perennis filiqua hirfuta. At Hadley Castle two miles from Lee in Effex.

\* Lathyrus major angustifol. flore pallide rubro Hampsbire, Mr. Goodyer.

\* Lathyrus major perennis, near Tewis in Suffex, idem, and at the Devices in Wilishire, in the way to Lavendon in a shady Lane.

\* Lathyrus flore ex caruleo & rubro mixto. In a wet marsh ground on the left hand of Puk-

bam Field from London.

\* Lathyrus perennis siliqua lata glabra. Neara Stone-bridge in the way from London to Harrom of the bill.

Laureola, Spurge Lawrel, G. 1104. P. 205. In sylvis & sepibus.

de

Lens major, & minor, Greater and leffer Lentils. G. 1224. P. 1068. Læte proveniunt ex femine in agro Oxoniensi.

Lens palustris, Ducks-meat, G. 829. Lens paluftris five aquat. vulg. P. 1262. In omnibus

aquis stagnantibus.

Lepidium Ægin. Lob. & raphanus sylv.officinarum, Dittander and Pepper. wort, G.241. Piperitis five lepidium vulg. P. 855. On most of the Ditches in the Ifle of Thanet.

Leucoinum luteum vulg. feu Keyri, Wild Wallflowr, P. 625. Viola lutea, G. 456. In muris

paffim.

9.

m

1

; 5,

ne

17

0

In

k-

2

7.

¥.

Leucoinum marinum latifolium, G. 460. P. 623.

Leucoinum marinum purpureum, Lob. G. 460. P. 662. On the Rocks at Aberdoye in Merioneth-Shire.

Lichen arborum, Tree Lung-Wort, G. 1566 mufcus pulmonarius seu vulg. P. 1313. Antiquis adnascitur quercubus & fraxinis.

\* Lichen seu Hepatica minor umbellara, G. 1565.

P. 1314. in areis hortorum.

\*Lichen humiditate intumescens, siccitate evanefcens. On the Downs west from Andover.

Lichen five Hepatica terrestris, Ground Liverwort, G. 1565. Lichen five Hepatica vulg. P. 1315. On Ditch fides, and in Woods.

Lichen marinus rotundifolius, Round-leav'd Oy-Myr. weed, G. 1567. Opuntia marina, The Sea-

Carland,

Garland, P. 1294. offrearum testis adnascitur.

Lithen sive Hepatica minor stellaris, Liver-wort,
with round heads, P. 1314. G. 1565. In a
pond near Twitnam by the Bridge.

Lichen petræus purpureus Darbiensis, Cork, or Arcel, In Laucashire they use it for Dying.

\* Lichen Capillaceus ex plurimis capillamentis nigricantibus constans, At Middleton-tire, and in a Lane a mile from Heptamstall at Greenwood-Lee in York shire,

\* Lichen tubulatus, At Avery-Farm in the way

from Westminster to Chelfey.

Lichen marinus, Sea Lung-wort, or Oyster-green, G. 1566. sucus marinus, lactuca marina dictus, P. 1293.

Arum vulg. P. 1446. In sepibus & sylvis.

Lignum lithoxylon, Petrefied wood, P. 1276. G. 1587. In variis Britanniæ locis, de quo postea.

Lilium convallium, Conval, or May Lillies, G. 410. On Hampsted-heath plentifully.

Limonium, Sea-Lavander, G. 411. Limon. vulg. majus, P. 1234. ubique in maritimis.

Limonium parvum, Rock Lavender, G.411. On the Cliffs at Margate.

Linaria vulg. lucea, Great Toad-flax, G. 550. Linar vulg. nostras, P. 458. In pratis.

Linaria adulterina, G. 555. Pseudolinaria montana alba, P. 459.

Linum sativum, Garden flax, G. 556. P. Linum sativum sive vulg. P. 1335.

Linkm

Jinum sylv. Catharticum, Mill-mountain, G. 560. Chamælinum. Clus. flore albo, sive linum sylv. P. 1336. In montosis & quandoque in pratis nonnullis.

Linum sylv. tenuisolium, G. 558. Linum sylv. latisol. P.1335. Near Compton in Surrey, and

in Devenshire.

ir.

rt,

a

or

tis.

e,

at

iy

15,

1.

5.

0

n

•

n

Linum latifolium sylv. G. 559.Lin. Caruleum, P. 1335. on New-Market-heath, and near Stanford in Linconshire.

Lithospermum majus, Great Gromell, G. 609. Lithosp. vulg. majus, P. 431. In collibus.

Lithospermum minus, G. 609. Lithosp. minus vulg. P. 432. In variis Devoniæ locis.

Lithospermum Anchusæ facie, Purple-flour'd Gromel, G. 610. fmall Gromel with tusted tops, P. 432.

Lolium album, & rubrum, white and Red Darnel

G.78.P.1144.Inter fegetes passim.

\*Lolium foliis angustioribus brevioribusque, In Oxford shire and in Kent: Icon ludg. 417. convenit.

Lonchitis aspera, Rough Spleen wert, G. 1140. Lonchitis aspera minor, P. 1042. In asperis & umbrosis.

Lorns quadrifolia, Four leav'd Grass, G. 1198.

Ad pratorum margines.

Lunaria minor, Small Moon wort, G. 405. P. 507. In montosis & ericetis. In several places near Oxford.

Lunaria minor ramosa, Three nales from Ox-

ford near the Blind-Pinnosks.

\* Luna-

\* Lunaria minor fol. dissectis, Plentisul in John Nuns Cow-pasture adjoyning to his house at Methley in Yorksbire. An lunaria minor sol, rutaceo. pin. 355. (b. 4?) Mr. Witham.

Lupulus Salictarius, Hops, G. 885. P. 177. In

Effex, Kent, Surrey and Suffex.

\* Lupulus sylv. & floribus sparsis & compactis, In sepibus.

Luteola, Dyers-meed, or Yellow-meed, G. 494. Lutea vulg. P. 603. In muris, seritur in quibusdam locis pro tinctorum usu.

Lychnis fylv. flore rubello, Red wild Campions,

G. 469. P. 631. Passim ad sepes.

Lychnis marina Anglica, G. 469.P. 639. Plentifully in the low Marish-grounds in Thanet.

Lychnis fegetum parva, viscosa, flore albo, Ph. \* Lychnis marina floribus albis pendulis, foliisq, inferioribus hirsutis. Betwixt Margate and Sandwich.

\* Lychnis sylv. flore carneo odorato, near Peters Fiela in Hampsbire. Ex Misto Gooderiano.

Lysimachia galericulata, Hooded Loos Strife, or Willow herb, G. 477. Gratiola Cærulea sive latifolia major, P. 221. Passim ad rivulos.

\* Lysimachia spicato flore amplo lacteo, Aoundantly near Castle-Eaton in Wiltshire, Mr. Fenner.

Lysimachia Cærulea, G. 477.

Lysimachia campestris, G. 478. Lysimach. Caru-

lea spicata siliquosa major, P. 548.

Lysimachia omnium minima annua, In the Combetwixt Donington and Marish in the Isle of Ely.

Lismachia lutea, G. 474. Lysimach.lutea major

vulg. P. 544.

×

It

1.

n

,

j-

rs

0. 1

C

۲.

1.

of i-

Lismachia lutea minor, G. 474. P. 543. 2. Listimachia luteo flore globoso, G. 475. P.544. Lifimachia purpurea trifolia cauleHexagono, Spigel. Ifag. I. c.10. CC.

Lifimachia filiquofa glabra media feu minor an-

gustifolia, G. 478. 12. P. 548. 4.

Lismachia purpurea spicata, G. 476. P. 546. \* Lysimachia siliquosa hirsuto magno flore, CB. Lysimac siliquosa hirsuta majore flore purpureo, IB. CC.

# M.

A Alva arborea marina nostras, English Sea-tree mallow, Mr. Morgan received it from the Ifte of Wight.

Malva fylv. G 930. Malva vulg. flore purpureo, P. 299, variat flore albo passim in hortis

& sepibus:

Malva fylv. pumila, G. 930. Malva fylv. minor, P. 299.

Malus Sylv. The Wilding or Crab tree, G. 1461. P. 1502. funt infinitorum pene generum.

Marrubium album, White Hore-hound, G. 693 . Marrubium album vulg P. 44. In locis faxofis & aridioribus.

Marrabium nigrum, flinking Hore bound, G. 701. MarruMarrubium nigrum fætidum Ballote dicum. P. 1230. Passim juxta vias & sepes.

P. 1230. Pallim juxta vias & lepes.

Marrubium aquaticum, G.700. Marrub.aquat.

vulg. P. 1230. In fossis & aquis stagnantibus, Matricaria, Fevefem, G.652. Matricaria.vulg.

P. 83. In muris & ad fepes.

Medica marina Trifolium cochleatum marinum, Sea-medick, G. 1200. P. 1114. At Rumney betwixt the Town and Cony-Warren.

\* Medica cochlea simplici, Below Branford where

they burn Chalk to Lime.

\* Medica spinosa globosa, By the Wind-mills at New market, where the Sefamoides grows, CC.

Medica, vide Trifolia.

\*Melampyrum cristatum flore purpureo, Purplecrested Cow Wheat, N.D. In Madingley and Kingstone Woods, and almost in all other Woods in Cambridge shire plentifully, CC.

Melilorus Germanica, German Claver, or wild Melilot, G. 1205. Melilotus vulg. P. 719.

in agris.

\* Melissa Fuchsii flore albo & purpureo, Bastard Balm with white and purple flours, G. 690. Melissophillum Fuchs. P. 41. In the Woods and Hedges a mile from Exeter, near the great stone-bridge going over the River Ex, and divers other places of Devonshire.

Mentha aquatica, sive Sisymbrium, Water-mint, G. 684. Mentha aquatica rubra, P. 1243. Ad

rivulos & aquas stagnantes.

\* Mentha Aquat. tota nigra, In many places of Lancashire, on Ingleborough-hill side towards S. tle. \* Mentha

Mentha odorata flore cinertio. Five miles from

Glocefter, in the way to Hereford.

\* Mentha Balsamita vel latisolia odorata. In the way betwixt Pemsey and Lewis in Sussex. Diutissime durat odor ejus suavissimus.

' Mentha incana fœtida.

Memba felina five Cattaria, Neap, or Catmint, G. 682. Calamintha vulg, P. 36. Juxta fepes in locis aridis.

Menthastrum, Horse-mint, G.684. menthastrum hortense sive mentha sylv. P. 33. In pratis humidis.

Menthastrum valde ramosum flore violaceo rubro, At Dartford in Kent.

Mercurialis mas, male Mercury, G. 332. P.

295.

e

r

1

Mercurialis fæmina, G. 332. P. 295. both by the Sea fide, at Ride in the Isle of Wight, and all about Dover.

Mespilus sylv. spinosa, wild Medler, In the Hedges betwixt Hams sted heath and High-gate, and in a Holt of Trees three miles west-

ward from Crediton, Devonshire.

Meum, Mem or Spignel, G. 1052. Meum vulgatius, P. 888. In Westmoreland, and in most parts of Scotland, and Sedbury, York-sbire.

Militaris aizoides, Fresh-water Souldier, G.825.
Stratiotes sive Militaris aizoides, P. 1249. In Lincoln-shire Fens, and those of Ely, and at Awdery Causey in the way from London to Cambridge.

Mille\_

Millefolium aquat, water Tarrow, G. 827 Millefolium aquat, vulgatius, P. 1256. paffin in locis udis.

Millefolium, five maratriphyllum flore & semine Ranunculi aquat. Hepaticæ sacie, Crow-soot, or water Milsoil, G. 827. Millesolium aquat, Ranunculi flore & capitulo, P. 1256.

Millefolium tenuifolium, Fennel leav'd mater

Milfoil, G. 828. ubivis in stagnis.

Millefolium aquat.floridum, water Gilly flour, P. 1256. viola palustris, G.826. In all the Ditches about Oxford.

Millefolium aquat pennatum spicatum, P.1257.
Millefolium palustre galericulatum, G.828. Millefolium aquat. flore luteo galericulato, P.

1258.

Millef-lium aquat. cornutum, P. 1257. In the Ditches about Oxford.

Millefolium seu myriophyllum aquat. minus Clus. G. 827. Millesol. aquat. minus, P. 1256.

- \* Millefolium aquat densioribus versus summitatem pennis colore rubro, In Putney Common near Wandfor, in standing pits near the Road side.
- Millefolium terrestre vulgare, G. 1072. Millefol. vulg album, P. 694. variat flore rubro fed hoc infrequens.

Millegrana minima, Dwarf-Allseed, G. 569.
Millegrana sive herniaria minor, P. 447. On
Putney heath or Common plentifully.

Monophyllon seu unifolium, One-blade, G. 409. Monophyllum, P.505. In Dingly wood Lancafaire, and at Homgill Yorkshire, in a field next Todberry in a moist ground belonging to Thomus Oddy.

Morsus Diaboli, Devils-bit, G.726. morsus Diaboli vulg. P. 492. In pratis, satis, & pascuis,

variat flore albo.

ne

t.

er

1-

P.

e

15

9-

n

d

j.

n

g.

0;

Morsus Rana, Frogg-bit, G. 818. Nymphæa alba minima sive morsus Rana, P. 1252. In

aquis stagnantibus.

Morsus Rana flore pleno odoratissimo, N.D.!n a Ditch by the side of Andery Causey, close by the great wooden Bridge plentifully, C.C.

Morus, The Mulberry-tree, G. 1057. Morus

nigra vulg. P. 1491.

Muscus quernus, Moss of the Oak, and other trees, G. 1558. Muscus arboreus vulg. & quercinus, P. 1312.

\* Muscus arboreus umbilicatus, Common in the

Orchards in Suffex.

\* Museus caule rigido ad instar fili Chalybei, Six miles from Bristol, in a shady lane near Winford where they dig for Oshre.

Museus alius terrestris, lycopodium Hist. lug. pag. 1325. On Putney heath on a bog near

the Wind mill.

Muscus aridus crustatus, P. 1313.7. On old

trees and pales.

\* Muscus aureus tenuissimus, Four miles from Basing-stoke, in the way through the Wood leading to Salisbury.

Museus polyspermus, On the North of Ingle-

borough near the under ground River.

\* Mascus maritimus ruber tenuissime dissetts, pin. 363. (a. 4.) Betwixt Deal and Dover.

\* Muscus terrestris denticulatus alter, & terrestris repens Clus. Vid. Hist. Ludg. 1326. On the East side of Rumble-moor, above Helmid. Lancashire, in the deep Heath.

\* Museus Triangularis aquat. CCapp.

\* Muscus repens virore splendens. On the old

walls near Banbury, Oxford Shire.

\* Muscus ex maxilla interiore ovilla quem'a musco Cranii humani distinguere nequeo on a stone-wall at Arlington, Glocoster-Shire.

Muscus corniculatus Knagged-moss, G. 1561, P.
1308. In herbosis montium jugis, & in Tud-

dington field near London.

Missous clavatus foliis Cypressi, heath Cypress, G.
1562. Muscus clavatus Cypressi formis, P.
1310. On Cheviot hills in Northumberland, and near the Mile-cross west from Halifax.

Muscus clavatus, five Lycopodium, Club moss of Wolves claw, G. 1562. P. 1307. On Hampt-

fed beath

In some Charnel-honses, chiefly in Ireland.

Muscus Corallinus seu Corallina montana, G. 1572. Muscus Corallinus montanus, P. 1316. On Hampsted-heath, and in dry Mountains of Banks. Variat color à cinerito ad candorem.

Muscus denticulatus major & minor, Toothid, or Dented-moss, G. 1561. P. 1307. in humidis. The greater at the Neat houses and Kingstontbridge by the Thames side.

Muscan

Muscus filicinus, G. 1561. P. 1310. On the East fide of Hamsted heath in the ditch neer the wood.

\* Muscus Hirlutus capillaceus, On Boulfover-

hill Yorkshire.

e-,

)n

ck

old

1

Ón

P.

id-

G.

P.

01

pt-

13.

G.

0.

or

or dis.

ne-

CHI

\* Muscus Lutescens fol. convolutis, On the houses and walls in Chismick.

Museus arboreus nodosus sive geniculatus, P.

1311.5.

Muscus marinus spiralis pennatus, N D. On the

coast of Anglesey.

Museus palmaris. 4. gestans in summitate solia ad instar Tormentillæ, Betwixt Plumfreet and Crayford in a dark lane leading to the Moor in Kent.

\* Museus pennatus longo latoque folio, By

Richmond Castle in Yorkshire.

Muscus pyxidatus, Cup or Chalice-moss, G. 1560. P. 1308. In muris & locis ficcioribus.

Muscus ramosus floridus, G. 1560. Muscus quercinus fæniculaceus, Fennel like Oak-moss, P. 1311. 2. at Midston tire Torkshire.

Muscus parvus stellaris, Small Heath moss, G.

1563. P. 1308.

\* Muscus stell. roleus. Half a mile from Crediton in Devon G. at a bridge in the way to Exen.

Muscus terrestris vulg. G. 1559. Muscus ter-

restris vulgatifimus, P. 1306.

\* Muscus terrestris scoparius, G.1559. P.1307.

\* Muscus terrestris minimus, P. 1309. 13. In areis humidis pedibus non multum tritis.

\* Muscus velutinus, Velvet-moss, sic dictus à
H pisca-

piscatoribus Lancastriensibus, quo purgant vermes Adnascitur plerumque veterum spinarum radicibus.

Myagrum, Gold of Pleasure, G. 273. Myagrum fylv. sive pseudomyagrum, P. 868. In sais

quandoque inter linum reperitur.

Myosieis scorpoides palustris, water Scorpiongrass, G. 338. Myositis Scorpoides repens, small creeping blew Monse-ear, P. 691.

Myofitis scorpoides arvensis hirsuta, G. 338. P.

691. utræq; in locis humidis.

\* Myofitis scorpoides hirta minor. In muris & locis siccioribus.

\* Myositis scorpoides latisolia hirsuta, In Charlton, and many other woods in Kent.

Myrrhis nova Æquicolorum, small Hemlod Chervil, G. 1038. 5. Myrrhis sylv. Neapolitana, atque etiam Anglicana, P. 935. Passim in sepibus. Vide Lyte, 616. & Col. 112.

Myrtus Brabantica, seu Eleagnus Cordi, Gal, Sweet willow, or Dutch-myrtle-tree, G. 1414. Rhus syl, sive Myrtus Anglica, P. 1451.00 Hanstow-beath, and Lincolnshire Fens.

# N.

N Arcissus, seu pseudo-Narcissus Anglicus, common yellow Daffodil, G. 133. In locis humidis.

15.

P.

&

rl.

lock

00-

af.

12.

all,

14.

On

Tus,

Nasturtium aquaticum, sive Cratevz Sium, Common Water Cresses, G. 257. Nasturtium aquat. vulg. P. 1239. In plerisque udis sossis & rivulis.

Nafturtium petræum, Stone-Cresses, G. 251. P.828. At Ham a mile on this fide Kingstone, on the dry Sandy way.

Nidus aris, five Satyrium abortivum, Birds-neft, G. 228. Orchis abortiva, P. 1362. On Roe-hill in Kent plentifully, and in some woods.

Nidus avis flore & caule violaceo purpureo colore, an pseudolimodoron, Clus. Hist. rarior plant. pag. 270. A mile from Alton in Hamp-fbire, Mr. Goodyer, Ph.

Nummularia, Herb Two pence, G. 630. Nummularia vulg. Money-wort. P. 555 In fossis

humidis & locis uliginofis.

Nummularia flore purpurascente, G. 630. P. 555. Cum priore sed rarius, in the meadows betwixt Deptford and Lemsham, neer the mill.

Nux vesticaria, Bladder nut, G. 1437. Nux vestic, five staphylodendron, P. 1418. In some woods in the farthest parts of York soire.

H 2

Nym-

Nymphea alba, G. 819. Nymphæa alba major vulg. P. 1252. The greater white-water Lilly, In fluviis & rivulis.

Nymphea alba minor, cum priore sed rarius,

G. 820. P. 1252.

Nymphea lutea, G. 819. Nymphæa major lutes, P. 1252.

Nymphaa lutea minor, G. 819. P. 1252.

\* Nymphea lutea minor flore fimbriato, JB. In the Rivers about Stretham-ferry, and in many Rivers about the Fens, CC.

### O.

O Enanthe aquat. Selini folio, ND. Water-

\* Oenanthe angustisolia Lob. P. 894. Filipendila angustisol. G. 1059. At East How in the Parish of Subborton Seven miles from Peters, field, Hamshire. Mr. Goodyer.

\* Oenanthe Aq. sive palustris, P. 895. In Mag-

dalen Colledge Meadows Oxford.

Oenanthe cicutæ facie Lob. Lobels Hemlock like Dropwort, P. 894. filipend. Cicutæ facie, G. 1059. In many places by the Thames fide. Vid. filipendula.

Onobrychis, five caput gallinaceum, Ordinary Cocks-head, or Medick Fetchling, G. 1241

P. 1082. In montofis.

6

Ombrychis flore purpureo, G. 1243. P. 1082. In some mountanous parts in Surrey plentifully

Obiogloffum, Adders tongue, G. 404. P. 506 Ludit quandoque, bilinguisque ex accident;

invenitur. In pratis humidis.

Orchis Andrachnites, Spiders Satyrion; G. 216. Orchis Arachnites, P. 1352. In an old Stonepit ground hard by Walcot a mile from Barneck, it flowrs betimes Dr. Bowles,

\*Orchis Antropophora autumnalis Col. mas

C B. & P. 1347. The man Orchis.

\* Orchis Antropophora Oreades altera Col.

pag. 318.

a,

in

Â

he

G. R.

ry F

MO-

\*Orchis Oreades trunco pallido, brachiis & cruribus faturate rubescentibus. These three Saz tyrions were found on several Chalkey hills, neer the high way from Wallingford to Rede ding on Bark fire fide the River by Mr. Brown.

Orchis Batrachites, Frog Satyrion, G. 224. P. 1352. by Barkway, and in many places about

Oxford.

Orchis Batrochoides autumnalis A. luteo parvo, three miles on the East fide of Lewis in Suffex. Orthis Hermaphroditica, Butterflie Satyrion G. 211. Serapias bifolia, five trifolia minor, P.

DI350.

Orchis Hermaphroditica supremo petiolo pupura maculato, By Vden Lodge in Pennystone Moor York Shire, Mr. Stonebouse, Ph.

Orchis Melittias, Bee-Orchis, G. 213. P. 1352. In Chalkey Holes on Bansted Downs in

H 3

Surrey,

Surrey, and many places about Oxford and in

York-Shire

Orchis Myodes Fly Satyrion, G. 213. Orchis Myodes minor, P. 1352. on a Hill Southwest of Bath.

Orchis Myodes lutea, G.214. In montofis pratis, Orchis Myodes purpurafcens, In Troadworth wood York-Bire, Mr. Stonehonfe. Ph.

Orchis Myodes floresoccineo, In Swanfcomb

wood, idem.

Orchis Orthinophora, Birds Satyrion, G. 215.

Orchis folio maculofo, P. 1353.

Orchis feu testiculus vulpinus secundus Sphegodes. Humble Bee Orchis, G. 21 3. Orchis fohegodes five fucum referens, P. 1350.

Orchis five Cynosoychis morio mas, Male Fulfromes, or Crow-toes, G. 208. Orchis morio mas foliis maculatis, P. 1346. In montolis pratis,

Orchis seu Cynosorchis, morio fæmina, serapia batrachites, G. 224. P. 1347. In pratis.

\* Orchis five Cynosorchis militaris rubra, P. 1345. 7. In a Chalkey close neer Cherrybin. ton, where they burn Lime, and in many other places of Cambridge-Shire, CC.

Orchis seu Cynosorchis major altera, white Dog-ftones, G. 205. Cynosorchis latifolia

minor, P. 1344.

Orchis five Chamzorchis latifolia Zelandiz, 1. 1354: In the watery places of Hinton and Feversham-moors, but rarely.

Orchis Zelandia minima. On the bogs betwist

Southampton and Rownams.

\*Orchis

\*Orchis Caryophillata spica longissima rubra, In many Chalkey grounds, bogs and old pastures betwixt North-fleet and Thames in Kent.

\* Orchis flore albo minimo, betwixt Weymouth

and Margate.

in

his

h-

is.

th

иb

5.

ŝ

ß

\*Orchis tota lutea, Betwixt Micham and Cafalton, in the first Close over the River, at

the Bridge-end.

Orchis Frisia lutea, Frisland Lady-traces, G. 219. Orchis Frisia littoralis, P. 1354. on Gads bill in Kent, and neer Greenbith.

\* Orchis militaris polyanthos, on Gads hill in

Kent.

Orchis seu Palma Christi mas, Male Satyrion

Royal, G. 220. P. 1356.

Orchis seu Palma Christi famina, G. 220. Orchis Palmata famina, seu Palma Christi famina maculato folio, P. 1357.

Orchis, five Palma Christi palustris, G. 227.

Orchis palustris maculata, P. 1360.

\* Orchis palmata flore viridi, P. 1359. 17. In fome Pastures and closes on the north-side of Chesterton plentifully, CC.

\* Orchis pufilla odorata, P. 1354. In a boggy place two miles from Stanford in the way to

Stilton.

\*Orchis palma quadruplici flore cineritio, At the East end of Wickham in a narrow lane.

Orchis palustris lophodes, N D. Tufted Marish Orchis, Ph.

Orchis radice repente, G. 227. P 1355. In fylvis & pratis humidis.

H 4 \* 0r-

\* Orebis grumofa radice, A mile South from Crediton in Devonsbire.

Orchis seu Serapias minor nitente flore, Red. handed Orchis, G.222. Orchis palmata minor flore rubro, P. 1358. In montofis.

Orebis, five Serapias candido flore, G. 222 two miles of Oxford from Frier Bacons, Study,

Orchis, seu Serapias palustris latifolia, G. 222. P. 1358. In humidis pratis.

\* Orchis spiralis flore luteo simbriato, On the South fide of Cherry-hinton, in the last of all the old Chalk-pits you pass through, C C.

Orchis seu testiculus odoratus, Lady traces, G. 219. Triorchis alba odorata major & minor. P. 1354. In montofis & pratis ficcioribus,

Orebis, five Tragorchis maxima, the greatest Gaot Stones, G. 210. P. 1348. Nigh the highway between Crayford and Dartford in Kent.

Orchis, feu Cynoforchis minor Pannonica & 207. Cynosorchis militaris Pannonica, P.

1345. On Gog magog bills CC.

Orebis abortiva ruffa. In the woods and grounds by Stoken Church in the way from London to Oxford, Mr. Witham.

"Orchis floribus purpureis. Near Hampalby

Robin- Hoods Well York Shire.

\* Orchis five testiculus sphegodes hirsutis floribus C Capp.

\* Orchis spicata fl. luteo. In cow pastures in mile from Long Prefton Lancasbire.

Orthis sphegodes major & minor. Near Stocken Church, and in many places about Oxford

Orchis fl. nigro sptendente. In the pastures where Setle Tork-shire.

Morchis strateumatica major & minor. In Stok-

en Church Woods, Mr. Witham.

In some grounds of Chiswell Berk Shire.

G. 666. Majoran sylv. P. 12. ad agrorum

and margines.

om

ed.

Ör

WO

2.

he

all

G.

ì,

eft

h-

nt.

d

m

y

i-

1

n

0

Ornithogalum, Star of Bethlehem, G. 165 On the top of a Hill three miles on this fide Briftol.

Ornithogalum luteum, sive Cepa agratia, G. 165. In a Corn field in Winetaunton Somer-

fet Shire.

Ornithopodium majus & minus, Birds-foot, or . Clam, G. 1241. P. 1092. In ericetis & locis aridis, On Black heath, and Hamfted-heath. Orbanche, five rapum Genistæ, Broom-Rape, or Orobanche, G. 1311. P. 1362. In ericetis & locis aridis, quandoque ex accidenti ramissicatur, uti observavit Dmus Loggins aliique.

\*Orobanche vasculo seminali pentagono, vel Hexagono, In Surrey near White stable.

Oralis Franca five Romana, Round-leav'd, or French Sorrel, G. 397. Oxalis fativa Franca five Romana rotundifolia, P. 743. On the Rocks of Snowdon.

Oxalis major forțe major Brockenbergensis, Montana maxima, CB.

Oxyacantha vid. Berberes.

Oxyacanthus, The Hawthorn tree, G. 1327
Spina

Spina appendix vulg. P. 1025. Passim in sepi-

Oxyacanthus folio & fructu majore. In Rich Park and elsewhere in Oxford Shire.

Oxys alba, White Wood Sorrel, G. 1201. Trifolium acetosum vulg, P. 746. variat flore rubro, at Chiselbart in Kent.

Oxys lutea, G. 1202. Oxys flore luteo, P. 746.

In fylvis.

#### P.

PApaver cornutum flore luteo, Yellow Herned Poppey, G. 367. In maritimis.

\* Papaver cornutum flore violaceo, G. 367. papaver corniculatum violaceum CC. In the Corn fields as you go to Burmel beyond Swasbam.

Papaver Rhaas, Red Poppey or Corn Rose, G.

371. P. 363. Inter fegetes.

Papaver Rhæas flore purpureo saturo, Murrey flowr'd Corn-Rose, In my own Corn at Darfield, Mr. Stonehouse. Ph.

Papaver fylv. G. 370. Inter fegetes.

Parietaria, Pellitory of the Wall, G.331. Pariet, vulg. P. 437. In muris antiquis & ruderibus.

Parenychia vulg. Common Whitlow-grafs, G. 624. Paronychia Alfinefolia, P. 556. In miris & tectis passim.

Paronychia folio rutaceo, G. 624. Paronychia foliis incisis, P. 556 cum priore, sed rarius.

\* Paronychia Gnaphalii facie. At Clapdale in

Tork fire in the mid way betwixt Setle and Ingleborough hill on the rocks.

Pastinaca latitolia sylv Wild Par nips, G. 1025.

P. 944. Ad agrorum margines.

Pastinaca sylv. tenuifolia, Wild Carrot or Birdsneft, G. 1028. P. 901. In agris.

· Pastinaca latifolia sativa, Garden Par nips.

G. 1025. P. 944. CC.

lepi-

Rica

Tri-

Hore

46,

OT N.

pa.

the

ond

G.

ret

147-

et

DS,

G.

nu.

bia

S.

in

rk.

Pastinaca sativa tenuifolia, G. 1027. Pastinaca fativa lutea tenuifolia, P. 901. Tellow and

White rooted Carrot, CC.

Pastinaca sativa atro-rubens, G. 1027. Pastinaca fativa altera tenuifolia atro-rubens, P. 901. Hæ tres Pastinacæ seruntur in agris juxta oppida majora & urbes.

Pellen veneris, five fcandix, Sheepherds needle, or Venus comb, G. 1040. Scandix vulg. five

pecten Veneris, P. 916. Inter fegetes.

Pedicularis, Lons wort, or Red-Rattle, G. 1071. Pedicul, pratenfis rubra vulg. P. 713. in

pratis humidis, variat flore albidiore.

Pedicularis Tragi 97 cubitalis est altitudinis & folis Quercinis. Two miles East of Croydon below the windmill nigh the fish pond.

Pintaphyllum rubrum palustre. Marsh Cinquefoyl, G. 987. In the ditches betwixt Godli-

manin Surrey and the Almes-houses.

Pentaphyllum incanum, Hoary five-leav'd Grafs, G. 988. Pentaphyllum minus repens lanugi-

nolum, P. 399. in montolis.

Pentaphyllum five quinque folium Tormentillæ facie, in agris ficcioribus, & muris, G. 988.

P.

P. 399. Pentaphillum seu Quinquesol vulg. G. 987. pentaph. majus luteo store, p. 396. Ubique ad agrorum margines.

Pentaphyllum supinum Potentillæ facie, G. 988; P. 399. In Tuddington field and near Hide.

Park.

Peplus feu Efula rotunda, Petty Spurge, G.

503. P. 194. inter feget es.

Percepierre Anglorum, Parfley Break - Stone, G.
1594. polygonum felinoides, P. 449. Colickwort ab Herefordiensibus. In agris siccioribus.

Perfoliata vulg. Common Thorongh-wax, G. 536. P. 580. inter segetes sed rarius.

Perfoliata siliquosa, G. 536. Perfoliata sive Bras. fica campestris, P. 580. At Darfield, Mr. Stonebonse, Ph.

Periclymeum, Wood bind or Honey Suckles, G 891. Periclym. five Caprifolium vulg. P 1460. in fepibus & fylvis.

Periclymenum fol. Quercims non procul Oxo-

nio. Mr. fenner.

Persicaria Hydropiper, Arsmart, G. 445. Persicaria vulgaris urens, P. 857. In locis humidis

Persicaria maculosa, Dead or Spotted Arsmatt, G. 445. Persicar mitis sive maculata, P. 8578 cum priore.

\* Perficaria longissimo & angusto folio, At

Mariyns Abbey.

Persicaria pusilla repens. G. 446. P. 857. in sostis. Persicaria siliquosa, G. 446. Mercurialis sylv. noli me tangere dicta, P. 296. Within a mile of Montgomerie at Gwern Dhee, Mr. Morgan,

Morgan , and Near Bingley York Shire. \*Perficaria foliis Salicinis, In a pond in St. fame ('s Park.

Berficaria mitis maculofa spica viridi, at

Kine frome upon Thames.

efol

P

188

de-

G.

G.

Ck-

ıſ.

l.

G

P

)

ì

Patafitis, Butter- Burr, G. 814. P. 419. ad

Huminum & rivulorum ripas.

Pencedanum, Sulphur wort, G. 1054. Peuced. vulg. P. 880. Hoggs-fennel, betwixt Feverham and the Sea, and at whit fable in Kent. Pincedani facie pufilla planta Lob. Selinum pu-

milum montanum Clus. Dwarf Hogs-fennel, or Rock-Perfley on St. Vincents Rocks, G.

1053. 3. Peucedanum minus, P. 880.

Philaris, Canary-feed, G. 86. Phalaris vulgaris. P. 1163. juxta oppida ex seminibus ejectis.

Phillitis, Harts-tongne, G. 1 138. Phyll. five lingua Cervina vulg P. 1046 in asperis & umbrosis. \* Phyllitis multifida, G. 1138. Phyllitis laciniata, P. 1047. Near Southampton, ex fenel-

cente planta finduntur folia, ideoque forsan

a priore non distinguenda.

Pilofella major, Great Monle-ear, G.638. Pilofelaltera erecta, P.690.ad agrorum margines. Pilofella repens, G. 638. Pilofel minor vulg. repens. P. 690. ad pratorum & fossarum

ficciorum margines.

Pilofella filiquata, On the dirch fides in the way to Marybone, and in a Close on the left hand of the lane from Islington to King sland. Iloret mense Martio, vide optimam iconem apud Camer, pag. 7. Huic addit C Capp. minorem. PimPimpinella fylv. Wild Burnet, G. 1045; Pim pin. vulg. major, P. 582. in pratis.

Pimpinella Hortensis. G. 1045. Pimpin, vulg. five minor. Common Burnet, P. 582.

Pimpinella Saxifragra, Burnet Saxifrage, G. 1044. Pimpin. Hircina major, Great German Burnet Saxifrage, P 947. In pratis & pascuis.

Pimpinella faxifraga minor, G. 1044. Pimpin, Hircina minor, P. 947.

\* Pimpinella Saxifraga major nostras, P. 946

in pratis frequens.

Pinguicula five Sanicula Eboracenfis, Butterwort, or York Shire Sanicle, G.788. P. 532 plentiful about Oxford, and neer Chifelburt, which Dr. Bowles planted there.

\* Pinguicula minima fl. albo. In the mid way betwixt Oakhampton and Lance fon Cornwall betwixta great wood and the River in boggy

Meadows

Pinns, The Pine tree, G. 1356. P. 1534. ina wood on the left hand of Hamstead, called the Pine walks.

Pifum arvense flor, albo, aut roseo, fructu albido aut cineritio White and Grey Peafe, G. 1219.P. 1059 quamplurimæ habentur fpecies in fatis,

Pisum spontaneum maritimum Anglicum, P. 1060. G. 1250. in Norfolk, and on the West of Hastings in Suffex on the Beach.

Pifum aliud maritimum Britannicum, P. 1060.

G. 1250. on Effex floar.

\* Pisum in extremitate clavatum modo Trifolii fiquofi. At Bingley York hire.

Planta-

Plantago aquatica major. Great water Plantain-G. 417. P. 1245.

Plantago aquatica minor ftellata, G. 417. Plantago minor muricata, P. 1245. in Ditton-Common plentifully.

Plantago aquatica humilis, G.417. P. 1245. in

foffis.

im

ilg,

G.

tan

nis.

in.

46

er-

26

rt.

ay

II,

12

d

0

S,

i

Plantago incana, G. 419. Flantago maxima &

flore incano, P. 493. In pratis.

Plantago latifolia, G. 419. Plantago latifolia vulg. common Waybread or Plantain, P. 493. Plantago marina, G. 423. Plantago marina vulg.

P. 498.

Plantago quinquenervia, Rib-wort Plantain, G. 422. Plantago angust major & ferrata, Great Ribwort with plain and dented leav's, P. 496. Plantago panniculis sparfis, G. 420. P. 494. in

the Ifle of Thanet, but very rarely.

Plantago quinquenervia rosca, G. 422. P. 496. below Hamftead Church in the way to London, just against the Chefnut walks.

\*Plantago Aquat, minima Cluf. P. 1244. fine icone. On Hunflow heath, a mile off that

Town in the Way to Coln- Brook.

\* Plantago aq. major muricata. In a fmall pond betwixt Clapham and South Lambeth-Common. Mr. Morgan.

\* Plantago aq humilis flore purpureo . At Taxton five miles from Godmanchester in the way.

to London plentifully.

Pneumananthe, Calathian Violet, G. 438. neer Brunley in Lancashire. In a Close call'd Banierfts heal, plentifully.

Paonia famina, Female Paonie, G. 981. P. 1380. in a Close belonging to Mr. Stevenson at Sanhing-well in Bark shire of above fifty years standing, and in Stancomb wood near Winch. comb, Gloucester-shire.

Paenia mas, G. 980. P. 1381. in Mr. Field; well-close in Darfield, Mr. Stonehouse.

Polygala, Milk Wort, G. 563. P. 1332. variat flore caruleo, purpureo, violeceo, carneo, rubro, albo, in montosis siccioribus.

Polygonatum, Solomons Seal, G. 903. Polygo. natum majus vulgare, P. 696. in fylvis.

Polygonatum latifolium, 2. Clus. Sweet-smelling Solomons Seal, G.904. Polygon, majus vulg. P. 600 Ph.

\*Polygonatum bacca nigra simplici & flore unico sinibreato virid. At Clapdale in the mid way 'twixt Settle and Ingleborough hill in Tork-foire.

Polygonum mas vulgare, Common Knot-grafi, G. 565. Polygonum mas vulg majus, P.443. ubivis obvium.

\* Polygonum marinum, Near Lewis in Effex.

Polygonum Selinoides five Knamel, Perfley Piirt, G. 566. Polyg. Germanicum five Knaml, Knot-grass, or Knamel, P. 447. Inter segence.

Polygonum Serpillifolium, Small round lead Knot-grafs, G. 566. Polygon serpillifolium Lob. P. 1680. In maritimis.

Polypodium, Wall-fern, or Polypodie of the Wall, G. 1132. Polypodium vulgare, P. 1039. in quercubus & muris an tiquis.

Popu-

Populus alba, white Poplar. G. 1486. P. 1410.

Populus nigra, G. 1486.P. 1410. juxta rivulos. Populus Libica, The Aspen-tree, G. 1487. P.

1411. in sepibus & sylvis humidis.

portulaca sylv Wild-Purslane, G. 521. P. 723. about Ramsey mear, in the soot-paths by the Rape-mils, CC.

Phimogeiton latifolium, Broad leav'd Pondweed, G. 821. Potamog, sive fontalis major latifolia

vulg. P. 1254.

Potamogeiton angustifol. G. 821. Potamog. sive fontalis major longifol. P. 1254. In aquis stagmantibus, at the farthest end of Hunstow-heath. Potamogeiton 3. Dodon, Small Pondweed, G. 822. est perfoliatum. In fluviis.

Phamogeiton longis acutis foliis, G. 822. in flag-

mis.

0.

m-

6.

lds

iat

0,

0-

inf

ılg,

ico

ray

rk-

is,

43.

Pt-

vel,

tes.

n'd

mu

all,

10

PH-

\* Potamogeiton Ramosum caule compresso folio graminis canini, ND, in the River Cam near Cambridge. CC.

Potamogeston Pusillum folio gramineo, caule rotundo, in Thames near Oxford, CC.

Inamogeiton Capillaceum capitulis ad alas trifi-

\*Potamogeiton latifol flore albo, at Noftal-Abby four miles from Wakefield Tork-foire.

Potamogeiton affine gramen aquaticum, Water-

grafe with Small crooked Cods, CC.

Primula veris major, Field Comslips, G.780. in pratis.

Primula pratensis inodora lutea, Field-oxslips, G. 780. I Pri-

Primula veris flore rubro, Red Birds-eye, G, 783. Paralytica Alpina minor. The less Birds-eye, P. 536. variat flore albo & purpareo, in divers moist mountainous places of the North, and in a very low and squally meadow neer Knaesborough.

Primula veris Heskethi, Mr. Heskeths Primroft, G. 782. versipellis Chamæleon, Or Varion Primrofe, in Clapdale Wood three miles from

Setle in York- Bire.

Primula veris minor, Field Primrofe, G. 781.

Primula veris vulg. P. 535.

\* Primula veris Polyanthos, multis infignitur floribus, quoad colorem & formam exactè refer Primulam vulgarem, fed compacti flores celecunt ad instar primulæ pratensis luteæ modoræ, In great Woolver Wood in Warwick-hin, Mr. Morgan, qui transtulit in hortum sum instructissimum Westmonasteriensem.

Prunella, Self. heal, G. 632. P. 526. In pratis,

variat flore albo.

\* Prunella fl. carneo. Near Crediton Devon fin in abundance.

Prunus sylv. The Sloe-tree, G. 1497. The Black thorn or Sloe-tree, P. 1033. In sepibus & dumetis.

Pranus sylv. fructu majore, The Bullattu.

In fepibus.

Pseudomelanthium, Bastard Nigella, or Cockle. G. 1087. Lychnoides segetum sive Nigellastrum, P. 632. cum superiore.

Ptarmica, Snees-wort, G.606. Ptarmica vulg.P. 859. In pratis humidis.

Ptarmica duplici flore, G. 606. By Chilmark in wilts, and many other places.

Pulegium regium, Penny-Royal, G. 671. Puleg.

vulgare, P. 29. in locis humidis.

Pulegium densis furculis, In Wales, P. 29. 3. uall Pulmonaria foliis Echii , Bugloss Comflip, G. 808. By Holbury house in the new Forest in

Hampshire. Mr. Goodyer.

Pulmonaria Gallica five aurea latifolia, Broadleav'd, French or Golden Lung Wort. G. 304. Pulmonaria Gallica angustifolia, G. 304. Both these in the meadows on this side Hampstead. Pulmonaria maculosa, Spotted Cowslips of Hieru-Salem, G. 808. Neer Kings. Wood in Hamp-Shire, Mr. Loggins. Ph.

Pulsatilla rubra, Red Pasq-flour, G. 385. P. 341. On a Heath towards Barneck three

miles from Stanford, Dr. Bewles.

Pulatilla vulgaris, Purple Pafq-flour, G. 385. P. 341. At Hilder ham fix miles from Cambridge.

\* Pullatilla fl. Cæruleo. Neer Pontfract York-

bire.

le, G.

leffa

urpo.

es of

arofe.

crion

from

781.

r flo-

efert

cref-

odo.

bire,

uun

atis,

hire

ack-

5 &

rtti

kle.

12.

P.

17-

Pyrola, Winter-green, G. 408. Pyrola nostras vulg. P. 508. at North-bridge half a mile

from Halifax plentifully.

\* Pyrola alfines flore minor. On the East end of Rumbles. Moor neer Helwick in the bogs amongst the rushes, above a great Brook, CB. prod, 100. P. 509.

Pyrus fylv. variorum generum, Hedge-pears, or Choak pear, &c. G. 1459. P. 1500.

Quer-



# Q.

Quercus vulgaris, The Oak-tree, G. 1339, Quercus nostras, P. 1386.

\* Quercus serotina, processor foliis fructuq minoribus Dor-Oak, plentiful on Liuma bill in Bramshaw Parish Wilts. Ex Mss, Mr. Goodyer.

Querous natalitiis Christissorens, at Glassenburg Abby, and in the new Forest in Hampsbire. Querous marina, Sea-Oak, or Wrack, G. 1567.

Quercus marina Herbacea, P. 1293.

\* Quercus marina fol. inferioribus fimbriais fæniculaceis. At Bass Castle in Cornwall.

Quercus marina secunda, Sea-thongs, G. 1568. fucus marinus Dod. P. 1293.

Quercus marina tertia, The third Sea-wrack, G. 1568. cum priore.

\* Quereus marina longiore & latiore folio.

Querous marina quarta, Jagged Sea-wrack, G.

\* Quercus marina angustisolia in extremitate referens Chelas Cancrorum, an quercus mirinæ varietas, G. 1567. 4? P. 1293.

## R.

R Adix cava minima flore viridi, Small Bunnikins hollow-wort, G.1091.P.288. Hollow-Root, In the bank of a hedge by a grove of Elms, on the North-side of Chesterton Cambridge-Shire.

339

auq:

MIP HOL

, Mr.

bury

ire.

567.

riatis

68.

rack,

G.

itate

ma-

Agix

Ranunculus aquatilis, Water Crewfoot, G. 829.
Ranunc. Aquaticus Hepaticæ facie, P. 1216.
Rannnculus palustris rotundisolius, Marish
Crow foot or Spear-wort, G. 962. Ranunc.
palustris Sardonius lævis, P. 1215. in stagnantibus.

Ranunculus flammeus major, Great Spear-wort, G. 961. Ranunc. palustris flammeus major, P. 1215. juxta aquas variis in locis.

Ranunculus flammeus minor, The jagged Spearwort, G. 961. Ranunc. palustris flammeus minor seu angustifolius, P. 1215.

Ranunculus flammeus ferratus, Jagged Spearwort, G. 962. P. 1214. 3.

Ranunculus auricomus, G. 954. Ranunc. nemerolus dulcis secundus Tragi, Sweet-wood Crowfoot, P. 326. in dumetis & sylvis copiosé.

Manunculus bulbosus, G. 953. P. 329. in praus & pascuis.

\* Ranunculus bulbocastani radice, medium radicis crassius longius que reliqua, ad cujus extremitatem oritur bulbus & ex eo nova nascitur planta. Below the City of Exeter.

Ra-

Ranunculus globosus, Locker Gowland, or Glob Crow-foot, G.955. P. 331. in divers places of Tork-shire and Lancashire.

Ranunc, hirfutus arvensis fl. minimo C Cappi

\*Ranunc. rectus fol, pallidioribus hirsutis JB. In locis humidis præcipuè ubi aquæ per hiemen stagnarunt. Below Hamstead in the Meadow betwixt the Town and Heath.

\* Ranunculus bulbosus fl. pleno. Betwist the two Chemes in a hollow place on the lower

fide of Banftead Downs.

\* Ranunculus hirfutus annuus fol. Gerani Columbini, at the farther end of the bridge betwixt Micham and Casalton, and at them

of Camberwell.

Ranunculus Hederaceus rivulorum se extendens atra macula notatus, C.B. Ivielaria mater Cromfoot, In the entrance of a little lane leading out of Cotton road up to the bowling-green beyond Cambridge Castle, CC.

Ranunculus surrectis cauliculis, G. 951. Ranunc. pratenfis erectus acris vulg. andulis

P. 328. In pratis.

Ranunculus pratenfis etiamque hortenfis, Common Crowfoot, G. 951. Ranunc. Pratenfiste pens, P. 328.

Ranunculus arvorum, G 951. P. 327. inter fegetes ubique, Rough headed Crowfoot.

Raphanus aquaticus, G. 240. Raphanus Aqua. Lob. P. 1228. in fossis, folia in aliis magisia aliis minus dividuntur.

Raphanus rusticanus, Horfe-Radish, G. 241.P.

860.

860. in locis ubi ex hortis radicum particulæ projectæ fuerunt.

Rapistrum arvorum, Charlock or Chadlock, G.

233. P. 862. in agris.

Globe

ces of

capp.

B. In

men

down

the

19WO

Co

e be-

end

tten-

avid

little

M.

3.

Ra.

lcis,

010

sre-

nter

pat, is io

60.

Rapistrum arvense alterum minus soliis minus

incisis lenioribusque, G. 234. 2.

Rapistrum luteum siliqua hirsuta articulata, Charlock with a rough joynted Cod, inter segetes CC.

Rapiftrum luteum filaqua glabra articulata,

cum superiore. CC.

Rapistrum flore albo, as common in Corn fields in Surrey as the Yellow. Mr. Brown uti etiam in pluribus Cantii locis.

\* Rapistrum aliud non bulbosum, P. 862, in the broad street by White Chappel, Mr. Goodger.

\*Rapum ma jus & alterum radice oblonga, Great and longish rooted Turnip, G. 232. seruntur in agris juxta civitates & oppida ma jora.

Rapum sylv. Wild Turnips, G. 233. Rapum sylv. non bulbosum, The mild Rape or Tur-

mip, P. 862. fown in the Isle of Ely.

Rapunculus nemerosus, Wood Rampions, G.

454. P. 650.

\* Rapunculus sylv. flore rubro albescente. In the pastures and hedge sides on the Northwest of the Moor not far from the great bog neer Peters-field, Mr. Goodyer.

\* Rapuntium fl. purp. At Effaton a mile from

Wigmore, Herefordshire.

Rapunculus corniculatus montanus, G. 455. Between Selbury hill and Beacon hill in the

1 4 way

way to Bath, and in the Chalkey hills by

Maple-Durham Hampshire.

Rapunculus Scabiosæ capitulo cæruleo, Hairy Sheeps Scabious, P. 646. On the Downs in Surrey.

Rapuntium parvum, G.453. in sylv. & umbross. Reseda Plinii, Italian Rocket, G. 277. Reseda

minor feu vulg. P. 823. In montofis.

\* Reticulum marinum viride & album. Tis of the thickness of Cap paper six inches long and four broad, and seems to be made up of Plants matted together. Both were found on the Beach near Dover.

Rhamnus folutivus, Buck thorn, G. 1337. Rham, folutivus, seu spina insectoria vulg. the common Purging-thorn, P. 243. near Lewsham in Ken,

Rhamnus secundus Clus. Sallow-thorn, G. 1334.
Rhamnus primus, Diosc. Lob. sive littoralis,
P. 1006. betwixt Sandwich and Deal,

Rhodia radix, Rose-root, G. 532. Rose-wort, P. 727. On Ingleborough by the Brook sides in Lancashire, and on the moist Rocks in Farfels in the bottom of Lancashire on the highest Fell towards Hawks-head.

\* Ribes fructu nigro, Black Currans, P. 1562. Ribes tertium Gerardi folio & fructu male olentibus, G. 1593. 3. Squinancie berries, By

the Rivers side at Abbington. CC.

\* Ribes fructu parvo, At Wimbleton in Surre,

and many places of Lancashire.

Rosa solis folio rotundo, Sundem with round leav's, G. 1556. P. 1051 and Red rot in uliginosis. Rosa Rosa solis foliis oblongis, G. 1556. Rosa solis sylv. longisolia, P. 1052. Mr. Heaton sound some with leaves a span long plentisully in a bog by

Edendery.

in

1,

la

of

g

of

ñ

H

n

V

1

Rosa sylv. odora, Eglantine or sweet Bryer, G. 1269. in divers hedges by Darsield but with more smooth shining leaves than the Garden kind. This is like that of P. pag 1017. but the leaves are sweet as well as the flowr, Mr. Stonehouse.

Rosa canina inodora, the Bryer-rose or Hep-tree, G. 1270. Rosa sylv. inodora sive canina, P.

1017, in sepibus & dumetis.

\* Rosa sylv. odora flore duplici, Double Eglantine, G. 1270. invenitur duplici, & triplici serie petalorum, in sepibus præsertim prope Wigorniam, Mr. Brown.

Rosa Pimpillæ solio, The Pimpernel Rose, G.
1270. Rosa Pimpinellæ sol, sive pomisera

minor, P. 1018.

\* Rofa Pimpinellæ folio flore Rofæ mundi vari-

egato, On Barns common.

\* Rosa pimpinellæ fol, fl. rubro. In some barren fields near Worcester, Mr. Brown and in a barren field at Church-lench four miles beyond Eversham in great plenty.

\* Rosa Canina suavissimi odoris. In Brodards field going over hell ditch not far from the

River Coln, Lancashire.

\*Rosa Canina sylv. unico fl. & fructu. In the fields near Hackney in the way thence to London.

Rorif-

Rorismarinum sylv. Wild Rosemary, G. 1292, Rorismarinum sylv. minus nostras, P. 76. in Shrop-shire at Birch in the Moors of Elles. mere and at Bulsover York-shire at a place call'd the Hayslacks.

Rubia minima, G. 1120. Rubia minima faxati-

lis, P. 277. on St. Vincents Rocks.

Rubia fylv. G. 1118. P. 274. 2.

\* Rubia pratensis minor carulea, P. 276. inter

fegetes, CC.

Rubia Tinctorum Red Madder, G. 1118, Rubia major fativa, P. 274- on St. Vincents Rocks and at Aberdoy in Merioneth-shire, 'ils planted in the Fens.

Rubia marina, G. 1119. P. 275. 6.

Rubia Cynanchia, Squincy-wort, G. 1120.5.
Rubia Cynanch. faxatilis, P. 453. 2. incollibus cretaceis.

Rubus Idaus, The Raspis Bush, or Hind berry, G. 1272. in many places of the North and Wales, and 'tis as common as brambles in the Woods of Chesham-Bois in Bucking hamshire.

\* Rubus scandens instar Viorna. By Maider

head and near Slough.

Rubus repens fructu Cæsio, Small Bramble Demberry, G.1271.2. in arvis nimis luxuriat.

Rubus, The Bramble-bush. G. 1272. Rubus major vulg. Bramble or Black-bush, P.1013, in sepibus & dumetis.

\*Rubus morus, The Mulberry Bramble, so call'd by the Countrey people at Sutton in Effex.

Rubus fructu Cæsio magno non repens In

London road half a mile on this fide Mayden-

head Bridge.

\* Rubus suavis fructu rubro. In the woods of Mr. Parker call'd Runcle-hurst, and on the wals in Lancashire.

Ruscus five Bruscus, Knee holm, or Butchers-Broom, G. 907. P. 253. in sepibus & dume-

tis.

92,

· in

lef.

ace

ati-

iter

Q-

nts Lis

5. ol-

he

le

ıt.

115

d

### S.

Sagittaria major, Great Arrow-head, G.416. Sagittaria major latifolia, The greater broad Arrow-head, P.1247. In aquis stagnantibus. Sagittaria minor, G. 416. Sagittaria minor latifolia, P. 1247.

\* Sagittaria longiffimâ cuspide pedali, Arrowhead with a leaf a foot or more long. In the

ditches about Oxford, CC.

\* Sagittaria fine capitibus ex cujus radice prodeunt caules sex vel septem spithamas longi ferentes virides nodos ad instar nucum juglandum; hujus solia deceperunt doctissimum botanologum Casp. Bauhin qui in prodromo vocat gramen aquaticum, 'Tis sound near Hampton Court, solia insigniuntur iisdem Characteribus quibus utuntur Musici in describendis cantilenis ut in aliis sagittariis.

Salix pumila, five Chamæitea, The Dwarf Willow, G. 1391. Salix pumila angustifolia

recta, P. 1434.

Salix Caprea latifolia, G. 1390. Salix latifolia oblongior, P. 1432.

Salix Caprea rotundifolia, G. 1390. Salix lati-

folia rotunda, P. 1432.

Salix humilis repens, G. 1391. Most of these Willows grow on Hamsted heath.

Salix humilis, G. 1391. Salix humilis latifolia

& Alpina repens, P. 1433.

Salix rosea Anglica, G. 1390. P. 1430. Hanc non esse salicis speciem à reliquis distinctam pro comperto habet CC.

Salix, The common Willow, G. 1389. Salixar. borea angustifolia alba vulg. P. 1430.

Salix Aquatica, The Ozier or mater Willow, G. 1389. Salix aquat, humilis, P. 1434.

Salix folio laureo sive lato glabro odorato solio, N D. Willow-bayes, at Wolverhampton.

\* Salix folio laureo odorato angustiore, a quarter of a mile on the River side from Creditor

Devonshire seven miles from Exeter.

\* Salix cortice intus flavo, fol. angusto, N.D.

The bark is as yellow as that of Barberies, at

Rippon and divers other places of Tork-shire,

An salix humilior soliis angustis subcarules

ut plurimum sibi invicem oppositis, CC. De
his salicibus videbis, CC. qui optime distinxit

& descripsit.

Salvia agrestis sive Scoridonia, Wood Sage, G. 662. Salvia agrestis sive scordium alterum,

P. 111. in aridis pascuis & sylvis.

Sambueus aquatilis five palustris, Water Elder, or Maristo Elder, G. 1424. Sambucus rosea & aquatica sive palustris, P. 209. Sam-

Sambuens, The common Elder-tree, G. 1422. Sambucus vulgaris, P. 208. In sepibus.

\* Sambucus foliis laciniatis, G. 1423. P. 280.

Near Briftol, Dr. Joyliff.

Sanicula sive Diapensia, Sanicle, G. 948. Sanicula vulg. five Diapenfia, P. 532. In fylvis & umbrosis sepibus.

Saponaria, Soap wort, or Bruis-Wort, G. 444.

Saponaria vulg. P. 641.

Saxifraga alba, White Saxifrage, G. 841. Saxifraga alba vulg. P. 424. in pratis humidis, plentifully in St. Georges fields.

Saxifraga aurea, Golden Saxifrage, G. 841. P.

in locis humidis 425.

Saxifraga aurea major foliis longius incedentibus, hanc belle depingit hortus Eystetensis. Near Hedley Hampshire. Mr. Brown.

Saxifraga pufilla graminea flore parvo herbido

& muscoso, on the walls at Chefwich.

\* Saxifraga fl. pleno. On the West side of In-

gleborough hill.

Saxifraga Anglicana alsine folio Chick weed, Break fone, G. 567. Saxifraga Angl. Occidentalium Lob. P.427. Between Guilford and

Compton in Surrey.

Saxifraga palustris alfinefolia, small mater Saxifrage, G. 567. Saxifraga paluftris Anglicana. P. 427. Marish Saxifrage, now commonly call'd Pearl worts whereof I have three forts from Bass Castle Cornwall.

Saxifraga Anglicana facie Seseli pratense, English Saxifrage, G 1047. Seseli pratense nostras \* Sax

P.904 in pratis & pascuis.

\* Saxifraga antiquornm Lob. G. 605. By lower Cheme in a Chalk pit near Banfted Downs on the left hand from the Town.

Scabiosa major vulg Great common Scabious, G. 719 Scabiosa vulg. pratensis, P.484. In satis, & pratis: variat flore cæruleo aut albo.

Scabiosa minor sive Columbaria. G. 719. Scabiosa minor campestris, P. 484. In montoss,

variat colore purpureo & albo.

Scabiofa omnium minima, Sheeps Scabions, G. 722. Scabiofa minima ovilla dicta, P. 485. Invenitur flore albo.

Scabiofa media, Middle Scabions, G. 720.

Scabiosa campestris sive segetum, G. 720. Sabiosa major segetum, P. 485. 3.

\* Scabiosa vulg fl. pleno. Betwixt Redbourn and Stony-stratford plentifully. CCapp.

Scordium, Water Germander, G.661. Scordium legitimum, P. 111. abundantly only on the banks of the Isle of Ely, and neer Aristotles Well at Oxford.

Scrophularia major, Great Figg. wort, G. 716. Scrophul. major vulg. P. 610. In sylvishu-

midis.

Scrophularia flore luteo, G. 717. P. 611' In Oxfordshire.

Secale, Rye, G. 68. Secale vulgatius, P. 1128. Seritur in agris.

\* Secale semine nigro. At Leeds in York Shire.

Sedum, seu sempervivum majus, Great Houlleek, G. 510. Sempervivum majus vulg. P. 731. In tectis, & muris antiquis.

Scaum

sidum five Aizoon scorpoides, G. 513. vermicularis scorpoides, P. 734. cum priore.

Sedum Alpinum Clus. G. 516. Sedum Alpinum flore pallido, P. 737.

Sedum Divi Vincentii, N D. Mr. Goodyer.

Sedum medium teretifolium, Small Prick-madam, G. 513. vermicularis minor, Small Stonecrop, P. 734. in locis saxosis & asperis.

Sedum minus officinarum, G.512. vermicularis

five craffula minor vulg. P. 734.

Sedum tertium Diosc. sive illecebra minor, Wall-Pepper, P. 735. vermicularis sive ille-

cebra minor acris. G. 517.

Sedum tridactylites five Alpinum trifido folio, Handed Housleek. Tridactylites Alpina JB. tom. 3. pag. 762. All these small Sengreens are said to grow on the high mountains of Wales. This I had from the black mountain in Herefordshire

Sedum minus Hæmatodes, Prick-Madam, G.

512.

er

n

S,

j,

Ħ

n

S

\* Sedum five illecebra fol. oblongis. On the North fide of Ingleborough hill, neer a bog by the fide of an underground River.

\* Sedum minimum fl. mixto ex albo & rubro.
On the West of Ingleborough hill in the mud

of the hollow topped stones.

'Sedum feratum rotundifol. The Princes fether.
On Snowdon bill both on Rocks and Wellsprings.

\* Sedum majus Portlandicum arborescens Angl.

five maritimum Adv. pag. 162.

\* Sedum

Sedum seu Vermicularis flore albo, at Absum below Exeter and at Oakhampton Devensire

plentifully.

Selinum Sii foliis, Hone-wort, G. 1018. Selinum fegetale, Corn Persley, P. 932. On the bank of Thames as you go from the Duke of Norfolks Garden to Lambeth.

Serpyllum citratum, Limon Time, G.57.1.P.8.In montolis, at Hockliff in the hole near Oxford,

Serryllum hirfutum, G. 571. Serpyllum latifolium hirfutum, P.S. at Bradbourn in Yorkibire.

\* Serpyllum vulg. flore albo, Plentifull betwix

Tuddington and Hampton Court.

\* Serpyllum vulg. primum genus J B. CCapp.

\*Serpyll. magnum latifolium, a mile of Bradford in the way to Hamorth, and at Sarefden betwiest the house and the pond in Tork-shire.

\* Serpyllum fætidum, two or three miles from Petersfield on the Chalky downs. Mr. Goodyer.

Serpentaria minima, five Coronopus. Small Bucks horn Plantain. G. 426. holosteum minus, five serpentaria minor, P. 499. In maritimis.

Serpentaria major, sive Holosteum angustisolium majus, P. 499. Coronopus, Sea Buk

horn, G. 425. in maritimis.

Serratula purpurea, five alba. Sam mort with purple or white flowrs, G. 713. Serratula vulg. fl. purp. vel abo. P. 475. In Shadowy fides and hedges.

\* Sesamoides Salamanticum magnum. Great Bafrard Wood, G. 493. Muscipula Salam. major, the greater Spanish catch fly, P. 636. sine

icone

icone. Near the gravel pit as you go to the nearest Windmill on the North side of New-Market Town.

Sederitis Anglica strumosa rad. Clowns woundwort, P. 587. panax Coloni, Clowns All-heal, G. 1005. juxta fluvios fossasq; Londinenses, & alibi per totam Angliam.

Sideritishumilis lato obtuso solio, G.699. Sideritis alsines Trissaginis solio, P. 586. nulla icon vel apud G. vel P. In many places of Hamp-

Bire.

12

n

k

.

'n

d.

į.

e.

IC.

1

.

m

11

5,

s. '

k

h

la

y

4.

le

10

Sideritis arvensis ruba, P. 587. 13. Tetrahit angustifol. Hist. Ludg. — In the Corn fields in Kent, and at Purbeck in Esex.

Sinapi sylv. minus, Small wild Mustard, G. 245.
P. 831. On the banks of Ditches newly cast

up, and amongst Corn.

Sinari Genevense sylv J B. tom. 2.858. Beyond

Southwark in the way to Lambeth.

flard, G. 242. 2. fine icone an finapi fativum rapifol. P. 831. On the Ditch fides in many places about London.

Sissum vulg sive Amomum German. Small mild Parsley of Germany, P. 914. Petroselinum Macedonicum Fuchsii, G. 1016. ad fossa &c

lepes.

Simm majus latifolium, Great mater Parsnip, G. 256. Sium Diosc. sive pastinac, aquat, major. P. 1240. In the Ditches by Reading and in the River Cam near Cambridge.

Slum alterum Olusatri facie, G. 256. Sium ma jus

R alterum

alterum angustifol. P. 1241. In piscinis& aquis stagnantibus sed rarius.

Sium minus five Pastinaca aquat minor. P. 1240. Sium umbellatum repens, G. 256.3. In fossion & stagnis ubique.

Sium majus angustifol. G. 256. 2. cum priore.

\* Sium vernum foliis fæniculaceis, On Punny Common near Wandfor in standing pits by the Road side.

Sium latifolium foliis variis, At Marlborth Wilts, Mr. Bobart Junior.

\* Siam pusillum fol, variis, In Surrey near

\* Sium umbellis ad caulium nodos, At Petersfuld. Mr. Goodyer.

Sium medium foliis elegantissime dissedis, la fome Ditches about Oxford.

Solanum hortense, Garden Night-shade, G.339. Solanum vulg. P. 346. In fimetis & ruderibus.

Solanum lethale, Dwale, or deadly Night-shade, G.340. P.346. In locis umbrofis, & arenosis.

Soldanella mar. Sea Bind-weed, G. 838. Soldan, vulg. & major maritima, P. 168. passimin maritimis.

Selidago Saracenica, Saracens Consound, G.419.
Solidago Saracen.major, P.539. 2. sine icone.
On the 5. mile bank near Wilt-sea, and between Dudson and Gwarthlow, and other places. Dr. Bowles.

Sonchus asper, prickly Som thistle, G. 291. Sonchus asper laciniatus, P. 804. 3. ubique

ferè obvius.

Son-

S

5

So

So

So

Spi

\$5

lis

0.

is

e7 he

W.

ar

14.

In

39.

NUS.

de ,

Sis.

lan,

n in

29.

one.

and

291.

ique

Son-

suchus Izvis, Hares Lettuce, G. 292. Sonchus Izvis vulg P. 805. cum priore.

Smehns arborescens alter, G. 204. P. 807. Super muros.

Souchus folio finuato hederaceo, In dark and fhady Lanes.

Smithus fl. albo medio luteo. Mr. Stoneboufe,

'Sonches foliis cauleq; Aristochiæ Clematicis, In adark Lane at the Bridge end a mile from Briftol Ferry.

\* Sonchus tricubitalis fol. cuspidato, In the Meadows betwixt Woolwich and Greenwich by the banks of Thames.

Sophia Chirurgorum, Flixweed, G. 1068. P. 830 ad agrorum margines & in locis in cul-

Subm the Service-tree, G. 1471. Sorbus ligitima.
P. 1420. In fylvis & sepibus, præsertim in
Sussexia.

Subm Torminalis, G. 1471. Sorbus torminalis vulg. P. 1420. cum priore sed rarius. Both grow in the Pine walks by Hampsted.

Parganium ramosum, branched Bur-reed, G. 45.

Sparganium latifolium, G.45. Sparganium non ramosum, P. 1206. cum priore.

the East side of Screeby nigh a great Wood, where the foot way is cast up Noting hamshire.

Spartum Anglicanum, English Mat-weed, or K 2 Helm, Helm, G. 42. Spartum marinum nostras, P. 1198. in maritimis & falsis.

Spartum Angl. alterum, G. 42. Spartum nosira alterum, P. 1198. in montosis & Ericeis.

Speculum veneris minus, codded Corn Violet, G. 439. the leffer Corn Violet, P. 1331. Inter fegetes.

Spergula marina Dales champ. Sea Spurre, G. 1125. 4. fine icone. Sagina spergula minor, P.

A561. 2. nulla icon.

Sagina spergula, G. 1125. Sagina spergulam. jor, P. 562. inter segetes.

Spergula fl. rubro, icon, Hist. Ludg. 1 179, lub titulo Chamæpuces Plin. In Tuthil-fields new Westminster.

Sphondylium, Com Parsnip, G. 1009. Sphondylium vulg. P. 953. In practis & pascuis pingui

oribus.

Sphondylium majus aliud laciniatis foliis, P.954

In Shroffhire, Dr. Bowles:

Spongia ramosa, branched Spunge, G. 1577.
Spongia ramosa Britannica, 1304. In littoribus nostris.

Stellaria aquat. Water Star-wort, G.830. Starlike water Fennel, P. 1258. ubique in aquok

Symphytum fl. Ceneritio, Every where about Reading vid. Confolida,

7

7

## T.

T Anacetum, Tansie, G. 650. Tanacet, vulg. P. 81. in arenosis & rivulorum ripis, Plentiful by the Thames side near Kingston.

Tap/m barbatus, Mullein, or Higtaper, G.773. Verbascum vulg sive Tapsus barbatus communis, P. 60. ad sepes & agrorum margines

variat flore albo.

, P.

firas

G.

nter

G.

r.P.

ma-

fub

lear

dy.

ur

54

rj.

þ

Taxm, the Yem tree, G. 1370 in Cæmiteriis fylv.
P.1412. & montibus, præsertim Surriensibus.
Taxus tantum florens, G. 1370. On the Chalky

hills in Hampshire, Mr. Goodyer.

Telephium five Crassula major vulg. Common Orpyn, P. 726. Crassula, five faba inversa, G. 518.ad agrorum margines & in locis umbrosis.

\*Telephium semper virens, G. 520. Telephium flor. purpureis, P. 727. On old walls in many

places of the North.

Thalistrum sive Thalietrum majus, Great Baflard Rubarb, G. 1251. Thalic, majus vulg. Meadow Rue, P. 264. in pratis humidis.

Thalictrum minus, G. 1251. P.264 cum priore

& inter fegetes.

Thlaspi Diosc. Treacle Mustard, G. 262. Thlaspi Drabæ solio, P. 830. inter segetes.

Thlashi Hederaceum, G.271.P. 848. in maritimis.

K 3 \* Thlashi

"Thlashi minus Clus, G. 268. Thlashi persoliatum minus, In the Pastures above the Ebbing and flowing Well two miles from Gislewick in stony ground amongst the grass, York shire.

Thlaspi minus Bowiers Mustard, G. 262. In

maritimis.

Thlaspi vulgatissimum, G. 262. Thlaspi vaccariz

folio, P. 836.

\* Thiaspi aliud lunatis foliis sive Magorum Arabum Lob. P.848. On the top of Penigen

bill in Yorkshire, under the stones.

\* Thiaspi vulg. serratifolium, On a Rock over against the Minster at Durham, an Thiaspi mont. bursæ pastoris fructu primum Col.pag. 276.

Tilia famina, the Female Lime-tree, G. 1483,

P. 1407. In areis & compitis.

Tilia mas, G. 1483. P.1407. In sylvis & sepibus \*Tilia ulmisolio, semine Hexagono, At Whista

ble in Surrey, and near Darkin.

Tithymalus Characias amygdaloides, unfavour wood Spurge, G. 500. Tithymalus Characias vulg. P. 186. in fylvis & umbrofis.

Tithymalus helioscopius, G. 498. P. 189. inter

legetes.

Tithymalus Hibernicus, Makenboy.

Tithymalus paralius, G. 498. P. 184. On most parts of our English shore.

\* Tithymali marini species minima ex Cornu.

bia.

Tormentilla, Sept foil G. 992. Torment vulg.P. 394. In Ericetis & dumetis, pascuisq; quibusuam.

\*Tormentilla quadrifolia radice rotunda. At wiggen in Lancalbire, amongst Furzes and in Ditches.

tom

and

e in

· In

miz

rum

gent

ver

aspi

ag.

83,

bas

ta-

ias

er

ıll

ŋ.

P.

Trachelium majus, Blew Canterbury Bells, G. 448. Trachel. minus fl. purpur, P. in paradifo. In sepibus & sylvis variat flore albo.

Trachelium minus, G. 449. P. 644.in montofis & aridis cretaceis.

Trachelium majus Belgarum, sive giganteum, G. 448. P. 643. in divers places in York Shire.

Tragopogon luteum, Yellow Goats-beard, G.735. P. 411. In pratis, variat fl. albo.

Tragopogon purpureum, G. 735. P. 412. On the Banks of Chalder near the Lady Hescoths house two miles from whaley, Ph. and in many places of Cornwall, Dr. Gunthorp.

Tragopogon minus angustifolium, G. 735. 3. A mile on this side Epping in the Forest.

Tragopyrum, Buck wheat or Buck, G. 89. P. 1141. feritur in variis Angliæ locis.

Tribulus aquat.minor muscatellæ floribus, Smal. Frogs Lettuce, G. 824. Tribulus aquat. minor alter, P. 1248. passim in rivulis.

Tribulus aquat. minor quercinis floribus Water Caltrops, Tribul. aquat. minor alter, P. 1248. cum superiore.

Trichomanes mas, The Male English Mayden bair, G. 1146. P. 1051. In montosis & locis siccioribus.

Trichomanes ramosum, hac plantula quam Collegit in Cornubia hortum meum siccum auxit Dominus Heaton. Theologus Hibernicus.

K 4 Tri-

\*Trifolium album umbella filiquosa, White Tree foil with a Codded Umbel. Apparet sub yariis faciebus diversis crescendi temporibus, unde varia induere nomina facile poterit, videtur esse trifol si viridi soliacio Dom. Stonehous. Locis siccioribus satis familiare est.

\* Trifolium album repens capitulo spicato ex apicibus surrestris composito.

\* Trifol. Burgundicum, G. 1199.

Trifol. Cochleatum, Medic fodder, G. 1199.

Trifol. Cordatum, G. 1190. Medica Arabia Cam. P. 1115. In pratis.

Trifol. flore Cæruleo, three miles short of New Market from Ely.

Trifol. fragiferum, G. 1208. P. 1109.

Trifol. five Medica frutescens fl. flavo Clusinter legetes sed rarius.

\* Trifol. Glycyrrhites rad. magna, P. 1114.

Trifol. luteum Lupulinum, G. 1186. Trifol. montanum lupulinum, P. 1105.

Trifol. luteum minimum, G. 1186. P. 1111.in

pratis & pascuis.

\*Trifol. Lapatum CB. capite albidiori minula, aspero, two miles beyond Tilbury towards. Lee in Essex.

Trifol.luteum filiquâ cornutâ, G. 1191 in prais & fepibus.

G. 1185. 1186. P. 1104. cum priore.

Trifol. majus Corniculatum hirfutum, in pratis humidis & ad fepes.

\* Trifol. Pentaphylli ficie ex mari sabriniano.

Trifol

7

7

1

7

7

7

7

Trifol. pratense flore albo, P. 1110. ubique in pratis & pascuis.

Infol. pratenfe, G. 1185. Trifol. vulg. pratenfe

purpureum, P. 1110. cum superiore.

'Trifol, pratense capite sertaceo. In the Meadows seven miles on this side Oxford.

Trifol. paludosum, G. 1194. P. 1212. in locis

humidis & aquis stagnantibus.

'Trifol. palustre nodosum semine aspero. In the Meadows near Cardyff.

Trifol. montanum majus flore albo fulphureo, betwirt North-fleet and Gads hill in Kent.

\*Trifol. purpuream dilute glomerulis florum oblongis five pediculis cauli adnatis. In the Close leading from Cambridge to Cherrez-hinton Church. C.C.

Trifol. filiquosum minus, G. 1190. in pratis &

pascuis.

I

ca

7

tet

iol.

in

ilq;

irds

atis

reo.

ratis

ano.

rifoli

Trifol. supinum pumilium flosculis longis albis, in montosis & siccis.

Trifol. ftellatum glabrum, G. 1208.6. in the Salt

Marshes at Dartfort.
Tripolium vulg.majus, Great Sea Star-Wort, G.
13. P. 674. In sals & maritimis.

Tripol. vulg. minus, G. 413. Tripol minus, P.

673. fine icone cum priore.

Triscum spica mutica, White or Lammas Wheat, G. 65. bare or naked White Wheat, P. 1120.

Triticum aristis circumvallatum, Bearded or Red Wheat, G. 66. Trit. aristis munitum, P. 1120.

\* Tri-

\* Triticum spica mutica albicante granis ruses centibus Robus Columella, CB. Kenish Wheat, Red Wheat, Red Lamma, CC.

\*Triticum aristis donatum, spica albicante, grans rusescentibus, Red Wheat with a beardd

White ear. CC.

\* Triticum aristis circumvallatum, glumis hishtis spica albicante, Gray Wheat, C.C.

Triticum aristis munitum glumis suscis hirlats, Dunover. In agro Wigorniensi & Gloucestr,

\* Triticum aristis donatum granis & spica rule scentibus, Tritici hyberni primum genus ci-statum, CB. bearded Kentish Wheat, CC.

\* Triticum aristis & spica & granis rusescentibus, glumis præ Cæteris aristatis levibus & splendentibus, Red-ear'd bearded Wheat, C. C.

\*Triticum Conica figura hirsutum cum & fine aristis, unde dicitur Cone wheat in agro Oxoniensi.

Tubera terra majora fimbriata calici formia,

From Lancashire.

Turritis, Towers Mustard, G. 272. Turritis vulgatior, P. 852. Below Beacon-hill in the way to Bath.

Tuffilago, Colts-foot G.SII. P. 1220. in locishu-

midis & quandoq; inter segetes.

Typha, Cats-tail, G.46. Typha minor & minima, Reed Mace, P. in stagnis.

## V.

V Actinia nigra, Black worts, or Wortle berries, G. 1415. Vaccin, vulg. P. 1456. in montolis & Ericetis.

Vaccinia nubis, G. 1420, abundantly on Pendle bill in Lancashire, both in and out of Berrey

in June.

fu-

is,

dr.

OS,

en-

&

gro

nia,

ritis

the

hu-

ma,

Vac-

Vaccinia palustria, G. 1419.P.1229. On a bog two miles beyond Lyn, and on another bog sour miles beyond Guilford, and in many places of the North.

Vaccinia rubra, G. 1415. Vaccin. rubra foliis buxeis, P. 1456. In the wild Moors of Nor-

thumberland.

\*Vaccinia rubra foliis myrtinis crispis. Four miles from Heptenstall near Widdop on a great Stone by the River Gorlpe in Lanca-soire.

Valeriana major sylv. Great wild Valerian, G. 1075. ad rivulorum & fossarum ripas.

Valeriana minor, G. 1075. Valeriana fylv. P.

123. in pratis humidis & fylvis.

Valeriana Graca, G. 1076. P. 123. On the Rocks betwixt Man mater Tarn and Mananco, where the highest Rock standeth round like a Castle.

Valeriana petræa, P. 123. Valeriana annua Clus. G. 1077. inter segetes.

Verbascum nigrum, Base Black Mullein , G.

775.

775. Verbascum nigrum vulg P. 61 in montosis & cretaceis.

Vermicularis frutex, Shrubby Sengreen, G.513.
Vermicularis arborescens & fruticosa altera,
P. 731. in maritimis.

Verbena communis, Common Vervain, G. 718. Verbena vulg. P. 675. juxta semitas.

Veronica vera & major, Fluellen or Speedwell, G. 626. Veron. mas vulg. supina, P. 550. ubique in pratis.

Veronica mas recta, G. 626: Veron. erecta, P. 550. On St. Vincents Rocks by Bristol, Mr.

Goodyer.

\* Veron. assurgens sive spicata, G. 628. Veron. recta angustifolia, P. 550. in several Closes near New Market-heath. C.C.

Veronica recta minima, G. 627. Veron pratensis minor, P. 551. fine icone, At Barns near

London.

Viburnum sive lantana, Way faring-tree, G.1490
The pliant Mealy-tree, P. 1448. in sep-

Vicia, Tare, Fetch, or Vetch, G. 1227. Vicia vulg. fativa, P. 1072. ubique feritur, variat fl. albo.

\* Vicia siliquis glabris flore dilutè cæruleo Tim

tare, CC.

\* Vicia sylv. flore albo, P. 1072. G. 1227. Near

Oxford, Mr. Bobart Junior.

\* Vicia repens fl. rubro siliquis longis, folissque brevibus, In a Moor between Sunning and Maidenbead.

\* Vi.ia

· Vicia foli gramineo siliqua porrectissima, Dr. Dale, about Tyburn and Maribone Park.

Vicia Bathoniensis, vel maxima sylvatica, In Smoak hall Wood by the Bath, and at the Devizes, Wiltsbire, Mr. Goodyer.

Viola canina fylv. Doggs, or wild Violet, G. 851.

in fepibus ubivis.

Viola canina fl. albo, In Hamt fed Wood on that fide the Cheffnut walk, where the two waves meet.

Viola nigra five purpurea, G. 850. in sylvis &

umbrofis.

fis

3.

a,

8.

1

0.

P,

n,

cs

fis

ar

90

pi-

cia

at

n!-

ar

ne

nd

ia

Viola flore albo, G. 850. cum superiore.

Viola martia alba odoratissima, From Cornwal, Dr. Ganthorp.

Viola martia lutea, G. 851. On the hills near

Latham in Lancashire.

Viola fol. Trachelii serotina hirsuta radice lignofa. In Chariton Wood, and in the Lane leading to Sittingbourn, and in the way to Limsham in a great Gravel pit.

Viola Tricolor Panfyes, or Hearts-eafe, G. 854. in areonofis & flerilibus & inter fegetes.

Viola palustris stellata, in a standing pit a fourth' part of a mile from Bigles worth, in the way to St. Neots in Bedfordshire.

Viorna, Travellers joy, G. 886. Clematis sylv.

latifolia, P. 383. ubique in sepibus.

Virga aurea, Golden Rod, G. 430. Virga aurea vulg. P.542. in ficcioribus, & ericetis, Gicon. secunda ponenda loco primo.

Pifcum, Mifelto, G. 1350. P.1393. Nascitur in

Quercu,

Quercu, Pomis, Pyris, Tilia, Acere, Corylo, Rubo, Salice, Fraxino.

G. 1043. Ulmar. vulg. Common Meadon [weet, P. 592. in pratis humidioribus.

Ulmus folio glabro, Witch Elm, or smooth leav'd Elm, G. 1481. P. 1404. in sylvis & sepibus.

Ulmus folio latissimo scabro, G. 1481. Ulmus latiore folio, P. 1404.

Ulmus minor folio angusto scabro, G. 1478, ulmus minor, P. 1404. cum superioribus.

Ulmus vulgatisfimus folio lato scabro, G. 1478, Ulmus. vulg.cum famarris sive seminibus suis, P. 1404.

Umbilicus veneris, Wall Penny wort, G. 528. Umbil. veneris vulg. P. 740. In ruderibus & muris, flos quandoque purpurascit.

Urtica urens, Common stinging Nettle, G.706. Urtica media sylv. P. 441. in compitis & lous in cultis.

Vrtica minor, G. 706. P. 441. cum priore.

Uftilagines, Burnt Corn, Avena, Hordei, Secalis, Tritici.

"Vve marinæ similis planta, A mile from Wine chel sea Castle towards Hastings on the Sea shore.

## Z.

Zrie, Stinking Gladden, G. 60. Zyris sive spatula sænda Gladwin, P. 256. In sylvis humidis.



# De Plantarum Classibus.

MAximi fane est momenti res omnes natu-VI rales ad certas classes reducere. Sic enim halius ediscuntur, promptius edocentur, fideliufqueretinentur. Præterea, hanc viam ingreffus non committet ut sub variis nominibus eandem plantam tradat, quod à quibusdam fieri non diffideest observare, neq; ipsi opus erit in sua Catalogo lectorem fæpe remittere. Antiquiores planmum historiam confuse ad modum tradiderunt. Primus, qui inordinem reduxit erat Caspar Bauhinus peritistimus botanographus in admirando ho quinquaginta annorum Pinace. Quem fequati alii Joan Bauhinus Jonftonus & Spigelius. Nos ex Omnibus hisce methodum inchoavimus que mihi magno usui fuit, speroq; & aliis suturam. Quam in proxima editione Deo vitam largente observare decrevi. Hoe unum monenlus & lector me eandem fub duobus & quandoq; tribus loculis reposuisse, sub propriis Romano, subalienis Italico Charactere quod ea ratione lei ut inde scias sufficere ut ad plantæ ignotæ rognitionem pervenias si sub duabus tribusve Classibus inveneris, Ex.gr.si tibi offeratur planta

1

e 1)

aquis innatans rotundifolia dis classibus inspecting nymphæam esse certo pronuntiabis. Par & ratio de cæteris. Et si quis diligenter perpenderit de methodo singularum specierum tam summana quam subordinatarum definitiones tradere pot tuerit, easque in tabulas redigere ut hac in ante tyro suo marte brevi peritus evadat botanos gus. Hoc ipse tentavi & aliquousq; processi ne du bito quin brevi persecturussim. Quod iden in animalibus, sossilibusque aggressus sum specimentandem ad umbilicum perductum iri Intera in Animalibus tradendis Aldrovandi & Jonston vestigia premo in sequentibus.

Grulatus Alopecuroidea, Anthomethum Aquat. Arundinac. Avenac. Briza spica. Bromoides Bulbosa Canina. Caryophyllata, Cristat. Cyperoidea. Dactyloid. Echinat. Festua, Geniculata, Glumis variis. Hirsuta, Hordesca, Juncea, Junci, Loliacea, Marina, Montan. Milia. Murorum Palustre Nemeros, Nodos, Pannitusa. Phalaroid. Plumosum Piperinum Pratens. Section. Segetal. Sorghin. Stellat. Spartea, Spicat Strat. Sylvatica, Toment. Tremul. Triticea, Typhoid. Ustilagines.

Erealia Habent geniculatum culmen & fol-Harundinaceis constant semenq; panisci aut pultibus aptum, in spicis aut Jubis serum, funtq; cavicaulia, quarum partes. Culmus, Vagina, Spica in qua arista. Aruncus, Frit. Granom, Glum, đ

ntio hat

M

polite

IF.

cm

101

rez

m

THE PERSON NAMED IN

ica,

cea,

lat:

d

is in

oid

fol

ici

nt

agt.

AW,

mi,

Gluma, funtque Avena, Ægilops, Bromos, Hordeum, & fpurium & gr. -Hordea-cum, Triticea, Secale, & gr. Secaliaum, Lolium, Phalaris, Tragopyrum, Melampyrum,

Eptomacris foliis & flore pulchro. Hæ præter Typham & Gladiolum, bulbofam habent radicem Afphodelus, Colchicum, Crocus, Cynoforch, Gladiolus, Hyacinth. Iris. Nidus avis, Ornithogalum, Orobanche, Orchis, Palma Christi, Serap. Tragorchis, Tragopogon, Triordis, Typha, Zyris.

A Llium spirant Allium, & Ursi-

Oliquose. Hæ acres & mordaces pleræq, flores Dquadripetali

Alliaria, Barbaræa, Brassia, Bunias, Cardamine, Chamænerion, Camælia, Chelidonium majus, Eruca, Erysemum, Gemelampsana, Lepidium, Leucoium, Lysimadia quædam, Nastur. Oxys. Pecten Veneris, Persoliata, Pilosella siliquos. Raphanus, Rapidium, Rapum, Sinapi, Sium aquat. Sophia Chirurg. Turriris, Trisolia quædam.

Capfulata. His vascula rotunda pro seminis theca, seu bursula flores quadripetali busa pastoris, Cochlearia, Crista-Galli, Draba, beris, Lupulous, Myagrum, Nasturtium petræum, Pedicularis, Thlaspi.

I.

Eguminosa Anthyllis, Aphaca, A. Arthyllis, Aphaca, A. Arthyllis, Catanance, Cracca, Faba, Ferrum Equinum. Glaux, Lathyrus, Lens, Ochrus, Q. nobrychis, Ornithopodium, Pisum, Terra gland. Vicia.

P Accifera. Baccæ funt minores fructus, nec at modum conferti, molliori cortice praditi, Androsæmum, Asparagus, Bryon, Clymen. Dulcamara, Hedera, Herba Paris, Polygonat. Solanum.

Ampaniformes flores. Quorum flosculi instar Campanæ rotundi continui in mucronen, vel turbinatum conum desinunt. Canpanula, Convolv. Gentianella, Helleborastr. Hy. acinth. Hyoscyam. Lilium conval. Rapunt. Soldanella, Trachel. Umbilicus Veneris. His anneste quibus flos indivisus Antirrhinum, Aristoloch, Digitalis.

Hodostora. Quibus flos patulus quinq; constans foliis rosarum instar. Adonis Alcara, Althæa, Anemone, Argentina, Agremone Blattaria, Linaria, Linum, Lysimach. Malva, Nummularia, Pæon, Papaver, Pulsatil. Ranunculus, Verbascum. His Junge Ascyrum, Hypperic.

Corymbifera. Corymbi sunt extremitates la tiores storum. His slos densus &in capitulus aggestus, multis staminibus minutissimis stipatum.

Absynth. Arthemisia, Baccharis,

theris, Bellis, Chamæmel. Conyza, Cotula. Crithmum. Chryfanth. Doronicum, Gnaphal. Helen. Jacobæa, Millefol. Parthen. Pilofella, Parmica, Tanacet, Tripol.

mu

0.

TZ

出山

ly-

0

tar

ún.

ly-

ol-

de

ch.

108

00

on.

מתנו

115

Judg

slay

hig

tum

Bac-

aris,

Aliculata, quarum flores calici insistunt, & Charum plerumq; pedamentum longius est.

Armer. Armorar. Ballote, Behen. Caryophylli.

Cinvolv. Centaur. minus. Tentianella. Gratiola:

Melistophylion. Fuchs. Muscipula, Paralys. Pneumonanthe. Primulæ, Pseudomel. Saponaria, Soluella, Speculum veneris.

TErticillata. Harum flores instar verticilli V caulem ad alas ambiunt, interspersis foliolis, diinsuper funt galeati, viz qui galeam referunt mhantque ex supremo petalo galeam referen. duobus lateralibus velut labiis & quarto menmalfimilando galezq, opposito. Hæc quatuor forem construunt, omnesq; fortem spirant odo-Acinus, Alfine Hederac Ballote, Berem. m Bugul, Calaminth, Cannab, Chamadr, Chamipitys. Glinopod. Gratagog. Cufcuta Euphrag. Gleopis Hormin, Lamium, Marrub. Melam-Mr. Meliff. Menthæ, Mentastrum, Origan. Panax colon Prunella, Puleg. Scord. Scoridon, Scropular. Serpyll. Sideritis, Stachys, Urtica.

Mbellifera. Sic dicta ab umbella mulierum quam referunt quoad latitudinem& rotunmarem harum pleræq; tenuifol, feminaq; appamelingula cum bina fuerint nudaq;

L 2. Aneth.

Aneth. Angl. Anisum, Apium, Bulbocast. Benglarum, Carum, Caucalis, Ceresol Cicuta. Cicutar. Coriandr. Delphin. Fumar. Filipend. Herba-Ger. Hipposelinum, Meum, Millesol. Myrrhis, Pastina ca, Peucedanum, Percepierre, Petroselinum, Maced. Pimpin. Radix cava. Scandix. Selin. Seseliprat. Sium, Sphondl. Valerian. Ulmar,

S Picata, quarum flores à latiore basi in mucronem desinunt. Agrimonia, Bistoria, Blattaria, Blita, Cauda mur. Circæa. Cornu cervi. Coronopus, Holost Luteola, Lysimach, purpurea, Mentha, Persicaria, Plantago, Potamogetta, Resed. Serpentur. Sesamoides, Tapsus, Tista, Verbasc. Verben. Veron. Umbilis. Veneru.

Pappola. Hæ post flores lanosos, villossates quæ vento defluunt acquirunt, plerisq, flos luteus, & semini plurimi villi adhærent Baccharis- Monspel. Cardum. Crithmum. Den Leonis, Helennium, Hieracia, Hypocharis, Hyoseris, Lactuca, Pilosell. Senecio, Sonda, Tripol. Virga aur.

Ripetala, quorum flores tribus foliis confitant Hoc sensu utitur Petalon peritifiimas Botanographus Fab. Columna Galum, Militaris, Morsus Rana, Plantago aquat. Rubeola arvens, Sagitta, Stellaria.

Cardina, Card. vulg Globofus. Eriocephalus U P

M Sol Pri

0.

i,

1000

3

S TO THE STATE OF THE STATE OF

cohalus Mariæ, Stellatus, Solstitialis, Cirsium Dolacus, Eryngium, Virga paftoris.

Chinato capite. His caput minus ex femm. Casperiusculis constans. Agrimonia, Aparine, Caryophyllata, Canclais Circaa, Corom, Cynogloff. Diplacus, Echium, Medica, Myositis, Plantago aquat. Ranunc. Sagitt. sorpoides, Sanicula, Sparganium, Symphyt. Tribulus-terreft. Urtica.

Apite Squammoso. Capitatæ quæ capitibus Umajoribus craffioribufq; funt præditæ rotunlisvel ad rotunditatem accedentibus Bucharis, Cardui. Cyanus, Eupator aq Jacea, lacob. Lactuca. Morfus diab. Prunella, Scabiofa, Senecio Serrat. Succif.

Raffa & Succulenta, à foll. constitutione appellantur, quæ carnola pinguia, fucco nigentia quæquæ radice ablata diu vivunt, per-· Acetofa, Atriplex, durantq, fucculenta Beta Blitum, Cepea, Cochlearia, Cotyledon, Craffula, Crithmum, Fucus, Halimus, Kali, Paronychia fol. rutaceo, Pinguicula, Portulaca, Abodia radix. Saxifraga alba & aurea, Sedum, Spongia, Teleph.

Convolvuli & Scandentes . Hæ non nisi alterius U fulcro & ope fustinere se commode posfant, Alfine baccifera, Bryonia alha & nigra, Clematis, Convolvuli, Cuscuta, Dul-EL RET

L 3

Dulcamara, Hedera, Helxine, Lupulus, Pericly. menum, Polygonum, Polygonatum, Viorna, Volubilis, Vinca pervinca,

Allescentes que lacteum fuccum fundent Cichoreum, Dens Leonis, Eful Lactuca, Peplus, Rapunculus, Sonchus, Tithyma. lus, Tragopogon, folum Chelidon majus emitit fuccum luceum.

Tomiles & humi ferpentes. Alfine, Anagal Anthyll.mar. Canda murit Platine, Euphrag. Gentianell. Glanx, Gratisla. Kali, knamel. Lunar. Musens, Millegrana, Nummularia minor. Ophiogloff. Ornithoped. Paronych. Percepierre, Pneumonanthe, Polygala, Polygon Rubia, Ruta muraria, Rubia Cynan b Saxifraga, Seda minora, Serpyllum, Spergul Tribuluster. reftr. Veronica minor, Vermicularis.

M Olles & Und nofe. His folia mollia Unduofa,& glabra. Acetofa, Althan Atriplex, Beta, Blitum, Bonus Henricus, Cyno crambe Glastum, Lapatha, Malva, Mercurialis, Parietar, Portulac.

Apillares. His radices plerumque numerola capillaceæ fol. sæpe tenue & multiplex, & sne Aore femineve manifelto quod foliorum inferiori parti inftar pulvisculi adhæret. anth. A fplen. A dianth aur. Capill. Ven. Dentaria, Dryopt, Filix, Filicula, Hepatica.

ur. Lonchitis, Lunar. Muscus, Onopteris, Opuntia mar. Osmunda, Physlitis, Polyped. Rorella, Ruta murar. Trichoman.

Erto fol. numero constant. Monophyll-Ophiogloss. Bifol. Trifolia sunt Fragaria.

Lagopus.

¥e.

0

131

11

ia-

tit

M

na)

05,

Sil A

n-h.

1

4

M)

0

64

9

la ne

XI.

h-

a,

ŕ.

Medicæ, Meliotus, Oxys. His accedunt quæ foliæ singula tripartita habent. Aquileg. Crithmum Ruta, Thalictrum, Peucedanum, Paronych. fol. Rutac. Quadrifol. Iotus, Herba Paris. Pentaphyll. Tormentill.

A Sperifol. Asperum lanuginosum vel hirsu-Atum fol.habent. Anchusa, Bugloss. Cardus. Cichor sylv. Cynogloss. Echium, Hieracia, medam. Lithosperm. Lycopsis, Pilosella, Symphyt.

N Ervifol. Quibus infignes conspectuiq, evidentes nervi insunt, vel per longitudinem pralleli, vel liratim exarati.

Allium Ursinum, Bifol. Calceolus Mariæ Colchic. Gentumil. Lilium conval. Monophyll. Plantago. Polygonat. Saponar.

OTellate. Quarum fol. caulem vel instar stellæ Ovel, crucis forma ambiunt. Aparine, Asparagus, Asperula, Cruciat. Equiset. Galium, Hirba Paris. Lysimashia lut. Mollugo. Pyrola Alsnes fl. Rubia. Spergula.

Rotun-

Rorella, Ranunculus, Sanicula, Saxifrag. alb, & aurea. Soldanella, Tragopyrum, Tuhjmalm Helloscopius, Tuffilago Umbilic, ven.

foliis Allium & Ursin Bellis minor. Caltha palust. Caryophyll. mar. Carla muris. Chelidon. min. Colchicum, Crocus, Gladiolus, Hyacinth. Holostea, Limon. Oxys, Paralf. Petast. Plantago, Primula, Rorella, Saguin, Tussilago, Typha, & inter gramina Tomento. sum, Hirsutum nemoros, Marinum spicat. Junce um Marit. Palustre Cyperoid.

A Quis innatant. Conferya, Coyledon, Gram, fluviat. Lens palustris, Millefel. Militaris, Morsus Ranæ, Nymphææ, Potamogelton, Sagitta. Stellaria, Tribuli. Viola palustra

Alsine, Anagal Angel. Apium, Arundo Asphodelus, Calaminib. Cardamine, Eruca, Eupator, Gladiol Hyerola, Lapath. Lysimach. Marrubium, Mentha, Menthastr. Nasturt Ocnanthe, Panax Colon, Platago, Ranunc. Raphanus, Saxifraga, Sium, Spar

ad ur

Ma

t,

m.

4

la. &

lius

S.

Dr Zis

da di

k.

e-

e-

į.

3

b.

-90BB1

surganium, Symphyt. Typha, Vaccin. Valeriana.

Arina. Abies, Alga Asparagus, Cochlearia, Corallina, Corallium, Crithnum, Eryngium, Fucus, Glaux. Halimus, Kali, Imon, Opuntia, Quercus, Soldanella, Spongia, Iripolium, Vermicularis Frutex.

M Arina aliæ.

Alopecuros, Alfine, Althaa, Arthemifia, Brassica, Caryophyllus, Cyperus, Echium, Eruca, Filix, Gnaphaliumi, Gramina, Holost. Jacob. Lyhnis, Medica, Malva, Musci, Papaver, Pisum, Sedum, Spergula, Tithymalus.

L'Acrementa terre ex putridis lignis vel herbis orta quæ male dicuntur semine & radice destitui. Fungi, Musci, Tubera.

Arbores & Frutices.

D'Accifera. Alnus, Agrifol. Arbutus, Aria D'Theoph. Berberis, Buxus, Cornus erica bacif. Ebulus, Euonymus, Frangula, Junip. Laureola, Ligustr. Morus, Myrcus, Ornus, Oxyacanthus, Periclymen. Rhamnus, Ribes, Rubus, Ruscus, Sambucus, Soibus, Taxus, Vaccinia, Wilcum, Viburnum.

Onifera. Abies. Larix. Pinus, Pinaster.

Landifera. Castan, Fagus, Quercus.

J Ulifera. Acer, Alnus, Betulus. Carpinus, Castanaa, Populus, Salix & inter Plantas A. corus.

Malus fylv. Pyrus, Rofa.

Os Sei inten Frudus, Cerafus, Cornus, Pru-

Noifera. Corylus, Juglans, Nux vesicar.

R Hodoflora. Cystus, Helianth. Rosa, Ro. rismarinum. Rubus.

Siliquofe. Anonis, Genista, Genistell. Gly-

S Pinofe. Agrifol. Anonis, Berberis, Genifold Juniper. Prunus sylv. Oxyacanthus, Rhannus, Rosa, Rubus.

E Chinophora. Castan. Fagus, Myrtus Brand.

R Otundifolia. Buxus, Bruscus, Morus, Popul. Salix Caprea, Tilia, Ulmus.

Enuifolia. Erica, Juniperus Taxus vide



Rogatu Nobilissimi Viri præmisi nomina Anglicana Latinis in fequenti Alphabeto.

ř.

4.

IIA.

nA TA

Alc

tu

A Dders tongue, Ophiogloffum. Adonis flour. flos Adonis. Agramonie. Agrimonia. Alder tree. Alnus. Algood, or common Mercury. Bonus Henricus. n A Allheal. Panax. IIA. Allseed. Millegrana. 24 Anemone. Anemone. Angelica, Angelica. Anise. Anisum. Apple. Malus. 1 Archangel. Lamium. MA Arrowhead. Sagittaria. de Atmart. Perficaria. Asphodil. Asbhodelus. Alparagus. Asparagus.

Ash tree. Fraxinus.
Ashweed. Herba Gerardi.
Avens. Caryophyllata.

B

Arbery bush. Berberia. DBarley. Hordeum. Stone Balil. Acinus. Wild Basil. Clinopodium. Balm. Meliffa. Bean. Faba. Beet. Beta. Beech tree. Fagus. Bell flour, Campanula. Betony. Betonica. Bindweed. Convolvulus. Sea Bindweed. Soldanella. Birch tree, Betula. Birds neft. Orchis abortiva. Birds foot. Ornithopodium. Birthwort, Aristolochia. Bitter-sweet. Dulcamara. Blew.botle. Cyanus. Blites. Blitum. Box tree. Buxus. Bramble bush. Rabus fylv. Brook lime. Anagallis aquat. Broom. Genista. Broom Rape. Orobanche. Bryony. Bryonia. Buck wheat. Tragopyrum.

Bucks-

Chefra

Botte

Buckshorn plantain. Serpentaria.
Buckshorn. Cornu Cervinum.
Buckhorn. Rhamnus folutivus.
Bugl. Bugula.
Bugloff. Bugloffum,
Vipers Bugloff. Echium.
Wall Bugloff. Lycopfis.
Burnet. Pimpinella.
Burdock. Bardana.
Butchers Broom. Rufcus.
Butter Bur. Petasitis.
Butterwort. Pinguicula.

Buc

Buc

Bu

Bu

61

V

Bur

Bu

180

Bo

C

0

0

0

C

Abbage. Brassica. Calamint. Calamintha. Caltrops. Tribulus aquaticus. Cammomile. Chamameium. Canary feed. Phalaris. Carrot. Pastianca tenuifolia: Caraways. Carum. Cats-tail. Typha. Catch-fly. Muscipula. Celandine. Chelidonium, Centaury. Centaurium. Charlock. Rapistrum. Cherry. Cerafus. Chervil. Cerefolium. Chesnut. Castanea. Chervil Myrrhis. Chick weed. Alfine.

Cinque -.

Dill.

Dittand

Divels

430C

Dodd

Drop

Water

Dac

Die

Oak

B histre

Illax.

T Toad

Dill.

Sweet Corn

Hebar

Cinquefoil. Pentaphyllum. Clary. Horminum. Cleaver. Aparine. Cocl. Pseudomelanthium. Colts-foot, Tuffilago. Coralline. Corallina. Corall, Corallium. Coriander, Coriandrum. Corans. Ribes. Cornel or Dogg-tree. Cornus. Columbine. Aquilegia. Saracens Confound. Solidago Saracen: Cow-wheat. Crataogonum. Cowslips. Paralysis. Cranes-bill. Geranium. Cress. Nafturtium. Dock creis. Lampfana. Swines crefs. Coronopus Ruellii. Winter cress. Barbarea. Crocus, Crocus. Crow-foot. Ranunculus. Cross-wort. Cruciata. Cud-weed. Gnaphalium. Cumfry. Symphytum. Cyperus. Cyperus.

D

Andelyon. Dens Leonis.
Dane-wort. Ebulus.
Darnel. Iolium.
Daysie. Bellis.

Dill. Anethum.

d

Ò

D

Œ

0

(

Dittander. Lepidium.

Divels bit. Morfus Diaboli.

Dock. Lapathum.

Dodder. Cufcuta.

Drop-wort. Filipendula.

Water Dropwort. Oenanthe

Ducks meat. Lens palustris.

Dyers weed. Luteola.

E

Elicampe. Enula campana.

Elder. Sambucus.

Elm-tree. Ulmus.

Eyebright. Euphragia.

I

[Everfew. Matricaria.

I sow Fennel. Peucedanum.

Star-like water Fennel. Stellaria aqu.

191 fern. Filix.

10 Oak Fern. Dryopteris.

fig wort. Scrophularia.

lir-tree. Abies.

Sweet Flag. Acorus.

(Orn or Sword Flag. Gladioluse

Ilax. Linum.

11.

Toad-Hax. Linaria.

Heabane. Congza.

Flour de luce. Iris.
Fluellin. Elatine.
Tox glove. Digitalis.
Frogs bit. Morfus Rana.
Tumitory. Fumaria.
Fuss balls. Tubera terra.
Furz bush. Genista spinosa.

G

Arlick, Allium.

Gentian Dwarf. Gentianella.
Germander. Chamadrys.
Water Germander. Scordium.
Gladdon stinking. Zyris.
Glass wort. Kali.
Goats beard. Tragopogon.
Gold of pleasure. Myagrum.
Golden rod. Virga aurea.
Golden locks. Adianthum aureum.
Grass. Gramen.
Cocks-foot Grass. Ischamum.
Gromel Lithospermum.
Bastard Gromel Anchosa degener.
Groundsel. Erigerum.
Ground Pine. Chamapitys.

H

Hairy-weed. Conferva. Hares foot. Lagoque.

Harebell. Hyacinthus: Harts tongue. Lingua Cervina. Hawthorn. Oxyacanthus. Hawk-weed. Hieracium: Heath, Erica. Helm or Matweed. Spartum. Hemlock. Cicuta. Hemp. Cannabis. Water hemp. Eupatorium. Hellebor. Helleborns. Wild Hellebor. Helleborum. Henbane. Hyofcyamus. Holly. Agrifolium. Sea Holly. Eryngium. Hollow-root, Radix cava. Honewort. Selinum Sii foliss. Hops. Lupulus. Horehound. Marrubium. Horn beam tree. Betulus. Hors tail. Equisetum. Hortles, Vaccinia. Hounds-tongue. Cynoglossum. Houseek. Sempervivum majus. Hedge Hystop. Gratiola.

1

JAck by the hedge. Alliaria.
Jews-ear. Fungus Sambucinus:
Iron-wort. Sideritis.
Juniper. Juniperus.
Ivy. Hedera.

K

Ings spear. Asphodelus.
Knapweed. Jasaa.
Knotgrass. Polygonum.
Knawel. Polygonum Selinoides.

T

Adies bed straw. Galium. Ladies slipper. Calceolus Maria. Ladies smock, Cardamine. Lady Traces. Triorchis. Ladies Mantle. Alchimilla. Langue de bæuf. Buglossum luteum. Sea Layender. Limonium. Laurel Spurge. Laureola. Larks spur. Confolida regalis. Leopards bane. Doronicum. Lettuce. Lastuca. Lentils, Lenos. Lilly. Lilium. Water Lilly. Nymphea. Line or Linden tree. Tilia. Liquorice. Glychirrhiza. Liverwort. Hepatica. Lungwort. Pulmonaria.

### M

A Adder. Rubia. Maiden hair. Trichomanes. Mallow. Malva. Vervain Mallow. Alcea. Marth Mallow. Alchea. Maple. Acer. Marsh Marigold. Caltha palustris. Corn Marigold. Chryfanthemum. May weed. Cotula. Meadow-Sweet. Tlmaria. Melilot. Melilotus. Mercury. Mercurialis. Dogs Mercury. Cynocrambe. Milk wort. Polygala. Misseltoe. Vifcus. Mint. Mentha. Money-wort. Nummularia. Moon-wort. Lunaria: Mols. Moscus. Mug-wort. Artemifia. Mother-wort, Cardiaca. Mouse-ear. Pilosella, Auricula muris. Monse-tail. Canda muris. Moth. Mullein. Blattaria. Mullein. Tapfus barbatus. Black Mullein. Verbascum nigrum. Mulberry. Morus. Mustard. Sinapi. Towers Mustard. Turritis.

er.

Treacl.

M 2

Treacl. Mustard. Thlaspi. Arabian Mustard. Draba. Mushrom. Fungus. Myrtle. Myrtus.

N

Nep or Catmint. Cataria.

Netle. Urtica.

Stinking dead netle. Galeopsis.

Night shade. Solanum.

Inchanters night-shade. Circaa Lutetianav.

Nut Hasse. Avellana sylv.

Bladder Nut. Nux Vesicaria.

7

Ak-tree. Quercus.
Sea Oak. Quercus marina.
Oats. Avena.
Wild Oats. Bromos.
One blade. Monoshyllum.
Orach. Asriplex.
Orpine. Telephium.
Origanum. Origanum.

P

Ansyes. Jacea & viola tricolor. Parsley hedge. Caucalis. Wild Parsley. Sison. purley break stone. Perce pierre Anglorum.

Persnip. Pastinaca.

Water Parfnip. Sium.

Cow Parsnip. Sphondilium.

pique flour. Pulfatilla.

Peas. Pisum.

Wood or heath Peas. Astragalus.

Everlasting Peas. Lathyrus.

Pear. Pyrus.

Pennyroyal. Pulegium.

Penny or Navel wort. Cotyledon.

Wall Pennywort. Umbilions Veneris.

Perwinkles. Clematis Daphnoides.

Pellitory. Parietaria.

Pronie. Paonia.

Pine tree. Pinus.

Pile wort. Chelidonium minus.

Pinks. Caryophyllus.

Pimpernel. Anagallis.

Plantain. Plantago.

Buck horn Plaintain. Cornu Cervinum.

Plowmans spikenard. Baccharis Monst.

Plum. Prunus.

Polypodie. Polypodium.

Pondweed. Potamogeiton.

Poplar. Populus.

Poppy. Papaver.

Spartling Poppy. Beben. Bastard wild Poppy. Argemone. Potatos - Battatas

Prick Timber, or Spindle tree, Euonymus. Theoph. Primrose. Primula veris.

Privet. Ligustrum.

M 3

Pur-

Purslain. Portulaca. Sea Purslain. Halimus. Water Purslain. Alsine fol. subrotundis.

Q

Olicken tree. Fraxinus bubula.

R

R Adish. Raphanus. Ragwort. Facobaa. Rampions. Rapuntium Rapunculus. Ramfons. 'Allium ur finum. Raspis. Rubus Idens. Ratl, or Coxcombe. Crista galli. Red ratl. Louswort. Pedicularis. Reed. Arundo. Bur-reed. Sparganium. Reed grass. Calamo groftis. Rest Harrow. Anonis. Rocket. Eruca. Italian Rocket. Reseda. Rose. Rosa. Dogg-rose. Cynorrhodon. Rose-root. Rhodia radix. Rosemary, Rosmarinum. Wall Rue. Adianthum album. Rhubarb. Rhabarbarum. Meadow Rhubarb. Thalist rum.

Rush. Juncus. Ryc. Secale.

S

Maffron, Crocus. Meadow Saffron. Colchicum. Sige. Salvia. 9. Johns-wort. Hypericum. Sampire. Crithmum. Sanicle. Sanicula. Sergrium. Satyrion. aw-wort. Serratula. wifrage. Saxifragia. Sabious. Scabiofa. Sorpion grafs. Scorpoides myofitis. Survy grafs. Cochlearia Herba Britan. Self-heal. Prunella. Service tree. Sorbus. Wild Service. Aria Theoph. Sheepheards purse. Bursa Pastoris. Sheepheards needle. Petten Veneris. Silver weed. Argentina. Smallage. Apium palustre. Snails or Medicks. Medica. Snap dragon. Antirrhinum. Sneese-wort. Ptarmica. Solomons feal. Polygonatum. Soap wort. Saponaria. Fresh water souldier. Militaris. Sorrel. Acetofa. Wood Sorrel. Oxys.

Speed-

M4

Speedwel. Veronica.

Spignel. Meum.

Spleen wort. Asplenium, lonchitis.

Spurge. Tithymalus.

Spurrey. Spergula.

Star of Bethlehem. Ornithogalium.

Water Star wort. Stellaria aquatica.

Sea Star wort. Tripolium.

Stichwort. Holosteum.

Stone corp. Vermicularis.

Strawberry. Fragaria.

Succory. Cichereum.

Swines succory. Hypocharis.

Sundew. Rosa solis.

Sunflour. Chamacistus Angl.

T

Tare. Vicia, vel Arachus.
Milk Tare. Glaux.
Teasle. Dipsacus.
Thistle. Carduus.
Carline Thistle. Cartina.
Cotton Thistle. Acanthium.
Sow Thistle. Sonchus.
Single headed Thistle. Cirsium.
Thorow wax. Perfoliata.
Thrist. Caryophyllus marinus.
Throat wort. Trachelium.
Tooth-wort. Dentaria.

Tormentill. Tormentilla.
Travellours joy. Viorna.
Trefoil. Trifolium.
True love and one berry. Herba Paris.
Turnep. Rapum.
Tutlan. Androfamum.
Tway-blade. Bifolium.
Wild Tyme. Serpyllum.

#### V

VAlerian. Valeriana.
Verch or Fetch. Vicia.
Kidney Vetch. Anthyllis.
Horf-shooe Vetch. Ferrum equinum.
Crimson grass Vetch. Catanance.
Yellow wild Vetch. Aphaca.
Violet. Viola.
Corn Violet. Speculum Veneris.
Calathian Violet. Pneumonanthe.

#### W

Wall-flour. Leucoium. Wallnut-tree. Nux Juglans. Wall pepper. Illecebra. Way faring tree. Viburnum. Wheat. Triticum. Cow-wheat. Melampyrum. Whitlow grafs, Paronychia.

Petty Whin. Genista aculeata.
Willow. Salix.
Willow weed. Lysimachia.
Wild Williams. Armoraria.
Winter green. Pyrola.
Woad. Glastum.
Great bastard Woad. Sesamoides Salamanticum.
Wood rough. Asperula.
Wood wax. Genistella.
Woodbind. Periclymenum.
Wormwood. Absynthium.
Sea Wrack. Alga.

Y

YArrow. Millefolium.

Synopsis omnium Classium quibus usi sunt Scriptores Botanici in suis de plantis Historiis, præter Alphabeticam.

Cidæ, Acres, Alexipharmicæ, Allium redolentes, Amara, Aquatica, Aspera, Bulbo-& fub his Carnofæ & Esculentæ, Campanulæ, Capfulatæ, Capitatæ, Capillares, Cardui & spinofa, Cathartica, Cichoracea & refrigerantes, Convolvuli & scandentes, Coronaria, Corymbifera Craffifolia & fucculenta, Echinophora, Epigeiophillæ, Exoticæ, Excrementa terræ, Foliis magnis, Flore placentes, Frumenta, & horum vitia, Galeatæ, Geniculatæ, Gramina, Hortenses, Humiles & serpentes, Indica, Junci, Lactescentes, Legumina, Marina, Miscellanea, Molles & Uncuosa, Narcotica & Venenata, Nervifoliæ, Nudicaules, Odoratæ, Olera seu Oleracæ vel Edules Dod. Orchis, Palustres, Pappifloræ, Rariores, Rotundifoliæ, Saxifragæ, Siliquofa, Stellata, Tenuifolia, Trifolia, Verticillatæ, Umbelliferæ, Vulnerariæ.

Arbores sunt Aromaticæ, Bacciferæ, Balsamiseræ, Coniferæ, Exoticæ, Glandiferæ, Indicæ, Lachrymiseræ, Leguminosæ, Miscellaneæ, Nuciferæ,
quibus ossei fructus intus. R hodostoræ, Scopariæ
Spinosæ. Trisoliæ quæ insuper leguminosæ, Vi-

rentes perpetuò.

Hisce classibus usi sunt. CB. IB. P. G. Lob. Dod. Recchus, Jonstonus, at Hortus Eystetonsis digessit secundum anni tempora in vernas, astivas, autumnales & Hyemales.

Opertet

Oportet futurum Herbarium in primis callere hæc Technica vocabula sub phantarum partibus dissimilaribus, tanquam capitibus disposita, Generalia Characten Romano, peculiaria Italico describuntur.

R Adicis sunt, Bulbi, Tubera, Filamenta, Stolones, Turiones & Asparagus, Fundus. 010

Caulis. Ala & sinus, Genua, Genicula su articuli. Nodi, Internodia, Medulla, Hinc Ra mus Thyrsus Capreolus, clavicula & crines su viticuli. Pedunculus, pediculus seu petiolus.

Foliorum. Caro, Nervus, Crenæ, Laciniz,

Lanugo.

Florum Alablastri, Palaustium & cytinus. Calix & perianthemum, Capillamenta & filamenta Gemma & oculus. Stamina cum Apicibus sa antheris. Spica, Stylus. Unguis, Petalon. Calca. Cupulus, Fundus, Juba, Tomentum, Unbilicus, ex mulcis diversisque compactis sium Serviæ & Coronæ.

Seminis Folliculus & utriculus, Loculamer tum, Pannicula, Samera, Pappus Valvalæ.

Frumenti. Acus & palea, Arista, Alica & halica, Amylum. Cor & corculum. Culmus, Fri. Gluma, & utriculus, Uruncus vel uruncum, Locusta, Granum, Spica quæ dicitur Mutia, Merges.

Frustunm. Acini & arilli. Bacca, Carola pulpa. Dastylus, palma, sen Caryotis. Cicus de cicus.

ciccum. Nucleus, Officulum, Pericarpum, Um-

bilicus, Corculum, Volva, Racemus.

al-

ta-

am

ere

ur.

0

fen lafen

Z,

(et

ar,

m-

unt

en-

iL

B,

CZ,

a

Arborum. Scapus, Truncus, Caudex, Cortex, Liber Malicorium, Matrix & Medulla & cor. Pecten. Circuli Surculi Rami Sarmenta Ramufuli, Spadix, Vimen. Virgultum & Cremium, Vitilia, Talea, Scobs. Termes, Turiones, Frondes, Cachryas & nucamentum & julus & Catulus, Comæ, Arborum species frutex & suffrutex.

synopis Etymologica que lectori inservire poterit vel ad inveniendum rationem nominum genericorum plantarum prima lectione, vel certe ad firmius memorie insigendum prius inventa que insuper non inutilis ad ipsarum herbarum cognitionem. Plura si velis adi CB.IB. CC. CO. & ceteros Botanographos.

Folio. Acanth. Agrifol. Alectoroloph. Anifum, Asperula, Argentina, Arnogloss. Auricula Judæ, Aparine Clivers, ab adhærendo. Bisol. Bugloss. Bupleure Hares-ear.

Chamædrys, Chamæpitys. Cardui, Coronop... Cornu Cervin. Clematis, Cotyledon, Cruciata, Cynogloff. Cochlear. Chærefol. Cinara. Diplacus, Dens I eonis, Dryopteris. Eryfimum, Equifetum. Filix, Fumar. Gnaphal. Gladiolus.

Hepta-

Heptaphyll. Herba Ger. Astreed. Jacobra, Ragwort. Lactuca, Lagopus, Laureola, Lonchitis,
Lingua cervina Lycops, Lunaria. Meum, Millesol,
Monophyll. Morsus Ranæ, Myositis. Nummular. Ophiogloss. Oxalis, Oxys, Oxyacantha,
Oxyacanthus. Pedicularis, Perforata, Perfoliata,
Periclymenum, Pimpinella, Plantago. Polytricum,
Populus, Pentaphill. Petasitis, Portulaca,
Pilosella, Pinguicula, Pes Columb. Persicar.
Pulmonar. Pyrola. Ranunc & slammeus,
Spearwort Rorella. Sagitta. Seratula, Sparganium
Spatula, Stellaria. Vermicularis, Verbascum,
Ulmaria, Umbil. Veneris, Ungula Cabali
(Tussilago.) Trisol.

Aule Cicuta. Polygon, Viburnum.

aFLore, Alopecuros, Anagallis, Anthyllis, Alnemone, Antirrhinum Aquileg. Androsam. Bellis Betonica, Calceolus Maria, Campanula, Cauda muris, Chamaemel. Chrysanthemum, Clinopod Crocus. Cyanus. Digitalis. Erigerum. CO. Flos solis. Galeopsis. Helenium Hyacinthi, Hares bells. Jacaa Knapweed. Ischamon, Cocksfoot. Iris. Luteolia, Lamium Oenanthe, Ornithog. Orchis Melittias, &c. Stachys. Speculum Veneris. Tragopogon. Viola tricolor.

Adice Bistorta, Bunias, Bulbosa, Bulbosastan. Caryophyllata, Chelidonium minus, Pilewort. Cyperus, Cyclamen, Dentar. Fabr. Filipend. AT-

tis,

fol,

IM-

ha;

ata,

ca,

ar.

us .

m,

Aª

m.

nla,

m.

ksog.

um

Ć2-

us,

or.

d.

filipend Glans terræ. Glycyrrhiza. Morsus Diab. Nidus Avis. Orchis. Peucedanum Sulpherwort. Mypod. Palma Christi. Pastinaca, Polygonat. Radix cava. Rhodia radix. Rubia, Rubeola, Raphanus, Rapum, Rapistrum, Rapunculus-Strophular. Valeriana.

Ructu, Baccif. Cereal. Cærasus. Fragaria. Forossularia. Legumina. Nux Vesicar. Poma, mua. Pyrus. Rubus, Black berry. Vaccinia massi baccinia.

Semine vel vasculo seminali. Acer. Astragalus Beta. Bursa Pastor. Caucalis, Cataputia. Edium. Faseolus, Fegopyrum, Ferrum Equin. Edium. Faseolus, Fegopyrum, Ferrum Equin. Edium. Melampyrum. Millegrana, Monomyl. One berrey. Nigella Nigellastrum. Ophiosoft. Ornithopod. Pecten Veneria Pussatilla, Edinaca syl. Birds nest. Serpentar. Scorpoides. Typha, Tragopogon, Tribulus, Thlaspi.

thum, Angelica, Argemone, Aristoloch.
Avium, Asplenium, Asphodel. Catananche, Cataria, Cardiaca, Cirsium, Chelidonum, Conyza,
Consolida, Erica Euonimus Euphragia, Galium,
Glaux. Gratiola. Hepatica, Hieracium, Hormimm, Juglans, Lampsana, Lepidium, Lens, Liden, Medicæ, Panax. Paralysis, Paronychia,
Percepierre, Polygala, Prunella, Ptarmica, Pulmonaria, Salvia, Saxasragiæ, Selinum, Sonchus,
Sum, Sorbus, Scabiosa, Sideritis. Telephium
Tor

Tormentilla, Trachelium, Tuffilago, Valeriana.

AM Ala qualitate. Anonis, Blattaria, Blitum, Circæa Cynanchice Eryngium, Fungus, Helleborus, Orobanche, Sinapi.

ASApore. Abfynthium, Daucus.

Ab Dore Baccharis, Ballamita, Caryophylli, Caryophyllata, Coriandrum, Myrrhis Sphondylium.

AD Vratione. Adianthum.

AM Odo Crescendi. Atriplex, Bryonia, Juniperus, Salix, Serpillum, Thalictrum.

AT Actili qualitate Cornus. Linum, Urtica, Cardui, Carlina.

A Loco, Alfine, Anagalis, à Gallo fluvio, CO. Carum, Cepea, Cerafus, Colchicum, Cytifus, Ebulus, Halimus, Limonium, Nymphaa, Parietaria, Fotamogeiton, Ruscus, Typha

A Jis Juglans, Serapias.

IC Antis Barbaraa, Jacobaa.

1,

is

nj-

ca,

O.

m,

m.

ha.

I Nventoribus, Armeria, Adonis flos, Arthemisia, Asclepias, Enula, Gentiana, Berba Doria, Gerardi, Hyacinthus, Mercurius, Myrtus, Narcissus, Nymphæa, Tanacetum, Teurium.

Su pro Animalibus. Cynocrambe, Hypocharis, Hyoseris, Hyoscyamus, Sa-

ana.
Ab usu in Mechanicis Juncus, Ligustrum,
Inteola, Lychnis, Saponaria.

compendium temporis in quo florere incipiunt planta nostrates, sc singulis mensibus separatim digestis, p. principium, f.
sinem mensis, denotat. Durationem autem intelliges ex notis arithmeticis Ex.
gr. Vide mensem Martium & ibidem
Acer cui postscribitur 3 indicans Acer
sorere a Martio ad 3. inclusive (1) Maium & in mense Aprili quare Arundinem quam siquitur Character 5 innuens
hanc slorere ab Aprili ad Augustum, &
sic de cateris. Alphabetice disposui in
singulis mensibus ut facilius indagares.

Dens Leonis & Erigera quovis Mense.

Martio Acer 3 Cheledonium minus, Petasites 2. Fopulus 2. Quercus, Radix cava,
N Rapi-

Rapistrum 5. Saxifraga aurea 2. Sorbus, Taxus, Tustilago 2. Vinca pervinca 3. Viola 2. Ulmus 2.

Aprili Alnus, Armeria 2. Arundo 5. Betula,
Betulus, Bugula 2. Cardamine 2. Castanea,
Cerasus, Chelidonium majus 4. Clematis 3.
Cornu cervin 5. Coronopus 5. Cynocrambe 5.
Equisetum 5. Euonymus, I aba 2. Fagus 2. Fegopyrum 2. Fraxinus 2. Gentium 2. Geraunium 4. Hedera ter. 5. & aquat. 3. Herba Paris,
Hieracia 5. Nux Juglans, Lujula sive Oxys 3.
Malus, Ophiogloss. 2. Ornithogal. 2. Paralysis
2. Pisum, Primulæ 2. Prunus, Pulsatilla 2. Pyrus,
Rubes, Rosamrinum & in Augusto, Salix, Sam.
bucus 2. Stella aquat. 2.

M Aio Alchimilla 2. Antirrhinum, Aquilegia 3. Arachus, Aria, Aristolochia 3. Armoraria 2. Afphodelus 2. Auricula muris 2. Barbaria, Behen 4. Bifol. 2. Bistorta, Bryonia, Bugloss 4. Bursa pastoris 4. Caltha 2. Camelina 2. Campanula 4. Caryophillat. 3. Caucalis 4. Cerefolium, Chamædrys, Christophoriana 2. Cochlearia, Convolvuli 4. Cornus 2. Cratæogonon 4. Crithmum 2. Cyanus 4. Cynoforchis 2. Dantaria 2. Draba 2. Echium 4. Eleagnus 2. Erica 4. Euphrosine 4. Filipendula 3. Flos Adonis 3. Fragaria, Fumaria 4. Galeopsis 4. Genista 4. Genistella 4. Glaux marit. Gratiola 4. Helleborine 2. Holostea 2. Hyacinthus 2. Hyoseris 4. Hypochæris 4. Iris 2. Juniperus, Lamium 4 Lampfana 4. Ligustrum 2. Lillium conval. Lunaria,

.

1,

1,

3.

5.

e-

1

5,

3.

18,

n.

le.

ar.

Bu-

12.

Ce-

20-

non

an-

a 4.

5 3.

4.

lle-

15 4.

n 4.

Lu-

aria,

mria, Lychnis 4. Mercurialis 4. Millefol. 2. & terrest. 5. Millegrana 4. Monophyllon. Morus f. Myagrum 2. Myrrhis, Myrtus 2, Nummularia 3. Oenanthe 2. Ophris 2. Orchis 4. Oxyacanthus, Alma Christi 2. Papaver 4. Pecten Veneris Peicularis 2. Pentaphylla 2. Perfoliata 2. Periclynenum 2. Pilofella 2. Pæonia, Polygala 4. Polionum 4. Polygonatum, Ptarmica 4. Ranuncu-154. Rapunculi 3. Rapuntium 3. Rhamnus, Rorella 2. Rosa 4. Rubeola 4. Rubus 2. Sagit-12. Sanicula 2. f. Saxifraga alba 2. Serapias 2. Serpentar 2. Serpillum 3. Sefamoides 2. Staphydendron, Synanchice 4. Teucrium 4. Thlaspi , Tilia, Tormentilla 4. Trachelium 4. Trifo-14. Tripolium 2. Turritis 2. Vaccinia, Valerima 3. Veronica 4. Vicia.

Unio Acinos 2. Acanthium 2. Adianth, aureum 3. Agrifol. Agrimon. 2. Alliaria 2. Allium Alopecuros, Alfine 2. Anagallis 3. Anchufa Anthylis 4. Aparine 2. Apium 2. Argentina 2. Mparagus 2. Afperula 2. Aftragalus 2. Atriplex Baccharis 2. Beta 2. Blitum 3. Bonus Henric. Mbocastan 2. Bunias 3. Calamintha 3. Calceo-Mariæ, Cannabis, Cardiaca 3. Cardui 3. hamamel 3. Chamænerion 2. Chamæpytis 3. licza 4. Clinopod. 2. Conyza 3. Confolida 2. unciata 2. Cynogloss. 2. Daucus 2. Digitalis 2. Dofacus 2. Doronicum 2. Ebulus 3. Enula 2. ylimum 2. Eruca 2. Eryngium 2. Elula 2. Fer-Equinum, Galium 3. Gladiolus 3. Glastum Glaux, Gnaphalium 3. Hepatica 2. Herba Gerardi N<sub>2</sub>

Gerardi 3. Hipposelinum 2. Horminum 3. Jace 2. Iberis 2. Irio 2. Kali 3. Knawell 3. Lactuca 2. Lagopus 3. Lapatha, Oxalis 2. Lathyris 2. Lepi: dium, Lichen 4 Limon 2. Linaria 3. Linum 2. Lithospermum, Luteola 2. Lysimachia 2. Lysopsis 3. Malvæ 2. Medicæ 2. Melampyrum 2. Me. liffa 3. Meum 2. Militaris 3. Mollugo 3. Morlus Ranæ, Muscipula 3. Myosit. scorpiod. 3. Nidus avis 3. Nymphæa 3. Ochrus 2. Origanum 3. Ornithopod. 3. Orobanche, Parthenium 3. Peplus 2. Persicaria 3. Peuced. 3. Pimpinella 3. Pinguicula, Plantago 2. Portulaca, Potamogei. ton 2. Prunella 3. Pfyllium 2. Pulegium 3. Pulmonaria 3. Pyrola 2. Raphanus, Reseda 3. Salvia 3. Saponaria 2. Scrophularia 2. Scabiola 3. Scordium 2. Scorodoprasum 2. Sedum 2. Sideritis 3. Sinapi 2. Sison 2. Solanum 3. Soldanella 2. Sonchi 3. Sophia Chirurg. 3. Spartum'3. Speculum Veneris, Spergula 2. Sphondilium 2. · Stachys 2. Tithymalus 2. Tragopogon 3. Tribu lus aquat. 3. & terreftr. 2. Viburnum 2. Ulma. ria 3. Umbilicus Veneris 2. Urtica 3.

Julio Abfynth. 2. Acorus 2. Ægilops bromoid.
2. Alcæa 2. Althea 2. Androsemum 2. Angelica 2. Anonis 2. Aphaca, Aribaccæ 2. Arbotus 2. Arthemisia, Ascyrum 2. Aster 2. Bardana Blattaria 2. Brassica, Bromos 2. Bupleurum, Catanance, Centaurium 2. Chamæcistus 2. Chaysanthemum 2. Cicuta 3. Cicutaria 2. Cirsum, Cotula 2. Cymbassica 2. Cyperus 2. Dulcamara, Elaphoboscum, Eupatorium 2. Festuca 2. Galega

Galega 2. Glycyrrhiza, Halimus 2. Jacobæa 2. Juncus 3. Lathyrus 2. Lens 2. Lolium, Marub. 4. Melilotus 2. Menthæ 2. Menthastr. 2. Nasturium, Onobrychis, Parietaria, Pastinaca 2. Phabris 2. Pseudomelanthium, Rhodia radix, Selimm, Sii foll. Serratula 2. Sium, f. Solidago, Tancetum 2. Tapsus 3. Telephium 2. Thalictrum 2. Typha 2. Verbascum, Vermicularis frutex 2. Verbena 2. Viorna 5.

A Ugusto Anethum, Agremone, Avens, Cichoreum, Elatine 2. Euphrasia 2. Genmuella, Lupulus 2. Morsus Diab. Panax Coloni, Pneumonanthe 2. Sparganium, Tragoriganum, Virga aurea, Zyris.

SEptemb. Berberis, Colchicum, Laureola, Oxyacantha, Pulfatilla.

Ctob. Leucoia 3.

Icea

a 2.

epi.

0 2.

/co-

Me-

rlus

idus

n 3. Pe-

3.

gei-

Pul-

Sal-

2 3. ide-

nella

n 3. m 2.

ibuma-

An-

otus

dana rum,

heyium, ulca-

ca 2. Ilega D Ecemb. Helleborastrum, & quandoq; pri-

Anuario Anemone 4. Bellis 8. Corylus 3. Cotyled. 4. Paronychia 2.

Ebruario Buxus 2. Crocus, Narciffus 3.

N 3

Qua-

# Quadrupeda Britannica.

Solidungula, Whole-Footed.

Equis, a Horse, I. 8. Ald. 21. G. 404. Cantherius, I. ib. a Gelding. Pullus equinus a Foal. In Equilibus Nobilium, Equus Friscus, I. tab. 2,

Afinus, an As, I 23. tab 6. G. 5. Mulus, a Mule, I. 23. tab. 6. G. 702.

Bisulca ruminantia, Cloven Footed, chewing the Cud.

Taurus, a Bull, I.42. Ald. 36. G. 102. Vacca lactans, I.ib. a Milch Com, Bos, I. tab. 14.15. an Ox. Juvenca, an Heifer, Vitulus, I. tab. 15. a Calf, ex Vitulorum Ventre sæpe eximuntur pilæ ex deglutitis pilis propriis Ald. 40.

Aries, a Ramor Tup, I. 60. Ald. 397. G. 771. Agnus, I. tab. 22. a Lamb. Vervex, a

Weather, I. tab. 22. 110.

Caper, a Weather-Goat, Hircus, a He-Goat, Hzdus, a Kid, 1. 80. tab. 27. Ald. 635. G. 288. in Montibus Cambrobrit.

Cervus, a Stag, I. 86. Ald. 774. Hinnuleus, a Hind. Hinnulus, a Fawn. Dama, a Doe. Dama vulg. a Fallow Deer, in vivariis quorum ingens copia per totam insulam.

Capreolus, the Roe-buck, I. 80 tab. 33. vulgo

habetur in Scotia.

A per, a Boar, I.108. Ald. 1006.1025. G. 918. Porcus, a Hog, Sus, a Sow, Scrofa, a Sow with Pig, Maralis, a Barrow-hog, I.ib.

Digitata Vivipara Semifera.

Vulpes, a Fox, I. 128. tab. 56. Ald. 221. G.

Herinaceus & Frinaceus, an Vrchin or Hedg-

M. 1.148. tab.68. In vepretis & fepibus.

Vitulus mar. a Seal, or Sea-Calf, I. 148. tab. 8. Ald. 724. ---- In borealibus tractibus & woia.

Lutra, an Otter, I. 148. tab. 68. Ald. 295. G. 684. in plerifq; fluviis majoribus, præsertim in Sussexia.

Mus domesticus, the house Monse, I 164 tab. 66. Ald. 417. G. 714. observavi, album, Cinerum & sufcum.

Mus Major seu Sorex, a Rat, I. ut supra, Ald.

406. G.731.

Mus aquaticus, a Water Rat, I. 156 tab. 73.
Mus Araneus, the Erdsbrew, or Field Monse,
1.164. tab. 66. Ald. 442. Invenitur nigri vel
issi coloris, & alter dorso nigro ventreq; albo,
& parvus griseus & in partibus borealibus magnus cauda brevi.

Mus Avellanarum, a Sleeper, or Dormonse, I. tiupra, Ald. 440. G. 735. raro occurrit.

Mustela, a Weasel, I. 155. tab 72. Musteasylv. Ald. 327. G. 752. In domibus, & horteis.

Viverra, 'chis, Furo, a Ferret, G. 762. at Sine kone.

Martes, a Martern, or Marteron, 1.155. Martis sceleton, Ald. 333. G 765. In agro Wiltomens & borealibus.

N 4

Putorius

anus 4 cus,

ng

ac-15.

tur G.

at,

88.

us,

ens Igo

18.

Putorius, a Fitchem, or Pole-cat, I, 155. tab.

64. Ald. 330. G. 767.

Taxus suillus, Meles, I. tab. 72. a Badger, Gray, Brock, or Bawson, I. ubi supra, Ald. 266. 267. G. 686. Cryptas incolit.

Lepus, a Hare, 1. 157. tab. 65. Ald. 351.G.

605.

Cuniculus, a Coney, I. 157. tab. 65. Ald. 385. G. 362. Estq; grifeus argenteus & niger.

Sciurus, a Squirrel, I. 164. tab. 66. Ald.

398. G. 845. In sylvis & arboretis.

Talpa, a Mole, Want, or Wont, I. 164. tab. 66. Ald. 451. G. 931. Repperitur quandoq; tota alba, nulla invenitur in Shepey insula,

Digitata domestica.

Canis leporarius & Scoticus, a Grey-hound,

I. 174. Ald. 545. G. 269.

Molossus, Canis bellicosus Anglicus, a Mafiff, Ald. 559. Canis, I. 174. tab. 79.

Canis fagax ad aquas, a Water-Spaniel, Ald.

556.

Canix Scoticus fagax, a Blood-hound, & alter furum deprehensor, Ald. 553. 554.

Canis sagax ad Coturnices capiendas, a Land-

Spaniel, Ald. 555.

Canis minor celer, a Beagle.

Canis Getulus, an Islana Dog, I. tab. 69.

Canis Melitensis, a Foisting Dog, Ald. 542. & alter brevioribus pilis, id. 541.

Canis pastoris fidelis, a Shepherds Car.

Canis Villaticus, a Cur Dog.

Canis Leporarius, a Hound, I. 174. tal., 70.

Canis pugnax, a Butchers Bull or Bear Dog. Canis avium venatica, a Setting Dog.

Canis focius & fidelis, a good House-keeping

Dog.

tab.

ger. 66.

.G.

Ald.

ger. ld.

ab.

eq;

nd,

la-

ld.

ter

nd-

12.

nis

Canis mixtus, a Mongrel.

Felis, Catus, a Cat, 1. 174. tab. 72. Ald. 57. Felis Sylv. a wild Cat, 1.174.tab.72.Ald.583.

Felis Syriaca multis punctis inter stinca, the Caprefs Cat, Ald. 556. Enutritur in adibus nobilium.

Digitata ovipara cute tecta.

Rana, a Frog, I. 187. tab. 75. Ald 591. eft vel Terrestris major flava, vel minor nigricans, secundo aquatica vulg. vel maculata.

Ranunculus viridis seu Dryopetes, G. 60.

Ranæ fætus caudatus, Tadpoles, G. de quadrup. oviparis, pag. 46.

Bufo, a Toad, 1.187. tab.75. Ald. 609. Rana Rubeta tum palustris, tum terrestris, G. ib.64.

Lacertus, an Eft, or Newt, I. 190. tab. 76. Ald. 627. G. 31. Hujus species apud nos, 1. Terrefris vulg ventre nigro maculato, 2. Terrefris anguiformis in Ericetis, 3. Parvus terrestris fufos oppido rarus, 4. Aquat. fuscus, 5. Aquat. niger, Lacertus Hibernicus ex Catal, Musai Tradescantiam.

#### Aves Britannica.

### Terrestres Carnivora.

Quila, the Eagle, I.10. tab. 1.2. Ald. 110. A G.149. quandoq; huc migrat ex Hibernia ubi abundat.

Accipiter, the Hawk, I. 20. tab. 7. Ald. 225. 228. G. 3.

Halixtus, the Sea Eagle, vel Ofprey Turn. Quandoq; conspicitur in Cornubia, I. 12.tab. 2.

Ald. 188.190. G. 177. fine icone.

Lanarius, the Lanar, mas vocatur, the Lanaret, Ald. 381.382. I.24. tab. 9 .-- in Shir wood Foreft, in agro Notinghamenfi, and in Dean Foreft, in agro Glaucestrensi.

Accip. Palumbarius, the Gofbank, mas dici-

tur the Taffel, Tertiolus, G.43.

Accip. Fringillarius, & Nisus, the Sparrow-Hank, I. 22. tab. 8. Ald. 346. 347. G.44. mas appellatur, the Muschel, In plerifq; locis fylvaticis.

Tinnunculus mas & fæmina, a Stannel, or Stonegall, I. 22. tab. 8. Ald 358. a Kefbrel, or Kaftrel, in tractibus Auftral. G. 46.

Falco, the Fant on, I. 30. tab. 12. speciem

nescio, in Pembrook-Sbire.

Coccyx, Cuculus, the Cuckoe, or Guckoe, I. 24. tab. 10. Ald. 414. 416. Sub medium Aprilis nos advolat.

Lanius, the Butcher, or murdering Bird, 1.24. tab. nb, 10. Ald 389. G. 520. vidi juxta Kingsland, astivo tempore ter. quaterve. Lanius Cinereus

Anglice, a Skreck. G.520.

Laniorum duas alias species observavit nobiis vir D. Willoughly totius naturæ diligentissimus Callentissimusq; scrutator non solum per Britanniam sed maximam partem Europæ.

Milvus, the forked tail'd Kite, I. 24. tab. 11. Ald. 395. G. 549. Turn. a Glede, a Put-

tock.

0.

1-

5.

n.

2.

1,

of,

i-

7-

25

1.

17

97

m

I.

is

Subuteo, the ring-tail d Kite, I. 24. tab. 9. Buteo Triorchis, the Buzzard, Ald. 367.

Peronos, the bald Buzzard, or Kite.

Noctua, the Night, or little grey Owl, I. 48. ab. 18. Ald. tom. 1. 544. Bubo Turn. a like Fowl.

Ulula, the white hooping Owl, or Owlet, or Hewlet, tab. 19. Ald. 538. G. 700.

Strix, the Screech, or screeching Owl. I. tab.

19. Ald. 563.

Corvus, the Raven, I. 38. tab 16. Ald. 694.

in ulmetis juxta ades nobilium, G. 294.

Corvus, J. 38. tab. 16. our common or Carnin Crow, G. 282. Cornix nigra, Ald. 736. & Cornix simpliciter Turn.

Cornix frugilega, spermologus, a Rook, I. 40.

tab. 17. Ald. tom. 1.753.

Cornix aquat. Hanc videt Turn. apud Morpetenses in ripis fluminum, G. 293. suspicor este, the mur Cornubiensium.

Cornix Cinerea, the Royfton Crow, I. ubi fu-

pra, Ald. 755.

Graculus

Graculus vel Monedula, a fackdaw, a Chongh, Turn. a Caddo, a Ka, I. 38. tab. 16. Ald. 771. G. 467.

Coracias Arist. the Cornish Chongh, I. 38, tab. 16. Ald. 768. In omnibus oris maritimis a Cor-

nubia ad Doroberniam.

Pica Glandaria, a 9 av, I. 40. tab. 17. Ald. 789. Garrulus avis, G. 634.

Pica, the Magpie, Pyot, Py-anet, I. 40. tab. 17. Pica varia seu caudata, Ald. tom. 1.85. G. 628.

Pica Marina, the Sea Pye, I. ut supra Ald.

tab. 792. 794.

Vespertilio, a Bat, Flittermouse, Rearmouse, I. 52. tab. 20. Ald. 574. G. 604. vesperi apparet astate. Hyeme vero latet in cryptis, & rupibus.

Loxias, the Shell-Apple, Ald. 2.877. I. 46. fine Icone in agro Warwice in Pomariis, Mr.

Willoughby.

Caprimulgus, the Goat-Sucker, I. 52. tab. 20. Ald. 568. G. 215. Hunc coepit Dominus Cole, in agro Hantoniensi, an. 1664. rara admodum avis.

Aves Granivora non canora.

Pavo, the Peacotk, I. 52. tab. 22. Ald. 219. G. 594.

Gallopavo, the Turkey cock, I. 58. tab. 24.

Ald. tab. 2. 39. G. 425.

Phasianus, the Pheasant, I. ut supra Ald. tom. 2. 49. G. 619. Horum pulli vocantur Pouts, Est albus & alter suscess.

Urogallus,

Urogallus, major Cock of the Wood, I. 60. tab. 15. Urogallus seu Tetrao major Ald. 2. 64. in Hibernia occurrit.

Gallina Coryllorum, the Hasel Hen, Grons, I. 60, tab. 25. Ald. 2. 82. Bonosa Albert, G. 203. Gall. Asricana, the Guiney Hen, I. 58. tab. 24. Gallina Guinea, Ald. tom. 2. 337. meleagris rel Gallus Numidicus, G. 424.

Otis, Tarda, Bistarda, the Bustard, I. 62. tab. 16. Ald. 288. G. 430. On Newmarket Heath,

& in Campestribus Sarisburiensibus.

Attagen, a Godwit, I. 62. tab. 26. Ald. 275.

Perdix Ruffa, the Partridge, I. 62. tab. 27. Ald. 2. 139. G. 606.

Coturnix, the Quail, I.62. tab. 27. Ald. tom.

2.153. G. 311.

gb.

71.

ab.

)r\_

ld.

b.

8.

d.

e,

1

.

.

Rusticola minor, the Snipe, or Snite, I. 62. ub. 27. Scolopax. Gallinago minor Ald. tom. 3. 479. Gallinago sive Rusticola minor, G. 448.est altera Hujus species nuncupata, the fack snipe.

Rusticola major, Scolopax. Gallinago, I.88. tab. 31. the Wood-cock, Ald. tom. 3. 473. Rusticola vel Perdix Rustica major, G. 445. Utraq;

Hyeme huc migrant. ex Hibernia.

Ralla-Anglor, the Rail, or King of the Quails,

Ald. 3. 455.

Upupa, the Hoopee, 1.62. tab. 27. Ald. 2. 704. G. 703. In the New Forest in Hampshire, & in Essexia, sed raro invenitur.

Pulveratrices domeffica.

Gallus, a Cock, I. 82. tab. 29. Ald. tom. 2.

... Hirfutis pedibus, ib.

Angl. Ald. 2. 341.

Gallina Rustica Turn. quam variis de causis

Attagenem esse conjicit.

Pulveratrices Lavatrices.

Fulica, a Coot, I. 88. tab. 31. Ald. 395. G.

Ispida, the Kings fisher, I. 88. tab. 31: Ald.

3.520. G 513.

Gallina Aq. I. 88. tab. 31.

Gallina ferica, I. 88. tab. 31. fic dicta à splen-

dore, Ald. 3. 470.

Columba vulg. Livia, the common House. Pidgeon, or Culver, I. 88. tab. 32. Aid. 2. 462. G. 245.

Columba Gutturosa perperam dicta Cropper,

Ald. 2. 479.

Columbæ Cypriæ, Jacobins, I. 88. tab. 32.

Ald. 2. 471.

Columbæ Turcicæ, Coloris sunt Betulini, cum oculis rubris, Ald. 481.

Columb. Tabellaria, Carriers.

Columb. Tremulæ, Shakers, funtq; vel acuti vel lati caudæ.

Columb. Hirsutis pedibus, rough-footed Pidgeons, I. 88. tab. 32. Ald 2. 466.

Columb. Angl. & Ruffica, G. 245. inter has majores vocantur, Runts.

Columb. Galeatæ, Helmets.

Columb.

Columb. maculis nigris & aliæ rubris decora-

Columb. Percustores, Smiters.

Columb. Gyratrices, Tumblers, Omnes hæ Columbæ in Columbariis aluntur præterq; has icuriosis educantur Turcicæ, Barbaricæ, the Finkin, Cornew, Bastard bill, Light Horsman, Dragoon.

Turtur, the Turtle Dove, 1.88 tab. 32. Ald.

1.509. G. 277.

2.

nna

usis

G.

ld.

n-

Ce-

2.

er,

2:

ni,

ıti

d-

as

Ь.

Palumbus major torquatus, a Ring Dove, or Quist, Ald. 2. 4. 7. In sylvis, Turn, a Comstot, tringed Dove.

Oenas seu vinago, a Stock Dove, or Wood. Pid-

geon, Ald. 2. 499. I. 88. tab. 3.2.

Passer domesticus, the House-Sparrow, I. 92. ab. 34. G. 581. Ald. 2. 534. idem quandoq; abus invenitur, I. ibid. Ald. 556.

Passer pusillus in Juglandibus degens, I. 96.

ine Icone, Ald. 2. 563.

Junco, the Reed Sparrow, I. 166. tab. 53. Hujus datur minor species in Arundinetis prope Kingstoniam.

Granivare Canore.

Carduelis, a Gold-finch, I. 69- tab. 36. Ald.

1.801. G. 215. Aurivittis Turn.

Calandra Ald. 2. 847. est Alaudæ persimilis sed ipsa paulo major, Ramis ar borum intidet an Passer torquatus G. a Bunting.

Coccothraustes, I. 98. tab. 37. Coccothrau-

fles mas Ald. 2. 846.

Fringilla, the Common, or Chaffinch, I. ib Ald. 2.817.

2, 817. G. 342. Turn. a Sheld apl. a Spink. Monti fringilla, the Bramble, or Brambling, I. 96. tab. 33. Ald. 2. 822. G. 343.

Chloris, the Green finch, I. ib. Ald. 2. 811.

G. 226. Turn. Acanthis, Spinus, Ligurinus. Citrinella, the Yellow hammer, I. 96. tab. 36, Ald. 2. 859. Emberiza flava Turn. a Youlring, G. 591.

Linaria, the Linet, I. 96 tab. 36, and 98 tab.

37. Ald. 2. 824. G. 550.

Luteola, a Siskin, rara apud Anglos avis nec uspiam fere alibi quam in caveis cernitur semel in agris Cantabrigianis se vidisse recordatur, Turn.

Alauda, the Lark, I. 98. tab. 37.38. Ald. 2.

845. G. 67. Turn. Galerita.

Alauda pratensis, the tit-Lark, I.ib. Ald. 2.840. Alauda cristata, the wood- Lark, G. 72. Ald.

2. 841. I. 98. tab. 37.

Rubicilla, a Bull finch, a Hoop, and Bul Spink, a Nope, I. 120. tab. 43. Pyrrhula five Rubicilla Ald. 2. 745. G. 662.

Baccivore.

Turdus vulg. the Song-Thrush, I. 98. tab 37.

Ald. 2. 600. & Turn. a Thruffel.

Turdus Viscovorus, the Misletoe Thrush, or Saith, I. 102. tab. 39. Ald.2. 583. G. 688.& Turn, simpliciter, a Thrush.

Turdus Illas, the Wind Thrush, I. 102. tab. 39. Turdus minor Ald. 2. 598. Turdus minor Illas vel Tylas, G. 689. Turn. a Wind Thrus.

Trichas, the Feldefare, Turdus pilaris Ald. 596. tantum untum hyeme apud nos reperitur.

Merula, Collyrion Turn. the Black-bird, or Hack Oufle, I. 104. 140. Ald. 2. 604. G. 543. Sturmus vulg the Stare, or Starling, I. 104. 1,40. Ald. 2. 632. G. 677. circa turres & alti-

ora ædificiorum culmina.

Sturnus Cinereus, Ald. 2. 638. I. 104. t. 40. Caruleo, a Clot bird, a Smatch, or Arling, a Stone-check, nidulatur in Cuniculorum foveis, & sub lapide in Anglia Turn.

### Insectivora.

Picus viridis, the Green Wood pecker, or Hickvall, I. 110. t. 41. Ald. 1. 849. G. 642. Chlonon, Virio, a Witwoll, Turn.

Picus varius major, I. 110. t. 41. Ald. 846. Picus varius minor & mas, I. ib. Ald. 1.

8474

ık.

no,

I.

36,

ng,

b.

nec nel

ur,

2.

10. Id.

nk,

illa

37.

07 . &

26.

100

96.

um

Picus murarius, the Creeper, or Wall-Creeper, lib Ald. 1. 852. G. 644.

Picus Cinereus, I. 110. t. 41. Sitta feu Picus

Cinereus Ald. 1.853.

Juynx, seu Torquilla, the Wryneck, I. 114. 1.42. Ald. 1. 866. G. 515.

Certhia, the Ox-eye Creeper, J. 114. t. 42.

Ald. 2. 870. Certhia Turn. G. 223.

Paffer Troglodytes, a Wren, 1. ib. Ald. 2. 61. G.588. Trochilus, Senator, Regulus Turn. Curruca, the Hedge Sparrow, G. 326. Ald. 1.753. Hypolais feu Curruca, 1. 122. t. 45.

Hirundo, the House Swallow, I. 114. t. 42. Ald. Ald. 2. 662. G. 492. vivit per Hyemem in mineris stanneis Cornubiensibus & in Rupibus marinis.

Hirundo Riparia, the Sand Martin, or Shorebird, J. ib. Ald. 2. 695. a bank Martnet, G. 508.

Hirundo agrestis five Rustica Plinii, a Mar-

tin, Ald. 2. 693. 1. 114.t. 42.

Hirundo apus, a black Martin, or Martlet, Ald. 2.699. I. 114. t. 42. a Rock or Church Martnet, G. 507.

Parus major, the Common Titmonse, I. 120.t. 43. Ald. 2. 713. G. 578. the great Titmonse,

vel the great Ox-eye Turn.

Parus Cæruleus minor, I. 122. t. 44. Ald. 721. G. 579. the less Titmoufe, Turn.

Parus ater, feu Carbonarius, the Coalmonfe,

I. ib. Ald. 2. 723 G. 579.

Parus Caudatus, the leaft, or long taild Titmouse, I. ib. Ald. 2. 716. G. 580.

Motacilla, a Water Wagtail, 1.122. t.44. Ald. 2. 727. G. 557. Culicilega, a Wag tail, Turn.

Motacilla flava rostro longiusculo nigricante,

1. 122. t. 44. Ald, 2. 859. G. 559.

Rubetra, the Stone-Chatter, or Blackberryeater, & Turn. mortetter, I. 122. t. 45. Ald. 2. 740. a Moortising Aquilonaribus.

Rubecula, the Ruddock, Red breast, and Robin Red-breaft, I. ib. Ald. 2. 742. G. 661.

Ruticilla, Phanicurus, the Red fart, I. 120. 43. Ald. 2. 747. a Redtail, G. 663. ex Turn.

Oenanthe, the Wheat ear, or White tail, 1.122,

t. 45. Ald. 2. 763. G. 567. in agro Warwicensi Fallow Smiters.

Luscinia, Lusciniola, the Nightingale, 1. ib.

Ald. 2. 777. G. 532.

Morinellus, the Dotterel, Ald. 3. 540. G. 554. in agro Lincolniensi certo anni tempore capitur jocose, vide Camden.

Aquatica Palmipides.

Cygnus, the Swan, I. 136. t. 48. Ald. 3: 8. G. 321.

Anser Domesticus, the Goose, mas vocatur, the Gander, I. 136. t. 48. Ald. 3. 102. G.125.

Chenalopex, vulpanser, a Bergander, nusquam slias vidi nisi in Thamesi sluvio aiunt tamen esse frequentem in insula Tenia (Thanet.) vocata & illic in scrobibus cuniculorum nidulari, Turn.

Anfer ferus, I. 136. t. 48. Ald. 3. 150. G.

140.

mibus

ore-

G.

Car-

let.

rch

). t.

use,

\ld.

use,

Tit-

Id.

m.

nte,

rry-

. 2.

Ro-

20.

122.

45.

Capricalca, Capricalze Scotis, Ald. 3. 164.

1,136. t. 48. G. 146.

Anseris speciem vidi in Cimelio Tradescanti fib nomine squeed una cum Ovo ex Insula Scotia Bass dicta, in qua quam plurima avium general stato anni tempore midificant uti etiam in insula Vecti.

Bernicla Brenta, the Brant Goofe, I. 136. t.

48. Ald. 3. 167. Brenchus, G. 95.

Gustarda Avis Scotica Ald. 3. 163. G. 145.

Anfer Baffanus, five Scoticus, a Soland Goofe, G. 145. ex infula Bafs non procul Edinburgo.

Anas Domefticus, the Duck, mas, the Drake, 1,142. t. 49. Ald. 2. 188. Anas cicur, G. 83.

O 2 Harle

Harle, the black Diver, I. 148. t. 49. a Shell Drake in Norfolcia.

Anas fera, I. ib. Ald. 3. 222. G. 101. the

Wild Duck.

Anas fera fusca, I. 142. t. 49. Ald. 3. 221. in Paludibus Lincolniensibus.

Anas Platyrhincus Ald. 231. in paludibus

Crowlandienfibus, I. 142. t. 49.

Querquedula, the Teal, I. 142. t. 49. Ald. 3.

549. G. 91.

A Gaddel, Ornithopolis nostris sic dictus est magnitudine Anatis, rostrum similimum rostro Querquedulæ, sed aliquanto magis cærulescit,

Penelope major, the Widgeon, Ald. 3. 219,

I. 142. t. 49.

Penelope fæm, Ald. 3. 220.

Columbus major, the great Ducker, I. 136.

t. 48. Ald. 3. 252.

Colymbus a Norwegis Lumme, a nostratibus, Razor bill, Worm, 304. ex Auctario Clus. pag. 367. Mr. Willoughby.

Colymbus Cristatus seu Auritus, Worm, ib.

fine Icone, idem.

Colymbus medius, the Dive-dapper, or Arsfort, I. 136. t. 45. Ald. 3. 258.

Colymbus minimus, the Dab Chick.

Mergorum serrati-Rostratorum species major & minor, in fluvio Tame in agro Warwicensi an. 1664. cum rigidissima suerit hyems, Mr. Willoughby.

Mergus Turn (ut fentio) qui vidit in rupibus marinis nidificantes, juxta Ostium Tinz fluvii in Norfolcia, Hoc me ditavit Doctiffimus affinis meus Ds. Jenner Sclopeto transfosso in agro Wiltoniensi.

Corvus aquat. the Cormorant, Ald. 3. 263. 1.t. 27. Carbo aquat. G. 121. in Cornubia Shags, Turn. mergus.

Onocrotalus, five Pelicanus, the Pelicane, I.

128. t. 46. Ald. 3. 47.

bell

the

11.

ous

3.

est

ro

t.

9.

6.

ıs,

g.

Ь.

1-

3-

n-

fr.

US

rii

in

Pelicanus sive Platza, a Shovelard, I. 128.

t, 46. ex agro Lincoln. Turn. a Spoon bill.

Larus major & minor albus, the Sea Mew, 1.126. t. 46. Ald, 3. 65. & simpliciter, Gul, Sea Gul, or Sea Cob.

Larus, quem Cornubienses indigitant, a Gant, forsan detorto nomine a Gavia vel a Gallicorum Gavian quod idem sonat, est par Anseri, palmipes, rostrum rotundum cæruleum, corpus grisei coloris, alte volat alausasq; minores solas captat.

Puphinus Anglicus, the Puphin, G. 657. ex Mona Linsula Anglesey, & Cornubia, Anas Artica Clusif & Fraterculus Ald. 3. 230. lin. 13, 14.

Fissipides Aquatica.

Ciconia, the Stork, I. 148. t. 50. Ald. 3.

111. G. 230. raro huc advolat.

Ardea Cineria, the Ash coloured Heron, or Hern, Hernsham, I. 148. t. 50. Ald. 3. 378. Ardea pulla sive Cinerea, G. 187.

Ardea alba, G. I. 152. t. 51. Ald. 390. G.

189. a Mire Drumble.

Ardea stellaris, the Bittourn, I. ib. Ald. 405. G. 100.

O 3 Ardea

Ardes minor, I. tab. 56. quam ad me transmist Ds. Jenner, ex agro Wiltoniensi.

Avis pugnax; I. 134. t.52. a Rough, eft tertia

in Tab.

Avis pugnax, quarta in dicta Tab. a Reev, utræq; ex agro Lincoln. est fæmina superior ris.

Hæmantopus mas & fæm. Red shanks, I. 154, t. 52.

Arquata, seu numenius, the Curliem, Ald. 3.

426. 1. 152. t. 51. G. 197.

Arquata congener, a Stone Curliem, huic rostrum breve, accipitrinum, pennæ milvi, Phasiano par magnitudine, Dilicatissimæ avis exagro Hantoniensi, Ds. Hutchinson Ornithopola Londinensis.

Vannellus, the Lapwing, bastard Plover, or Pewit, insula quadam ab its nomen sortitur in Essexia: Huc enim migrant practise ad diem Divo Georgio sacrum, vide Fuller, 318, I, 166. t. 53. Ald. 3. 526. G. 692.

Vannello congener capella vulgari proceior, at Turdo minor, pennis Caruleis & crista lon-

ga, ex Cornubia Ds. Gunthorp,

Pluvialis Cinerea, the Grey Plaver, I. 166,

t. 53. Ald. 3. 531. G. 647.

Pluvialis flavescens hujus meminit, J. pag.

Pluvialis vulg. the Whiftling Plover, or green

Plover, in Ericeto Lincoln.

Rallus Itallorum, I. p. 147. Ald. 3. 98. utribique fine icone.

Trynga

Trynga Ald. 3. 814. I. 166. t. 53. justa stage in Ichthyophylatica, & rivulorum margines, in agro Warwicensi, Mr. Willoughby.

Trynga paulo minor, Ald. 3. 482. 1, 166. t.

13. idem.

f-

ia

0.

4,

3.

ic

a-

a

7

ŋ

R

7

1

.

Ņ

Merulam aquat, vidit volantem in Comberlandia Ds. Willoughby, Ald. 3. 486. I. 166. t.

Charadrios ab incolis, Sea Lark, in Horibus Cambrobritannicis, præsertim in fatione Belli Mauritii, I. 166. t. 53. Charadrios five Hiatula, Ald. 3. 537. Ds. Willoughby.

Grus, the Crane, I. 166. t. 45. Ald. 3. 329.

G. 474.

Crex, a Daker Hen, est avis longis cruribus, catera coturnici, (nisi quod major sit) similis, qua in segete & Lino, vere, & in Principio and this non aliam habet vocem quam crex, hanc enim vocem semper ingeminat, quam ego Aristorecem esse puto, nusquam in Anglia nisi in sola Northumbria, vidi, & audivi, & an sit eadem am Ortygometra superius memorata nescio, suspicor tamen esse Turn. me talem vidisse audivisse ad wheatley quinq; Oxonio Milliaribus memini.

Tres sequentes aves cum nominibus & breviusculis descriptionibus mihi communicavit Ds. Hutchinson Ornithopola Lond. quas se vidisse ait in agro Lincoln.

Non est avis aquatica querquedula paulo minor Rostrum ei rotundum, tenue & augustum, superius paululum incurvatum, toto venti e albes-

Q 4

cit, dorsum nigrum, caput cristatum unde for.

fan ei nomen, Ic. a monacha velata.

Crickaleel, est priori æqualis, Cærulea in alarum supernis, caput collumq; maculata, adingluviem coloris grisei inde deorsum albescit vel contra quoad ventris colorem.

Gossander, palmipes & cristata ventre aureo, rostro longo & angusto, caro slavescit & coca tota facessit in oleum, non est edulis, exagro

Lincoln, videtur effe Puphini species.

# Pisces Britannici.

## Pelagii Squammofi.

A Sellus merluccius, Cabeliau, a Cod, G.84. Ald. 286. a Borealibus, a Keling, ab Occidentalibus, a Melwell, Turp. & hujus folliculus glutinosus Ichthyopolis, the Sound.

Afellus mollis, a Whiting, I. t. 2. qui in sole exiccatus nuncupatur, Buckthorn, & Scalpion Turn. omniscus sive Afellus & ad hunc alium

refert quem nominat, a Meerling.

Alellorum tertia species, a Haddock, G. 86. Ald. 282. Turn. in Epist. ad. Gesn. Onos sive

Afellus mollis nigricans, a Coal Whiting Chichestrensibus, & Pollock Cornubiensibus.

Afellus niger, the Coal Fish, Ald. 289. G.

89. Turn. Carbonarius minor.

Afellus sub mento barbatus, a Pout, Ichthyo-

Afelli

Aselli quartum genus, nunquam vidi in tota Anglia nisi in Northumbria prope oppidum, Bednel, a Codling vocatur, Haddocko est multo major, Goddo multo minor Turn.

Afellus fuscus Ichthyopolis Lond, a Hake.

Asellus varius molva vel morva, Ling a Longiudine, I. 226. t. 46. Ald. 288. G. 87. 88.

Harengus, a Herring, I. 8. t. 1. Ald. 189.G.

95.

for-

ala-

in-

vel

eo,

cta

c-

1-

le

n

m

5.

e

.

.

. 13

i

Alosa minor, a Pilchard, ingenti copia Quotannis capiuntur in tractu Cornubiensi & in Italiam Hispaniamq, vehuntur.

#### Pelagii Lav's.

Serpens marinus, a Sea Snake, I. 16. t. 4.

Conger, Congrus, a Conger Eel, I. 16. t. 4.

Ald. 349. G. 290. in Cornubia, Milwel.

Galeus, a Dog, or Dogger fish, 1. 26 t 8. Catulus Ald. 390 394. G. 167. 607. Hujus genus quoddam appellant Cornubienses, a Plusher, Carium.

Aquila piscis seu Pastinaca mar. the Poison fish, the Fire, or fierce Flam, the Cat-fish, I. 30. t. 9.

Ald. 426. G. 75. &c.

Raia, a Thorn-back, I. 36. t. 10. Ald. 450. G. 790. Raia junior vocatur, a Maid.

Raia Clavata, Ald. 459. G. 796.

Raia lævis a Homelyn, or White Cunt, a Scate, &in Cornubia, a Guilt head.

Raia exiccata & formata, the Dragen, Ald.

443.

443. Raia exiccata & concinnata ad formam monstrosi piscis volantis, I. t. 12.

Squatina, a Kingstone, I. 36. t. 11. Ald. 472.

G. 899.

wid George Rana piscatrix, the Monk, Toad, Nass, or what. Devil fish, 1.36. t. 11. Ald. 466. G. 813.

Lumpus Anglorum, a Lump, 1.42. t.13. Ald.

479. G. part. 2. 25.

Turdorum variæ species, I. 42. t. 13. Ald. 2. 16. in quamplurimis Cambrobritaniæ locis & prope insulam Anglesey, ubi appellantur, Sea Carps, Mr. Willoughby.

Acus, the Needle, or Garfish, and Horn-keck, I. 52. t. 15. Ald. 107. G. 9. in Cornubia, an

Old Wife.

Mugil imberbis, a Mullet, I. 68. t. 17. Ald.

131. G. 567.

Faber, a Doree, a Dorn, I. 60. t. 17. Ald. 112. G. 369.

Cuculus, a Gonrnet, a grey Gournet, I. 60. t.

17. Ald. 739.

Lyra, a Rotchet, I. 114. t. 24. Ald. 146. G. 516. in Cornubia, an Illeck, a Red Gournet.

Lyra speciem Cornubienses vocant, a Tub, a capitis sig. majori, depressiori, breviori, cui caro pinguior & friabilior.

Apuia vera, a Dace, I. 74. t. 19. Ald. 211.

G. 64.

Sarda, a Sprat, I. 82 t. 20. Ald. 221. Merulæ meminit Turn. quam se vidisse purat

in

inlittore occiduo quam Piscatores illic voca-

Solea, a Sole, I. 82. t. 20. Ald. 235. G.

667.

is

,

18

1.

d.

t.

3.

,4

cui

11.

rat

in

Citharus, a Dab, I. 82, t. 20. Ald 240. G. 127. Hunc vidi prona parte album, supina suf-

Gandils Angl. I. 92. t. 21. Santals, Ald. 254.
G. part. 2. 5. Sand Eels, quafi Arenose Angu-

Araneus, a Weaver, or Wiver, I. 92. t. 21. Sorpius Ald. 96. 256. G. 77.

Scomber, Scombrus, a Mackrell, I. 92. t. 21.
Ald. 270. G. 842. a Cornubiensibus nuncupa-

or Bucks Horn.

Paffer niger a Flounder or Flool

Paffer niger, a Flounder, or Flook, Fletuletus Ald. 243. 249. Fleffus fluviatilis Bellon.

Passer aureus, a Plais, Passer lavis Ald. 243.
Rhombus lavis, G. 663.

Passer Britannicus, a Holyburt, Holybut, or Holybut, Ald. 243. G. 666. sine I cone Fletule-ms Bellon, ex Cornubia missus erat R hombo 3. no major quinq; pedes longus, cum triplici denum serie, cauda longissima, prona parte Galum, supina soleam quoad colorem referebat.

Rhombus, a Turbut, 1.98.t. 22. Ald. 250.

G. 670.

Rhomboides, a Pril. Ichthyopolis Lond. I t. 20. Passer asper sive squammosus Rondel, Ald. 42. Limanda, G. 665.

## Squammosi in Mari & in Fluminibus.

Salmo, a Salmon, I. 106. t. 23. Ald. 483. G. 825.

Salmoneta, a Salmon Trout, parvus Salmo, G. 828.

Acipenfer, Sturio, a Sturgeon, I. 106. t. 23.

Ald. 526. G. 6.

Acipenseri congener, cui valde similis excepto capite, saporis delicatissimi captus erat in Insula Vecti anno 1664. Ds. Cole, qui ipsum delineavit exiccavitque.

Experlanus, Viola, Smelta, a Smelt, I. 114.

t. 24. Ald. 536. G. 365.

Lampetra, a Lamprey, I. 114. t. 24. Ald. 540 mustela seu Lampetra, G. 590.

Lampetra fluviat, a Lampern, Ald 581.G.598. Lampetra parva fluviatilis Rond, 202. Here-

fordienfibus, a Stone Grig.

Anguilla, an Eel, I. 114. t. 24. Ald. 4.

G. 41.
Congri pufilli, Elvers, in Sabrina non procul
Glevo Urbe, G. 292.

### Fluviatiles Squammofi.

Trutta, a Trout, I. 130. t. 26. Ald. 589. G. 1.007. minorem Truttæ speciem Cornubienses vocant, a Shote.

Truttæ genus in Northumbria vocamus, a Bull Trout, Truttam taurinam, ab infigni magnitudine, in aliis provinciis vocatur, a Grey Trout,

paliis, a Skurf. Turn. in Epist. ad Gesn. meminit & ibidem alterius Truttæ Allerfange dictæ and profundius fub alnis dilitescat.

Truttæ species Londinum affertur, ex agro Lincolniensi, quam Shard vocant, Ds. Wine.

khthyopola regius,

.G.

mo,

23.

cep-

t in

de-

14.

Ald.

198.

ere-

14.

ocul

.G.

rout. in

Truttæ species quam nostrates Scag vocant, a Thamisi non procul Kingstonio.

Umbra fluviatilis, an Umber, I. 130. t. 26. Ald 596.

Barba & Barbus, a Barble, I, ut supra Ald.

198. G. 124.

Capito, Cephalus fluv. a Chub, a Pollard, a Chevin. I. 130. t. 26. Ald. 601. G. 396. & partt 2. 8.

Leuciscus, a Dar, or Dart, I. 130. t. 26. Ald.

607. G .26.

Gobio mar. fluviatili viridior, & maculis maresperus apud Cumbrenses, qui vocant, a fundlin, Turn.

Gobio & Gobius fluviatilis, a Gudgeon, or

links, I. 130. t. 26. Ald. 612. G. 399.

Gobites barbatus, a Loch, I. 30. t. 26. Ald. 18. fine Icone.

Asper pisciculus, the broader Sharpling, I. 10. t. 26.

Asper pisciculus Gobioni persimilis Rond.

enses 14.615. in the North.

Pungitius, the Prickling, Banfted, I. 146. t. s, 4 & Ald. 628. Pisciculus aculeatus, G. 8. in sof-

Cernua

Cernua fluviatilis, a Ruff, G. 192. propter horum piscium copiam Celebratur fluvius Yare non procul Norwico in Essexia.

Rutilus feu Rubellus, the Roch, I. 130. t. 26.

Ald. 62 1. 732. G. 821.

Gobio capitatus, Cottus Rond. a Millers thamb, a Cull, I. 130. t. 26. Ald. 614. G.400.

Fluviatiles Laves.

Alofa feu Clupea, a Shad, I. 146. t. 27. Ald. 501. G. 19.

Phoxinus, a Minoe, Minom, I. 146. t. 27.

28. Ald. 582. Mæna. G. 715.

Cyprinus, a Carp, I. 146. t. 27.29. Ald.

637. G. 310.

Cyprinus latus, Abramus, a Bream, or Barn alicubi I. 146 t. 29 Ald. 642. qui si crassior suerit a Cornubiensibus vocatur a Bass.

Mustela, an Eelpout, I. 146. t. 28. Ald. 290. 577. G. 599. Occidentalibus, a Coney-fish.

Perca, the Perch, I. 146. t. 29. Ald. 622. G.

699.

Lucius, a Pike, I. 146. t. 29. Ald. 625.G. 501. maximos vocat Gesner. Luces, parvos Pikrels, & adultos Luls.

Alburnus, a Bleik, or Bleis, I. 146. t.29. Ald.

629. G. 23.

Tinca, a Tench, I. 146. t. 29. Ald. 646.G. 984.

Cetacei Generis.

Balena, the Whale, I. 216. t. 41. Ald. 676.
G. 114.

Cetus

Cetus Britannicus, G. 212, I. 216, t. 41.

Cetacei generis piscis quem nostrates Grampus vocant nullibi extat icon. (quod sciam) caput refert illud quod venit sub nomine Delphini

apud I. t. 44. G. 321.

cr

re

6.

rį

O.

d.

7.

d.

rn

or

10

G.

G.

ck

ld.

G.

16.

tus

Porcus marinus (quem nautæ Harengivorum nuncuparunt) littus nostrum appulit præterita hieme anno 1642, una cum fatulactante. In diffecti exterioribus apparuit pinguedo ad 7. vel 8. pollicum craffitiem ex qua oleum elicuerunt rustici & carnem sale Condierunt & comederunt. Hu jus sapor carni bovillæ non multum cedebat, uti etiam adeps. Totus piscis muccaginis ingenti copia oppletus erat, Ds, Gunthorp.

Maculo Cetus Anglicus, G. 216.

Phocana, a Porpus, a Porpes, 1.216. t. 41. Ald, 720. Turfio, G. 712. Bell 16.

### Molles feu Mollufii.

Loligo, Sepia, the Cutle, Prete, I. 4.t. I. Ald.

49. &c. G. 825.

Stella mar, minor, the Sea Star, Rofes, Prizers S. Fingers, I.t. 8. & Sea pad, G. 919. Invenitur 3. bus. vel quatuor radiis in oftio The mifis.

Polypus, J. 438. Ald. 14. G. 739. ab incolis Vectis Infulæ, a poor Cutle, Turn.

# Crustacei.

Loculta mar. malè Ichthyopolis nostris, a long gd sm Offer, I. 14. t 2. Ald. 102. G. 485.

Aftacus mar, the Lobfter, I. 14. t.2. Ald. 112.

G. 98.

G. 98. recens capti (antequam coquantur cum fale underubescunt) funt nigri cærulei, & quan. doq; flavescentis coloris.

Aftacus fluviatilis, Gammarus, the Crey fif,

1. 18. t. 3. & 20. t. 4. Ald. 129. G. 104.

Fagurus decem libras pendens, a Punger, G.

155.

Squilla parva, the Prawn, I. 20. t. 4. Squilla parva Rond. Ald. 155. G. 907. squilla fluv. à nostratibus vocatur, a Shrimp.

Squilla fluviatilis, Rond, 212, in fluviis rece-

dente æstu & in rivulis.

Cancer marinus, a Crab, I. 424. t. 5. Ald.

174. G. 148.

Cancellus crusta superna faciem humanam referens cum duabus antennis palmaribus, Dr. Ball Collegii Medd. Lond. Socius honorarius dignissimus.

Cancer parvus, I. 36. t. 7. Cancer fluv. Ald.

207. G. 138.

Cancer in tellis degens, I. 20. t. 7. Wrong beirs.

Araneus mar the Sea Spider, I. 16.t. 6, Ald. 201.

Echinus mar. the Sea Urchin, or Sea Hedgebog , I 48. t. 12. Ald. 403.

Echinus spinis spoliatus, the Sea Button, Ald.

403. G. 350.

Echino Spatago nudo in partibus maritimis Cambrobrit. Meermaids heads, & obsceno nomine Cants, Imperat. pag. 781. Hunc mihi dedit Dr. Ball, in tractibus occidentalibus ab ipfo Testacei captum.

Teftacei.

Ostrea, the Oyster, I. 56. t. 14. Ald. 482. G. 645. Distinguint Ichyopolæ nostri in rupicas soleæ equinæ æquales & minores inter quas prædellunt Colcestrienses & Wainsteetenses, ex his quædam in stagnis quæ mare singulisæstibus inundant, aluntur & inde pinguiores viridioresq; sunt.

Peden, a Scollop, I. 56. t. 14. Ald. 500. G.

690.

à

.

١.

n

S

1.

g

1

.

١.

15

).

6

ei

Patellæ Limpyn & Lympets, Ald. 546. I. t.

16.

Pectunculi, Cook fift, for fan Turn. I. 56. t. 14. pectunculi auricularum forma variantes.

Lepades faxis adharentes, Cockles, G. 688.

elt Patellæ Species.

Mytuli, Muscles, I. 66. t. 15. Ald, 509. G. 277. majores à nostratibus appellantur, Horse Muscles.

Cochlea mar. Whilks, or Periminkles, Ald.

389.

Cochleæ fluviatiles , G. 214.

Balani five Polycipides, Center Shells, I.66.t.

15. Ald. 5 23. G. 121.

Concha Venerea minores, I. 68. t. 17.

Solen, the Sheath, or Razor fift, I. 66. t. 15.

Ald. 528. G. 1035.

Turbinum minorum 3 vel 4 species, I. 46. t.
11. Turbinis tuberosi Icones Rond. Ald. 353.

Trochorum parvulorum duz species, 1.46.t.

P

Tubili

Tibuli in quibus vermes, Worm-shells, I.68. tab. 17. Ald 562. G. 1026. in Ostreis antiquis.

Concha ex dimidia pene margine profunde

ftriata.

Conchæ Anatiferæ, Barnacles, Ald. 544.Divulsi a navibus ex India Orientali reducibus.

Zoophyta.

Tethyæ Thætyæ I. 74. 19. Ald. 576. 583. G. 954. Hæ fæpiuscule adnascuntur antiquis ostrearum testis.

Urticæ vel Pulmonis marini speciei meminit Ds. Gunthorp, litteris ad me datis ex Cornubia his verbis, Blubber, est Gelatina quædam lucida, variis mixta coloribus, rubro, viridi, cæruleo, albo, est globosa subtus cavaq; in cujus medio apparet quadrilatera, quasi quis digitos pollicemq; contraheret, denuoq; aperiret & hoc modo nutrimentum suum apprehendit: Magnitudo ejus quanta vola manus continere potest, exiccata ad nihilum sere redigitur. Aquis innatat, & huc illucq; à ventis, æstuq; desertur, certisq; tantum anni temporibus conspicitur, semel quæsivi in loco duabus à littore leucis ubi prius milliones videram neq; unicum inveni, Maio mense apparet.

Meminit Camdenus in sua Britannia quatuot piscium sequentium, Guini ad in agro Mertonethensi in palude Bala sexaginta ingereum,

Jerosch Lithostrata, 2. Torcoch in Caernarvenstin palude truba comp Limparey, 3. Chare in Wimander Meer quosinsed srubger; colæ etiamnum affirmant illic reperiri sed certis rubic under commen sonat – casden oio audur; tantum ul nomen Cambrica sonat – casden oio audur; tantum charry diet. in

Agr. Lincostr.

tantum anni temporibus, & alterius unoculi meminit in Montibus Caernarvens, capti quod mihi dubiæ est sidei.

Et vir ornatissimus Ds. Carem in libro suo de Cornubia hos enumerat, a Barn, a Scad, a Shad, Cudles, a Bass, a Millet, a Horpool, Dorns, Polleck, Illeck, Cunner, Rockling, Wroth, Becket, Squarrey, Rough-hound, (quem suspicor esse Galeum vulgarem) qui qualesve suerint ignoro atq; etiam hos Chichestrenssum, Bland, Pulver, Bresen, Sea rats, & Farthing-Trout.

Insetta nostratia primum in genere sub suis

daffibus postea in specie proposita.

le

j-

3.

uis

nit

bia

ıci-

ru-

ne-

itos

hoc

lag-

test,

nna-

cer-

emel

orius

Maio

tuor

erto-

eum,

alude

os in-

certis

ntum

Terrestria favisicantia alata, pedata, Anelytra, quadripennia pennis vel membranaceis instructa, viz. Apes, sucus, Bombylus vespa crabro, Tipula que non favisicat vel pennis farinaceis, viz. Papilio qui diurnus, vel nocturnus, uterq; maximus, medius, & minimus, Bombyx,

Anelytra bipennia, Muscæ.

Coleoptera seu vaginipennia. Locusta, Grylalus, Gryllotalpa, Scarabæus, Proscarabæus, Capricornus, Cantharides, Cicindela, Forficula.

Terrestria pedata non alata sex pedam, formica, Cimices, pediculus, Syrones, Ricinus, Pulex, Lendes, vel osto pedam, Araneus, vel multipeda, viz. Erucæ, Circulio, Ambulo, Scolopendra, Julus, Asellus.

Terrestria apoda sunt vermes 1. Plantarum, 2. Fruticarii, 3. Leguminarii, 4. Herbivori, 5. Animalium, 6. in Homine, 7. Limax & Cochleæ.

Aquatica paucipeda. Squilla, pediculus & pulex mar. P 2 Aquatica Aquatica multipedi, Scolopendra.

Aquatica apeda. Hirudo, Stella, Lumbricus mar. Seta, vermes in navibus, tubulis, tellis tellaceorum, Teredines.

Favificant.

Apis, a Bee, or Been, I. 4. t. 1. Ald. 191. Apiarium, I. 25. t. 2. inter has Regulus. Dantur sylvestres minores, nigriores.

Fucus, a Drone, or Dran, I. 4. t. 1. Ald. 191.

hu jus variatur color ut in apibus.

Bombylus, a Humming, or Humble-Bee, I. 4. t. 1. M. 53. Horum nidus, I. 25. t. 2. Repperitur major minorq; & alter nigricans, cum

pedibus flavescentibus.

Vespa, a Wasp, I. 4 t. 1. M. 42. Ald. 203. Vespeta Ald. 209. &c. I. 25. t. 2. Habemus flavam & nigricantem utrumq; majorem & minorem, & alteram longiorem, & insuper pseudo sphecas aculeo carentes degentes in parietibus ex limo constructis.

Crabo, Tenthredo, a Hornet, or great Wasp, I. 4. t. 1. Mouf. 50. Ald. 226. Crabroneta,

300. &c. I. 25. t. 2.

Libellæ, Tipulæ, perlæ, Brises, or Breeses, Adder bolts, Dragon styes, and Water Butterstyes, I. 41. t. 4. 138. t. 27. Ald. 303. ad 306. M. 66. ad 71. quarum species.

I. Virescens cum maculis in collo, summilg;

lateribus ex fusco viridibus.

2. Priori fimilis sed colore obscuriore.

3. Oculis margaritis similibus, alis argenteis macula susca simbriatis.

4. Lutea

Lutea cum 4. vel 5. spinulis in caudz extremitate.

5. Corpore & capite cœsiis, ore magno, cum
3. spinulis in cauda.

6. Cum Macula nigra in medio alarum & 2. spinulis in cauda.

7. Rubea capite collo & scapulis luteis.

cus

ftis

91.

an-

10

I.

ep-

um

3.

nus

&

per pa-

sp,

eta,

fes,

06.

ifq;

teis

itea

8. Brevis toto corpore lutea, & cauda quasi abscissa intersegetes.

Media, 1. Capite Cæruleo aliis violaceis.

2. Cafia cum alis albescentibus, casiis lineis in longum ductis, medioq; purpurascentibus.

3. Subviridis cum alis striatim sanguineis, ver-

4. Fusca cum viriditate fatiscente.

5. Viridis cum oculis cæruleis.

6. Viridis ex toto, sed in alis leviter.

Minima, I. Cyanea alis argenteis.

2. Rubra alis argenteis.

3. Crocea, cauda protuberante & fanguineo colore maculata.

4. Minima, cauda exili longa, capite magno.

Lacustres, 1. Cauda bifurcata, corpore lon-

2. Lutea, caudà bifurcata.

3. Fusca longis alis cruribus, antennisq; Juxta paludes & stagna. Tipulæ ortum ex pediculo marino Clarè demonstravit mihi, D. Ball.

Papilio, I. 46. t. 5. & 56: t. 6.7.8. Ald. 237

ad. 253. M. 89 ad. 108. eftq; vel diurnus, a But. terfly, vel nocturnus, a Miller, qui phalana di-

citur cujus species sequuntur.

Phalæna major, 2. exigua argentea nigris maculis notata, a Mosh, 3. minor cum aliis externis nigris, quinq; maculis internis rubris, 4. minor gracilis cum quatuor fanguineis maculis in aliis exterioribus, 5. minor cum guttis fanguineis duabus circa extremitatem alarum, 6. major cum exterioribus alis fuscis internis aureis nigra linea fimbriatis

Papilio diurnus maximus, maximam partem flavescens, reliquam pene nigricans, 2. Cum antennis porrectioribus oculisq; nigerrimis.

1. Papilio diurnus medius ex pallido flave-

2. Cum corpore & alis externis nigrioribus, fuscis hic illic maculis notatis.

3. Cum spinosis crenis suscus.

4. Dorso nigro, ventre thoraceq; albis.

5. Corpore & antennis livescentibus, capite alisq; pallidis.

6. Oculo nigro, pupilla candida, alifq; Dra-

contii modo variis,

7. Alis æreis micantibus venulis, & fimbria maculis nigris conspersa.

8. Antennis ex luteo rubescentibus, & alarum

limbis multicoloribus.

9. Capite, alisq, lacteis quibus maculæ susca, & nigricantes.

10. Ultima parte alæ exterioris clypeolo ni-

gro, quem medium punctus eburneus ornat, deorata.

11. Alis arctiffiffimis, cauda lata & quali plu-

12. Corpore Cinereo, caudaq; nigricante.

Papiliones minimi, 1. Cum alis externis coc-

2. Alis cæsiis, & æruginosis, intus oculatis.

3. Pedibus purpureis, & corpore fuscis punfulis pleno.

4. Alis oculatis cyanum cælestem spiranti-

bus.

utdi-

ris

·X-

4.

n-

6.

Q.

m

n-

e.

S,

e

è

2

n

- 5. Corpore gruino, in medio alarum luteo Cineritia.
- 6. Corpore, pedibus, capitulo, antennis Cineritiis.

7. Summe nigra, primifq; alis deargentatis.

8. Alis omnibus pallescente flavo nitenti-

9. Alis conchatis, & clavatis, ex albo & obfurè rubente variis.

Muscæ, Flies, a volando vid. iconas apud, I. 68. t. 9. I. 70. t. 10. I. t. 19. 142. Ald. 346. ad 349. 471. &c. M. 58. ad 67. Harum species sunt.

1. Carnivora in macellis usitata, the Flish sty, 2. Hippoboscus, a side or Horse sty, 2. Tabanides, a Burrel sty, Stout, Brees, Clog, or Cling. 4. Culiciformis crassifuscula ex rubro stavescente. 5. Culiciform. Capite longo, productis pernigrisq; cruribus inter volandum pendulis, posteriora

riora crura anterioribus multo productiora.

6. Curvicauda, briftle-tail'd, with two hairs, or more.

7. Scorpiurus.

8. Bipilicauda.

9. Tota nigra, pedibus ex croceo flavescentibus.

10. Pedo, Gruina, the Crane-sty.

11. Pedo sæmina, Shepheards.

12. Stercoraria.

13. Humipeta, a gray Path-sty, pro piscibus capiendis.

14. Intermalvas & urticas degens.

15. Bibio vel Vinula, the Wine-sty.

16. Herbivora max.

17. Nocturna minima viridescens examinibus volunt inordinate.

Eulæ, Maggots, Eularum ova, Flyblows.

18. Musca ex phryganio orta pro truttis alli-

ciendis, the Cados fly.

19. Musca in Calicibus herbæ Ben reperta.

20. Musca nocturna minima viridis alis porrectissimis ex foll. fraxini.

21. Musea priori similis nisi quod susci sit co-

loris ex foll, aceris majoris.

22. Culex, the Gnat, qui si minor Midges dicitur.

23. Culices pyramidales duo in Anglia Aquilonari, Monff.

24. Culex ex pomis Quercinis ortus, idem.

Locusta, a Grashopper, cujus datur apud nos major & minor ubiq; in pratis pinguioribus.

Gryllus, a Cricket, I. 78. t. 12. Ald. 443. M.

124. in Caminis & Pistrinis.

Gryllotalpa, Evochur, Fen. Cricket, & Churworm, I. 78. t. 12. 88. t. 13. M. 164. In hortis & stercoretis.

Scarabzus, a Beetle, Bug, or Clock, 1. 90.

t 15. & 94. t. 14. & 100. t. 16. Ald. 450. ad

461. M. 148. ad 164. 327. 328. 330.

Scarabæus cervus volans dictus, a flag, or flying Fly, M. 148. I. 94. t. 14. Ald. 451. In agro Hantoniensi.

Capricornus, a Goat Chafer, M. 149. I. 90.

t. 15. Ald. 452.

Scarabæus stercorarius vel fimarius, a dung Beetle, or Sharnbug, M. 153.

Smaragdus vel viridulus, a green Chafer. M.

160.

ra.

or

ota

De-

na,

ter

la,

ır-

r-

ly-

lli-

ı.

or-

:0-

ges

ui-

105

M.

ur-

or-

10.

5.

Scarab. arboreus Dors, M. 160, in lignis aridis habitat.

Hydrocantharus major Anglicus vel nostras, M. 327. Water Clocks.

Scarab. Pilularius melanocyaneus, ubiq; oc-

currit in agro Colchestrensi.

Scarabæus minor ruber maculis nigris notatus, a Lady cow, M. 161. Hujus aliam speciem observari paulo majorem sine maculis, & alteram ex rubro nigroq, pulchrè tessulatam.

Scarab. minor nitore splendens, & alter Ca-

ruleus Cyaneusve...

Proscarabæus, an Oil-beetle, or Oil-clock, M.

Cantharides ex cicutaria, M 145.

worm, viz. fæminæ quæ vermes sunt, mares autem sunt alati, nec lucent, Mouff. 109.

mogeiti aliarumq, plantarum infidet, M. 137.

Bruchus, a Field Cricket.

Blatte

Blatta fatida, Londini in Cellis vinariis, M.

139.

Blattam observavi depascentem solia ulmorum novellorum relictis sibris quæ elegantissime dissecta erant.

forden Robin Ballock, M. 171. Datur vulgari major & corpore nigricantiore.

Formica, an Ant, Emet, or Pismire, I. 114.t. 17. Ald. 517. Inter has aliæ pennatæ, aliæ im-

plumes.

Cimices domestici impennes, Wall-lice, Punies, M. 269. I. 118. t. 3. Ald. 542. In parietibus,&

lectorum spondis.

Pediculus, a Louse, M. 259. in capite simpliciter, a Louse. 2. Morpiones, Crab er peirt lice, 3.

Corporei maculati, Body lice.

Pediculus marinus, Ald. 712. I. 188. tab. 14.
Pediculi ovium seu Ricim vel potius Reduvii, Ticks, Sheep-lice, M. 272. & etiam in aliis animalibus sibi peculiares generantur, viz. in Equis, bobus, vitulis, porcis utii dicti. Canibus, Capris, Cervis. In Salmonibus, Accipitribus, Gallinis, Columbis, Anseribus, Coturnicibus, Phasianis, Perdicibus, &c. In plantis, nempe Abrotano, phynthio Nymphææ floribus, & potissimum Aquilegiæ soliis, fructu pruni majusculi.

Pulex, a Flea, Videfis iconem in Microgra-

phia Di. Hook.

Afellus mar. fquillam refert, M.322. Hos confertissimos vidi in statione navium Billingsgate dicta cymbam ascensurus post æstus restuxum.

Pulex

Pulex aquaticus, minutissimum animal in agro intra S. Jameses & Mary-bone, juxta aquadudum.

Lendes, Nits, sunt pediculorum ova.

Asellus, a Sow, Tylers lonse, Thurs-lonse, Cheese-bug, Cheese-lip, M. 203. I. 107. tab. 23.

Afellus arvensis Julio hunc in agro Colcefrensi vidit Pennius inter segetes, alias nunquam

repperit, M. 247. Hexapus eft.

50

do

10

n-

14

CX

Araneus, a Spider, M. 234. I. 130. t. 18. Ald. 608. Estq; Domesticus vulgaris, 2. Fuscus non nullæ pelluciditatis, 3. Retiarius inter rosarum mmulos rete artificiociffimum expandens, vel sylvestris, 1. Sepiarius crassiorem telam texens. adum in agro Colcestrensi observavit Pennius inter origanum sylvestre muscis insidiantem, 3. Opilio, a Shepheard, 4. niger breviusculis pedibus ovum nigrum fub ventre portans, 5. nigernmus magno corpore pedibus breviffimis in putridis & cavis arboribus degit, 6 oblongus cauda unta, 7. Araneola ruberrima boves mordens, Blain worms, Twings, & in Borealibus, Tings, que inter buprellas numeratur, a Monf, 141.9. Lupus maximus fuscus globosus, 10. Lupus medus tinctura grisea, 11. Lupus nigrior in dorso Cruciger. Lupus non inepte vocari potest, the leaping or dancing Spider, 12. Araneus parvus luteus ventre nigro ex Rosis Damascenis, Jun 13. 1666. - mido de Lyster - Thely. Gransach. Eruca, Caterpillers generali nomine, hirfutas

Erucæ, Caterpillers generali nomine, hirfutas vocant Boreales Oubuts, Meridionales, Palmerworms, M. 179. I. 142. 144 t. 19. 20. 21. 22. Ald. 166, &c.

rides sunt, 1. Ligustrina nobilissima. 2. Sambucina priori similis. 3. Lactucaria. 4. Mespilaria.

5. Quercina ha omnes decempeda.

Erucæ flavescentes, 1. Tiliaria. 2. Salicina. 3. Porcellus in trifol. paludoso. His adde sphondylen seu verticillum, Borealibus Andever, Meridionalibus, Whurl-worm nuncupatam, M. 196. I. 114. t. 17.

Staphylinus, the poysonous Caterpillar, M. 197.

I. 114. t. 17. cujus duas species proponit Monff.

Prima tota cum splendore nigricat, corpore gracili & longo. Secunda Caput parvum puniceo suscum fere globosum, os sorcipatum & exiguum habet.

albo viridis in incessu gibbosa. 3. Nigerrima pedes caudamq;. 4. Costis miniacei Coloris. 5. Populnea ex Ochræ colore & albo. 6. Striata alba ex slavo. 7. Undulata Cinerea ex nigricante. 8. Ranunculata, tota varia. 9. Glauco-nigrz.

Erucæ hirsuæ densius pilatæ sunt, 1. Ambulones, Palmer-worms, & Bear-worms. 2. Flava collo ventreq; piloss. 3. e Fusco slava cum capite puniceo surca invea ornato. 4. Flava ex susco dorso pilisq; dorsi, ventre ejusq; pilis utrisq; suscis, 5. Corpore vario pusillis guttulis sub cæruleis decorata. 6. Lucophæa setis valde sertaceis, rigidisq; 7. Peniculata cum cunei formibus in dorso maculis 8. Murina cum pilis in dorso cuneatim dispositis. 9. Singulis incisuris variis ab invicem coloribus. 1 c. Ornata liris nigris, Cæruleis, viridibus

vi.

u-

ia.

3.

y-

ri-

6.

7.

f.

a-

eo

m

Ex

na

is.

ta

n-

.53

u.

va

ni-

co q;

æ-

e-

in

e-

·i-

15.

US

viridibus & luteis. 11. Cum in cifuris albicantibus & prastinis. 12. Corylaria saturate viridis. 13. Corylaria pallidius virescens, 14. Mesoleuca. 15. Urticaria cum pilis erectis rigidiusculis. 16. Brafficaria capite cyaneo. 17. Sepiaria major crocescente vultu. 18. Sepiaria minor cum facie subcarulea. 19. Antennulata. 20. Jacobaica capite, pedibufq; obsoletis. 21. Echinata caruleo virens. 22. Echinata mesoleuca. 23. Rubicola nigrocineritia liris utring; tribus pallide flavefcentibus, 24. Verbascina, 25. Lupulina. 26. Scrophularina. 27. Alnina. 28 Ulmina. 29. Tithymalina. 30. Glycyrrhizina. 31. Salicina. 32. His adde Bombycinam cum suo papilione & singulis metamorpholibus quas vide belle depictas apud the first M. 181. 1. 153. t. 22. & 171. t. 23. Ald. 280. 282. - pighio sed hæ Erucæ non nisi à curiosis aluntur.

Ad Erucas spectant Aureliæ, quæ vel vellere cooperiuntur, vel glabræ sunt, utræq, varii coloris, & nunc auro illitæ (quæ verè Chrysalides) nunc vero (quæ spuriæ) aureo colore vacuæ.

Scolopendra, Many feet, & Scolopender, M.

199. L. 171. t. 23. Ald. 636. 714.

Scolopendra nocticula, in Ericetis muscosis toto corpore lucens.

Scolopendra mar, alia grifei alia rubicundi co-

loris in offrearum areis. M 322.

Julus, I. 171. t. 23. a Gally worm, est brevis scolopendra cujus species Lactucarius glaber capite nigerrimo dorso slavo. 2. Totus niger, alba linea excepta à capite ad caudam usq; recta ducta per dorsum. 3. Obsolete slavescens cum capite pedibusq; rubris.

Juli-

Julius Hirsutus, albicans cum pilis brevissimis innatis nigris, 2. ventre sublivido, dorsóq; inamæna slavedine maculato, 3. Exilis spadiceus pe-

dibus exiguis & numerofis.

Vermes in homine, M. 249. I. 175. tab. 24. & 194. tab. 25. lumbrici simpliciter, Gut worms.

2. Cucurbitini, Tæniæ, Flat-worms, 3. Ascarides. 4. Syrones in cute, Wheal-worms, in sacie Barons nominantur, inventi etiam in Cerebro pulmonibus vesica (quos excretos non semel vidi) in auribus dente Lingua, liene, fellis vesicula, renibus, utero, sub unguibus in panaritio, Acarus, a Hand-worm.

In aliis animalibus sc. in Equis, bobus (quos bots appellant) vitulis, canibus, porcis, Salmonibus in quorum intestinis hi soli sine Chylo repperiuntur, in hepate ovium quos Lanii Flooks vocant. In radice cornuum Cervorum, Caprarum, ovium. Lytta sub lingua Canum rabidorum. In mullis, Gobiis sluviatilibus, leucisco, Ostrearum testis.

Vermes Terrestres majores, Earth-Worms, 2. Minoros rubri pro piscatoribus. 3. Sublividi. 4. Cauda flava, Yellow tails. 5. Torquati & pingues. 6. Minores appellantur, Meds & Grubs.

Arborei ligniperdes. 1. Deces in truncis crefcentibus. 2. Enxyla in aridis & exuccis. 3. Thripes in folidis ficcis. 4. Teredines Coss. 7. Timberworms, in cæsa materia. 5. Termites in lignis calidioribus, Teredines in navium tabulis. 6. Corticarii. 7. In omni frutice & suffrutice.

In fructibus, viz. Sorbis, Mespilis, pyris, Malis, Glande, Gallis, Nucibus, In

In plantis, Carlina, Pimpinella, Lupulo, Afphodelo, Violis, Perficaria filiquofa, Dipfaci capite, Rofarum pomis & cæteri herbivori, leguminarii vel fruticarii dicti.

Syrones in cera, & in caseo Mites. In frumentis nondum mola confractis & malta seu byna Curculiones, Wivels, Popes, 7. frumentarii, Meal-worms, Blatta, Tinea, a Book worm.

Vermes minimi rubri aquam stagnalem colore sanguineo inficientes unde vulgus dira porten-

dit.

e

0

i-

1-

,

15

j.

5

1-

).

li.

1-

2-

rnis

s,

Vermes sin aceto quos non sine arte semper custodit ornatissimus Collega meus Dr. Goddard verus naturæ mystes, similes sunt lumbricis Aquaticis, I. 194. t. 25.

Hirudo, sanguisuga, a Horse-leech, a Bloodsucker, estq; salutaris vel noxia, M. 323. Ald.

733. I. 194. t. 25.

Limax, a Snail without a Shell, the Honfe-

Snail, Ald. 703. I. 175. t.24.

Cochlea alba major cum suo operculo, sua-

vissimi est saporis in Sussexia.

Cochlea alba minor, ubiq, in hortis.

Cochlea vulgaris testa variegata.

Seta aquat. M. 325. Ald. 765. vulgo creditur oriri ex seta caudæ equinæ aquis immersa.

#### Serpentes.

Nguis, Coluber, a Snake, G.69. I.24.t.5. Ald. 270. Vipera, a Viper, G.125.1.10. t.1. Ald.115.

116.

frecat of

N.C.

Typhlops, Cacilia, a blind-worm, G.61. Ald. 245. I. 20. t. 4.

Metalla.

Aurum eruitur ex stanno at parva quantitate, Argentum ex minera in Cambrobritannia erusitur etiam ex minera stannea majore quantitate

m Cumbria & ex plumbea sed minore. Camden.n'

Stannum ex mineris Cornubiensibus & Devonia occidentali & in totam Europam distrahitur ntpote optimæ notæ metallum, stannurgi di-Ringuunt in Pyran & Murdick & Block Tyn,

& lapides stannei nuncupantur, Shoad.

Cuprum effossum erat Wenlochiæ in agro Salopiensi & Rich. 2di, tempore dives minera Cuprea Richmondiz in Episcopatu Dunelmens, & insuper Keswici & Newlandia in Cumbria, & minore copia in Cornubia, Edw. 1. & 2di tem-poribus. Im Walled - sat copies

Ferrum abundat in quamplurimis Angliæ locis utpote in Suffexia, in faltu Dean. dicto in agro Glocestrensi, in Cantio, juxta Bromicham in agro Warwicensi, uti etiam in Sommersetensi,

Stafforde Hantoniensi, Dunelmensi, Salopiensi, & Sheffeldiæ in Eboracensi, Distinguunt ferrarii in Cold-

fel, & Redfel, Scotia vocatur Slag.

Plum-

Plumbum in collibus Mendipensibus in agro sommertensi, Combertoni in Devonia, ad originem fluvii Istwyd in Cardigania, Richmondia, ad Abbatiam fontium in agro Eboracensi præcipue autem Wensdaliæ & Mashæ, optimum autem effoditur & maxima copia Brassenmoræ & Dovegangæ in agro Derbienfi, ex plumbo fiunt-Lithargyrus, Ceruffa, Minium.

Antimonium repperitur in mineris plumbeis

Derbiensibus.

Mineralis quædam gleba ex effusa aqua calorem igneum acquirens, cu jus meminit immortalis Boylæus in lib. de frigore.

Diaphani. Adamantes quos vocant Bristoll-Hones ex rupibus Divi Vincentii non procul dicta Urbe, ex mineris stanneis Cornubiensibus & rupe quadam inde the Diamond Rock, denominata, ex tractu meridionali Cambrobritannia.

Achates, the Agate, ex Cornubia uti referunt

quidam.

Ł

80

1-

a-

in

fi,

el-

m-

Crystallum effodi in agro Derbiensi scribunt inflores fed credo effe folummodo fluores.

Fluores, Spars, varii apud nos inveniuntur, ex agro Cantabrigiensi sexangulares & in aliis locis multum differunt quoad colorem, figuram, & pelluciditatem, Sequetes fervo in penu meo quos metallurgi Derbienses hisce nominibus distingunt in candidum, purpureum, Amethystinum leu violaceum. Dog (1) caninum, Tooth Spar dentatum, Aquaticum & rubrum. Quidam immixta continent plumbum, cuprum, stannum varivariosq; alios lapides præsertim Marcasitas. Hyacinthi impuri ex agro Eboracensi. Smaragdus ignobilis ex agro Eboracensi.

Ex animalibus. Alectorius in Stomacho Gallorum inventus.

Bufonius, The Toad-stone, falso sic dictus, demonstravi enim coram serenissima Ma jestate regia (in Collegio nostro qui interesse dignatus est exquisitissimis prælectionibus ornatissimi Collegia Dris Georgii Ent quem titulo Equitis aurati merito honoravit) hunc esse dentes molares Piscis Lupi, quem vide apud Sombenck felt & Jonst. de Pisc. t. 47. Et Aurisices summa cum admitatione confessi sunt hosces molares esse veros lapides busonios ab iis divenditos. Gesn. in appendice proponit sub nomine Aparrhica.

Ex animalium felle, in suspensa vidi multos seminibus Coriandri saccharo incrustatis similimos, in Vesica etiam Ovilla quem à seipso inventum mihi ostendit Dr. Hamey, ex intestinis, renibus, Pulmonibus, Cerebro, & aliis partibus. In vetula suspensa vidi diaphanum quendam, durissimum, ex crure extractum, & in senibus non semel observavi magnam partem Aorta petre-

factam.

Carpionum lapis. Cancrorum oculi male sic nominati, eximuntur enim ex Cammarorum capitibus certis anni temporibus. Limachum, qui videntur nil aliud esse quam horum animalium Cranium-Chelidonius in Hirundinibus. Margaritæ frequentes satis in quibusdam Ostreis prafertim fertim Scoticis, majores dicuntur extrahi ex Mitylis grandioribus ad Kerby Lansale in Welfmorlandia, Cornubia, Cumbria, & agro Salop. 7
Percarum inveniuntur in Capitibus omnium
Asellorum nostrorum supra memoratorum.

on Walling

mouse

de Thug 6.

Opaci. Alablastrum vel Alablastrites in agris Staffordiensi, Derbiensi & Lincolniensi ad insulam Axholm.

Ammites vel Hammites Sand-Stone, ubiq; obvius variorum colorum & qualitatum, qui pro
adificiis infervit appellatur Free-stone, & nomen Late qualitatur à latomia unde esfoditur, enigea, Purbeck, facile sciss
stone, ex quo construitur Templum Divi Pauli - iliz hic Londini, ex insula Purbeck.

Arena alba, lutea, Littoralis, rubra, omnes

crassiores vel minores.

1-

e-

e.

ft:

e.

u.

a-

&

d-

05

p-

05

li-

n-

e-

In

if-

fe-

re-

fic

ca-

qui

um

ga-

ra-

im

Astroites, the Star-stone Ald. 880. Boodt 300. Shugburiæ in agro Warwicensi & variis Angliæ locis, Lutescentem & cæruleum inveni in loco unde argilla essoditur ejusdem coloris Winchtombiæ in agro Glocestrensi.

Belemnitæ variorum colorum, quarum quædam femidiaphanæ aliæ magis opacæ. Vulgo

vocantur, the Thunder-bolt, Ald. 621.

Corallium feu potius Pseudo corallium quandoq; invenitur in littoribus nostris, maxime Cornubiensbus.

Cotes, novaculæ, Whet stones, & coloris nigricantis, & aliæ flavescentes ex Hibernia afferuntur, & agro Derbiensi.

Calaminaris, Calamie, ex quibus as conflatur

ex

ex collibus mendipensibus supra memoratis, ex hoc lapide Cadmia, botrytes, placites, &c. de quibus vide Math. in Diosc.

Lithoxyla. In agro Staffordiensi ager qui cito convertit quodcunq; lignum in lapides & variis

locis in Hibernia.

Magnes bonæ notæ in rupibus Dartmore in

Devenia, & alibi viliores.

Manganese ex collibus mendipensibus de quo plura in observationibus meis in artem vitrariam nuper editis. Magneria - of providuo. Magneria.

Marcasitæ variæ ex quibus Vitriolum conficitur, feu pyrites Copperas fones, & Fire fones, in toto littore Orientali nostro & plurimis locis terrestribus. Variant quamplurimum quoad colorem, Sc. Aureum, Argenteum aneum, ferreum, & quoad figuras innumeras quarum plures iconas exhibuerunt CB. in historia Balnei Bollensis Ald. 574. &c. cubica forma exhibent tracus nostri boreales, ænei plane coloris.

Marmor in agro Somertensi. In Scotia & Hiberniæ variis locis, in infula Purbech ad Egerton Com Sales in Episcopatu Dunelm. & in agro Hereford.

multiplic Marmoris maculati species in agro Ebora-multiplic censi. cham in calcary; suryon, monty in Agri ourning and Molares, Grinde, or Grindle stones, ex insula

Anglesey & Flint-Shire, & in agris Derbiensi,

Celtriensi, & Staffordiensi.

Sciffiles & cruftofi, Slates or Tyles, funt carulei, & coloris folii falviæ in Cornubia, grisei seu vulgares ad Colly Weston in Northamptonia &

in Cotes. Wold, optim; propo Bangor nignicales rulig - m Wallia - team m Com. Montgomr. Silex, Silex, the Flint. stone, nigricantes albi & quandoq, gemmas Diaphani etate & duritie imitantes, & alii margam continentes Ald. 214. Rusescentes sunt matrix pseudo adamantum Cambd.

Silex Dactylites ad littus Cantianum Ald.

730.

X

le

0

is

in

0

m

i-

in

is

)-

1,

is

S

-

71

a

u

Smiris, Emerie, or Emeril-stone in insula Sara

vel Guarnsey.

Syrtes seu sabulum vivum, Quick-sands, or Cocker-sands in Lancastria & in Cantio ad Vias regias.

Triorchis Viridescens hunc invenit Ds. Marmaduke Rawdon Essexiensis ad Whitby in agro

Eboracensi Ald. 484.

Præter superiores habemus Aluminosos varios in agro Eboracensi & aureos dictos ex aureo clivo Heresordiensi.

Calcarius ubiq; optimus censetur ad Barrow

in agro Licest.

Clivosi dicti Coloris grisei marmoris, aliiq;

nigri, instar gagatis in Cornubia.

Cylindrici, filicei intus extraq; ad Badmington in agro Gloucest.

Ceraunias Ald.: 611. & Ombrias id 616.

quandoq; fed raro apud nos invenitur.

Gypsum præstantissimum ex agro Darbiensi quo utuntur ad domorum areas & Concamerationes.

Hibernici, Irish-slate, pro inscribendis rebus memoriæ causa. Maculosi Punctis sanguineis ad sontem Sanctæ Winisredæin Flints. Moor-sones (1) Paludosi pro ædificiis in Cornubia,

Q3 Me-

Metallici testulati. In agro Richmond. Porus & Tophus ubivis pene occurrunt.

Ruber ad margines fluvii Erwell in agro Lancastriensi & in sundo sontis Sanca Winisrida, & ad Penrith in Cumberlandia.

Sabulum Pebles, Campos nostros nimis

infestat.

Sepulcrales duri Cærulei ad Pucklichurch Gloucestr.

Arietis cornu similis inter Bristoliam & Aquas Solis.

Saxum est pluribus calculis à natura productam ad Highgate montem Ascensuro.

Tartarum ex doliis vinariis.

Ex Fodinis plumbeis Dar biensibus missi erant sequentes sub hisce nominibus, Cank & Bastard Cank, Black Chert, Wheat-stone, Sheaf.

Lapides pro pavimentis platzarum, Rance, Rochester-stone. Silices, Ragman, quibus utuntur

pro platæis Lond.

Lapis spongiosus albissimus & lævissimus ex littore Essexiano.

Lapis schistus Cornubiensis diu calorem retinens, the Warming stone.

In Lapides Corversa. Sunt Cochleæ vulgares & Turbinatæ, Pectines, Ostrearum tellæ & Mytilorum ad Badmington in agro Gloucestr.

Bucardites primus Ald. p. 479. Hunc ad me transmist filius meus Robertus natu maximus Oxonio quem illic invenerat.

Echinites Ald. 455. 10, noster est plane silex albus creta obductus. GlosGlossopetra, a Tongne-frone, Boodt. 341. Ald. don't char 602. &c. funt dentata & sine dentibus in Variis Deany

matricibus latent ex mineris Carbonariis.

Ophiomorphites cum & sine aculeis, Snakefone, Ald. 540. ad Keynsham & Adderley juxta
terræ superficiem in agro Gloucestr. & ad Farnham in Surreia sed alterius coloris & ad Whithy
in agro Eborac. Ds. Marmaduke Ramdon mihi
Ophiomorphitem, Ostendit quem in horto suo
servat cu jus Diamiter 21. pollices æquat, Ex
agro Eboracensi. Ex varia constat materia.

Stelechites varii, quosdam cum nodis quales in arboribus apparent, alios sine his, quosdam radices plantarum referentes, alios animalium caudas, cæterasq; partes ex mineris Cambrobritann. attulit Regiæ Societatis ingens lumen Robertus Moray Eques auratus, vid. Ald. pag. 867. qui hisce auxit locupletissimum Cimelium

dictæ Societatis.

Stalagmites varii Dropping-stones, ex sonte Knaesborough in agro Eboracensi, vidi stramina, & integrum panem & Caseum (qui forsan Tyromorphus Ald.) muscum & plurima alia in hoc sonte petrificata, & ad puteum dropping-well appellatum, & in Bughan in Scotia.

Fungitem possideo elegantissimum ex agro

Eborac.

Ostra accomorphites Ald. 464. ad Bidford

Northampt.

Spatagoidem ostendit mihi Dr. Ball ex silice exemptum, in quo piscis modulum consecerat sux siguræ, & magnitudini conformem. In hoc Q 4 lapide

Am only

lapide comparuerunt singula lineamenta spatagi nudi, unde & ex aliis quamplurimis liquido mihi constat multos lapides pro naturalibus habitos ex animalibus eorumve partibus mediante succo quodam terrestri essistos, vel hac siguram suam Communicasse argilla vel terra cuivis molli, & postea periisse servata sigura, ut in lapidibus Islebianis liquido patet, in quibus sedulo observatori squamma occurrunt manifesta.

Hisce accenseri possunt Amygdaloides, Umbilicus marinus Eboracensis Mus. Trad. Agrisol. solia petrificata ex lacu Hibernico Loughneath, ib. Sideroites ex agro Cantabrigiensi & ad Sapmorth juxta Sharston in agro Wilton. Cochlea & Turbines in Norsolcia. De Testis piscium affirmare haud dubitarim nullas ex supra memoratis non alicubi apud nos inveniri petrificatas.

Lapis caudam felinam vel alterius animalis minoris referens ex plurimis articulis constat ad cujus articulationem plurimæ in ambitu striæ Ex Holy Island. Incolæ vocant St. Cutbeards beads Ds. Ramden.

Concha oblonga crassaq; coloris Lyncurii ex agro Wigorniensi ubi vocant Crow-stones & Crow-eups, & à forma oblonga Egg-stones, ex quibus autem conchis nostratibus proveniant, conjecturâ assequi haud facile est.

Ardentes. Carbo fossilis Pit-coal. Ex Novicastro, Tinmoutha, Episcopatu Dunelmensi, Insula Coquet dicta.

Lithantrax, Stone-coal in agris Derbiensi No-

ringhamensi in borealibus Leicestrens. Salop. stafford Pembrock. Caermarth. Sommers. ubi ænei sæpe apparent Coloris, referunt quidam mineram Carbonariam quondam accensam adhuc ardere ad Penneths-Chase in agro Salop. The proportion of the Agriffag

Lapis Obsidianus (in fallor) Kennel coal ex finel quo varia artificia fiunt. in Ag. Lancol Vitafforel.

Bitumen, Pitchfordiæ (1) Piceum vadum in agro Salop ubi fons perpetuo Bitumine opperitur, atq; etiam in Scotia non procul Edinburgo & Dunelmensi Episcop.

Cespites bituminosi Turfs, præsertim in agro Lancastriensi ubi Incolæ utuntur vice Cande-

larum.

i

Ô

n

IS

1.

1.

-

æ f-

)-

is

d

æ

X &

X

,

ala

0-

g-

Gagates, I et. In Norfolcia & prope Mulgravii castrum in agro Eborac.

Salia. Alumen nativum Cantabrig. Eborac. & Cantianum ex penu Tradesc.

Alumen factitium in agro Eboracensi.

Vitriolum copiosissimum in quamplurimis lo-

Nitrum ubivis & Aphronitum in cellis vina-

Sal marinus coquitur variis in locis, Camaloduni in Essexia, Fossilis albus & ruber, in Musao Tradesc.

Sal ex puteali aqua coctus albissimus, vid. fontes postea.

Lapides ex Snowdon monte altissimo, ex quibus

Ellicitur Alumen, Vitriolum, nitrum.

Terra. Fabrilis Rubrica, Red O. hre, Marking. frone, Redle & Rudle in agro Herefordienti.

Hantoniensi, Lancast, Rutland.

Nigrica fabrilis, Black ledd, Peculiaris hae terra Angliæ Europææ & Americanæ & hactenus nomine caruit, ideoq; favente Analogia hoc ipsi imposui ad Keswick in Cumbria.

Ochra, Tellow Ochre, yellow Stuff, Hereford. In agro Sommers. Oxon. & Scotia, denominan-

tur, à locis unde petuntur.

Azutum sive cæruleum fossile in Scotia

Umbra, Umbre ex Bristolia.

Lapis cæruleus Killow dictus ducendis lineis idoneus in agro Lancast. Figlinas habemus varias pro diversis Artificiis pro ollis vittariis & Metallis fundendis. Optimæ pro tubulis Nicotianæis ex Nonfuch in Surria ad Pool & ad insulam Vectim in Dorsettia.

Marga, Marle, observavi ruffam, griseam & luteam ex eodem loco. Lutea nobilis ad Brotheton in agro Eborac. & præstantissima in Suffolcia.

Smectis seu terra Fullonica, Fullers Earth, Fx Nutley in Sussexia & Woburn Berks, inde denominatur Woburn earth.

Rutlandsbire nomen sortitur à terra rubra lanas ovium inficiente, eadem invenitur in valle dicta de Red Horse Warwick.

Plastica, Creta, Chalk, ex qua album Hispanicum, & plurimæ pinguiores ubiq; locorum procurandæ,

Argilla

Argilla, Clay, pro lateribus conficiendis, & aliis ufibus. Invenitur alba, Cinerea, Cærulea, flava, rubra, &c. in agro Salop.

Agricolarum terræ plurimum variant in

fingulis regionibus.

c

G

I.

1-

12

is

15

.

is

.

8

ef-

le

a le

-

m

2

Marina. Succinum, Amber, raro occurrit. Meminit tamen Hettor Boeth, in Hist. Scot. ingentis massa equo majoris inventa in borealibus Scotia tractibus.

Ambragrisea, Ambergrise, nonnunquam ventis & æstu ad littus nostrum appellit, idq; quan-

dog: ingenti mole, & multis in locis.

Meteororum nomina ideo adjeci ut lector in fynopsi haberet quicquid in toto naturæ ambitu contineretur, & hic ansam, tradendi plurima quæ apud nos rariora sunt, prehendi, unde extranei & viatores curiosi occasionem capiant penitius examinandi, non quod vera hæc omnia esse asseriment, (excerpsi enim pleraq; ex scriptoribus nostris præsertim Camdeno) sed ut pro re nata viri harum rerum studiosi facilius in veritatem inquirerent.

Sequentur Meteora, ignita, viz Ignis fatuus, the Walking fire, or Jack of the Lantern. Castor & Pollux, Helena. Ignis lambens. Draco, Stella cadens: Est substantia quadam alba & glutinosa plurimis in locis conspicua quam nostrates star faln nuncupant, creduntq; multi originem suam debere stella cadenti hujusq; materiam esse esed Regia Societati palam ostendi solummodo oriri ex intestinis ranarum à corvis in unum locum congestis, quod aliis

aliis etiam ejufdem focietatis viri prastantisimi

postea confirmarunt.

Fulmen, Fulgur & Coruscatio, lapis sulmineus ex sulmine salso oriri creditus. Procellæ, Capra, Trabs, Bolis, Faces Dolia, Clypei, stellæ & milites discurrentes, Pyramis Galaxia quam tantum esse stellarum minutissimarum congeriem Telescopia plane detexerunt. Cometæ, Venti & hinc tetræ Motus.

Meteora aquea, Ros, Mel, Manna nostra ex ulmi foliis. Nubes Pruina, nix quarum generationem experimentis demonstravi in tractatulo de congelatione, Grando, nebula, Pluvia & fanguinis quam certissime constat esse tantum insectorum excrementa, Pluvia Tritici quam nibil aliud esse quam Hedera baccisera grana à sturnis devorata excretaq; comparanti liquidissime patet. Iris, Halo sive Corona, Parelii Paraselene, & Virga, spectra & phasmata. Maria nobis Oceanus Britannicus, Hibernicus Caledonicus Atlanticus & Germanicus.

Fontes, calidi ad Aquas Solis in agro Sommerfets, & alter ex clivis Divi Vincentii prope Briftoliam fingulis æstibus demersus, Buxtonæ in agro Darbiensi octo ex rupe erumpentes intra

feptem ulnarum spatium.

Fontes Purgantes ad Ebbelham in Surreia, ad Lewsham in Cantio, ad Barnet in Midlesexia, Spadani in agro Eboracensi prope Burntwoodin Essexia, non procul Stanes in Bercheria, ad Hailweston in Cornubia, ad Willinborough in Northamptonia Aquæ cujusdam deceni ab Exonia

onia milliaribus dissitæ meminit Ds. Gunthorp litteris ad me datis quæ ex ipso sonte haustæ blande purgat, at inde deportata ad aliquot milliaria hanc virtutem unà cum sapore & odore amittit. Hujus farinæ plurimi alii apud nos.

15

a-&

e-

X

a•

lo

1-

e-

il

r-

ic

200

in

a

d

a,

'n

d

n

-

Diuretici sunt Tunbrigenses & Sussexiensis & Astrofic Eboracenses, & ad Newnham regis in agroWarwicensi cui si sal admisceatur laxat ventrem si ferr saccharum sistit alvum.

Petrificantes, Knaesburienses in agro Eborac. ad Egerton prope Lenham in Cantia, & ad Abbatiam de Boxley juxta Maidstone, ad Kings-mill ex adverso Collegii Magdalensis Oxonii, ad Lutter-worth in Leicestria, ad Eldenhole in agro Darbiensi, ad Newnham Regis in agro Warwic. vid. Ger. emac. 1587. ad Boughton Northampt.cranium Fuller, 250. ad Shelly non procul Chiping Unger in Essexia, his addo puteum Dropping well-dictum in agro Eboracensi, & Wockey hole non procul à Bristolia, ad Tenderdon in Essexia intra 5. vel 6. annos lapis maniseste augetur, ab aqua incidente.

Salinus prope Harwich in Essexia & Lemingtonia in agro Warwic. Ex sequentibus Sal conficitur viz. ex sontibus ad Droitwich in agro Wigorn. ad Nantwich, Northwich, & Middlewich in Cestriensi, & sal niger vulgo Bay salt excoquitur in agro Hantoniensi.

Aqua Exotice natura, viz. Aqua fluvii Dee non miscetur cum palude de Bala in agro Merioneth, fluvius Levenney transit lacum I ynsava- ad sene vera ut q hodanu lacum than than lemanum plans net miscerij

than fine mixtura in agro Brecknock. Aqua in calefcit hyeme & frigida est æstate ad Luckington Hancocks in Bercheria. Vorago juxta Petriburgum adeo profunda & tam frigida ut nullus urinator fundum attingere poterit, ex quo ebullit fons nunquam congelascens. In aqua ad Saltas in Cornub, nunquam coquuntur pifa ad mollitiem ufitatam.

Rivuli fubterranei funt, Mole (1) talpa in Surreia. Deverel in Wiltonia per milliaris spatium. Recal juxta Elmley in agro Eborac. Lid. prope Lidfordin Devonia, Alen in Denbienfi, Medweri in Cantio ramulus prope Cockshead, Hans fluviolus in agro Stafford. Tribus milliaribus.

Tub terra cuniculos agit.

Æftus, Puteus ad Peak Foreft, non procul Buxtona in agro Derbiens. fluit refluitq; quater finnon Kitkongulis horis, & alius juxta Kilken in Flints. Staselhoc. fain tutis temporibus, ad Careg castrum fons observat æstus marinos, in agro Caermar, & in Cornubia lapis cavus in quo aqua contenta æstus ma-Ace Autor aut rinos imitatur. In portu Pool Devoniens. 4ter de die. Bristoliæ æstus ascendit ad 11. vel 12. ulnarum altitudinem. Ad Giglefwick in agroEborac. Scaturigines parvæ erumpunt, jadu disciab invicem haud distantes, quarum media sola singulis horæ quadrantibus afcendit ad novem pollices in refluxu autem unciam non superat. Juxta flumen Loder in Westmorlandia Cepius de die. Puteus ad flumen Ogmore prope Newton inagro Glamorgan distitus à Sabrina plus minus mille paffus aflate fingulis aftibus exurgit ad magnam altir

hunc-Hans

skitudinem. Observandi veniant æstus in Sabrina & Humbro quos nostrates. Hygras vocant. Eagers

П

i-

it B

r.

1.

e

gi ns

Z-

n.

a-

r-

r.

a-

er

2.

0-

b

nolta e.

ro lle m Catarrhactarum duorum meminerunt scri-pronuscial ptores nostri ad Kan prope Kendal in Nor-in Frent vinumbria, quorum si medius stridule magis, & Outo anti altius sonat certo expectant incolæ serenitatem qua apsio at Australis eodem sono presagit nebulas & plu-pim suant viam. Narravit mihi Ds. Williams doctissimus fuanti surisperitus se cum aliquot sociis judicem in itinerario suo Cambrobritann. comitantem observasse valle sereno aere in cacumine montis nigerrimam parvamq; nubem & inde pluviam. Qui miratus hoc ascendit montem viditq; nubem pluviamq; ortas ex Catarrhacta aquam in minimas guttulas dividente ex casu impetuoso. Simile quid alicubi resert incomparabilis Boylæus sibi contigisse (ni fallor) in Alpibus.

Sed ut telam retexam stemq; promissi in principio operis recensenda mihi loca celebriora in quibus arbores subterraneæ inveniuntur, viz. in Cornubia ad sinem terræ in agris Flints, Pem-Camdon brochiensi ubi ad littus maris tam consertæ cernuntur ut totum littus tanquam sylva cædua apparet, in Cestriensi, Cumbria & insula Anglesey & plerisq; locis Euro-borealibus. Vocant

Indicena, Noah's Ark.

FINIS.

A Ld. Aldrovandus.
A C C. Catalogus Cantabrigiensis.
C Capp. ejuldem appendix.
C O. Catalog. Oxon.
C B. Caspar Bauhinus.
I B. Ioann. Bauhin.
G. Gerardus emaculatus.
G. Gesnerus.
I. Ionstonus.
M. Moussettus.
P. Parkionus.
Pin. Pinax C B.
Turn. Turnerus.



